

EXCELLENT BROADLEAF WEED CONTROL IN PEANUTS.

PRODUCT ATTRIBUTES	BENEFITS
Reliable PPO Mode of Action	Aim® EC herbicide is a fast-acting PPO product for control of broadleaf weeds in peanuts.
Broad Spectrum of Activity	Aim EC herbicide performs under a wide range of diverse crop and environmental conditions.
PPO Chemistry	Different mode of action to increase efficacy and help with weed resistance management.
Crop Use, Application Flexibility	Aim EC herbicide can be applied preplant burndown, as a hooded spray and as a harvest aid.

TECHNICAL INFORMATION

Active Ingredient	Group	Formulation	Signal Word	Packaging
Carfentrazone-ethyl	WSSA Group 14	Emulsifiable concentrate	CAUTION	0.25 GA x 4, 6, or 12/case

APPLICATION RECOMMENDATIONS

- Aim EC herbicide can be applied as a burndown treatment no later than one day after planting up to 2 fl. oz./A.
- Aim EC herbicide can be applied to row middles using hooded sprayers up to 2 fl. oz./A.
- Aim EC herbicide can be applied to peanuts as a harvest aid to defoliate and/or desiccate troublesome broadleaf weeds such as morning glory and pigweed at 1-2 fl. oz./A.
- Add crop oil concentrate (COC) at 1 gal. per 100 gal. of spray solution, methylated seed oil (MSO), or nonionic surfactant (NIS) at 0.25% v/v. Coverage is essential for maximum efficacy on target weeds.
- **Maximum Total Seasonal Rate:** 6.1 fl. oz./A per season.
- **PHI:** Seven days
- Use at least 15 GPA for optimum results. The higher rate (2 fl. oz./A) may be needed if smallflower morningglory is present.
- Do not graze or feed peanut hay to livestock. Only one application of harvest aid per season is permitted.
- Rainfast in six to eight hours.

TECHNICAL INFORMATION

- Active Ingredient: Carfentrazone-ethyl

1 FL. OZ. AIM HERBICIDE WITH NIS CORDELE, GA

