
## **AUTHORITY® ASSIST** HERBICIDE BROADLEAF CONTROL WITH GRASS ASSIST

Group 2,14 Herbicide

APPLICATION GUIDELINES			
Application Timing	Apply preplant up to three days after planting.		
Use Rates	6 – 12 fl. oz./A. See label for specific soil, OM use rate recommendations.		
	Use high rates for soils of pH less than 7.0 and lower rates for pH greater than 7.0 within the rate ranges.		
	Do not apply more than 12.0 oz./A per year.		

BE		

- Can be used on LibertyLink®, Roundup Ready®, Roundup Ready®
  2 Xtend, or non-GMO soybeans.
- Can also be used in an Enlist Duo® herbicide system on Enlist E3™ soybeans.
- Excellent foundational broadleaf herbicide for unmatched residual control of small-seeded broadleaf weeds. Controls waterhemp, pigweed, nightshade, lambsquarters, kochia, morningglories, smartweed, velvetleaf, wild buckwheat and nutsedge.
- Provides excellent suppression of early-season grasses (foxtails, shattercane) to allow for reduced postemerge trips in-season.
- For resistant weed populations, use higher rate in recommended rate range from label for longer residual control.
- Tank mix with other herbicides for full-season grass control or use as part of a planned pre/post program.
- For no-till, add Aim® EC herbicide 0.75 fl. oz. + 2,4-D 1 pt./A + COC 1% or other appropriate burndown program.
- · Active Ingredient: Imazethapyr, Sulfentrazone.



USE RATES BY SOIL TEXTURE- FL. OZ./A					
ORGANIC MATTER %	COARSE	MEDIUM	FINE		
>1 - 2%	6 – 8	8 – 10	10 – 12		
2 - 4%	8 – 10	10 – 12	12		

ROTATIONAL CROP GUIDELINES			
Wheat, Lima Beans	4 months		
	4 months		
Rye	18 months (MN/ND north of Hwy #210)		
Tobacco, Barley	9.5 months		
Corn, Field* and Seed	10 months		
Alfalfa	12 months		
Cotton, Sunflowers	18 months		
Sweet Corn, Sorghum	18 months		

\*IR, Clearfield\* and IMR corn hybrids may be planted after four months where Authority Assist herbicide was applied at 4 oz. or less.



Untreated



Authority Assist herbicide 9 fl. oz./A SDSU. Pictures 30 days after planting.



