

Coragen[®] Insect Control

Powered by Rynaxypyr[®] active



STRAWBERRIES

Help realize optimal yields and quality in strawberries with reliable and consistent control.

Product Information:

Coragen[®] insect control powered by Rynaxypyr[®] active is a breakthrough mode of action insecticide that delivers remarkable protection. Help optimize yields and quality in your fields by achieving reliable and consistent control of key insect pests in strawberries.

The Benefits of Using Coragen Insect Control:

- Breakthrough mode of action versus current standards helps break the insect resistance cycle.
- Excellent control of major worm pests.
- No signal word.
- Works through ingestion or contact to deliver ovicidal¹, ovi-larvicidal, larvicidal and adulticide² activity.
- Fast acting crop protection. Insects stop feeding in minutes, helping improve crop quality and reducing routes for disease entry.

- Long-lasting protection provides extended residual control and reduces the number of spray applications across the season.
- Reduces pest populations and helps keep them down.
- Shortest re-entry (four hours) and short preharvest interval (one day for strawberries) for added flexibility.
- Application flexibility: apply by foliar spray, ground and air, whichever fits your production system.
- Minimal impact on most beneficial insects when applied at labeled rates; excellent fit with IPM programs³.
- Rainfast once spray has dried.

Active ingredient name: Chlorantraniliprole

Chemical class: IRAC Group 28, anthranilic diamide

Formulation: 1.67 SC

Packaging: 1-gallon container or 1-quart container

¹Ovicidal activity on selected pest species.

²Based on lab studies, adult mortality is dependent on species, application rate, exposure level and time depending on application method.

³In line with integrated pest management and good agricultural practices, insecticide applications should be made when pollinators are not foraging to avoid unnecessary exposure.



Coragen® insect control powered by Rynaxypyr® active for strawberries

Application Method	Targeted Pests ⁴	Application Rate Per Acre		REI (hours)	Last Application (days to harvest)
Foliar	Beet armyworm, cabbage looper, corn earworm, Japanese beetle (adult), light brown apple moth	0.045-0.98 lb ai	3.5-7.5 fl oz.	4	1

⁴ See individual product labels for specific pests controlled or suppressed by these products.

Make no more than four applications per acre per crop.

Minimum interval between treatments is seven days.

Do not apply more than 15.4 fl. oz. Coragen insect control or 0.2 lb. a.i. of chlorantraniliprole-containing products per acre per crop.

Do not apply more than 30.8 fl. oz. Coragen insect control or 0.4 lb. a.i. of chlorantraniliprole-containing products per acre per year. In NY, do not apply more than 15.4 fl. oz. Coragen insect control or 0.2 lb. a.i. of chlorantraniliprole-containing products per acre per year.

Contact your local FMC retailer or representative to learn how you can get more consistent control of key pests in strawberries with Coragen insect control. Visit us at [Ag.FMC.com](https://www.Ag.FMC.com).

Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo, Coragen and Rynaxypyr are trademarks of FMC Corporation or an affiliate. ©2023 FMC Corporation. All rights reserved. 23-FMC-1504 09/23

