

Mustang[®] Maxx Insecticide



CORN AND SOYBEANS

Better yields and more dollars per acre with Mustang[®] Maxx insecticide

Mustang[®] Maxx insecticide brings the pyrethroid difference to your corn and soybean crops, fighting yield-robbing infestations of cutworms, beetles, grasshoppers, aphids, stinkbugs and others before they take hold. It is especially useful for late-season seed corn protection.

The enhanced pyrethroid chemistry of Mustang Maxx insecticide works on contact and ingestion, and has strong residual control of many pests. It also makes a great tank-mix partner with other products.

For better yields and dollar-per-acre return on your investment, choose Mustang Maxx insecticide.

Quick Facts

- Enhanced pyrethroid chemistry works on contact with residual control of many pests.
- For at-plant, pre- or post-emergence application.
- Proven, broad-spectrum insecticide for higher yields.
- Controls more than 45 pests in corn and soybeans.
- Controls widest range of insects; few insecticides can make same claim.
- May be applied by ground, air and chemigation.
- Can be tank mixed with other labeled products such as fungicides, herbicides or spray adjuvants.



Product Information:

Active Ingredient: Zeta-cypermethrin

Chemistry Class: Pyrethroid, IRAC Group 3A

Formulation: 0.8 lb. ai/gal.

Signal Word: WARNING

Pests Controlled:

CORN (FIELD)	CORN (SWEET)	SOYBEANS	
Adult Japanese Beetle	Adult Corn Rootworm	Alfalfa Caterpillar	Mexican Bean Beetle
Adult Sap Beetle	Adult Japanese Beetle	Bean Leaf Beetle	Painted Lady Caterpillar
Aphid	Adult Sap Beetle	Beet Armyworm	Pea Leaf Weevil
Armyworm	Aphid	Blister Beetle	Plant Bug
Bean Leaf Beetle	Armyworm	Colorado Potato Beetle	Saltmarsh Caterpillar
Corn Borer (European, Southwestern)	Chinch Bug	Corn Earworm	Seedcorn Maggot
Cereal Leaf Beetle	Corn Borer	Cowpea Curculio	Cucumber Beetle
Chinch Bug	Corn Earworm	Cutworm	Silver-Spotted Skipper
Corn Earworm	Corn Silkworm	European Corn Borer	Southern Armyworm
Corn Rootworm Beetle	Flea Beetle	Flea Beetle	Soybean Aphid
Cutworm	Grasshopper	Fall Armyworm	Spittlebug
Flea Beetle	Leafhopper	Flea Beetle	Stink Bug
Grasshoppers	Tarnished Plant Bug	Green Cloverworm	Three-Cornered Alfalfa Hopper
Green Cloverworm		Grasshopper	Thrip
Hop Vine Borer		Hornworm	Tobacco Budworm
Hornworms		Imported Cabbageworm	True Armyworm
Meadow Spittlebug		Japanese Beetle	Velvetbean Caterpillar
Southern Corn Leaf Beetle		Kudzu Bug	Webworm
Stalk Borer		Leafhopper	Whitefly
Stink Bug spp		Leaf Miner	Woolly Bear Caterpillar
Tobacco Budworm		Leaf Skeletonizer	Yellow Striped Armyworm
Webworm		Lesser Cornstalk Borer	
Western Bean Cutworm		Looper	

Guidelines for Use:

	PRODUCT RATE (FL. OZ./A)	PHI (DAYS)	REI (HOURS)
Corn (Field)*	1.28-4.0	7	12
Corn (Sweet)**	2.24-4.0	3	12
Soybeans***	1.28-4.0	21	12

*Do not apply more than 16 fl. oz. (0.10 lb./A) per acre per year including at-plant applications.

**Apply at minimum three to five day intervals. Do not apply more than 24 fl. oz. (0.15 lb.) per acre per year. Use a minimum of 20 GPA finished spray by ground and 2 GPA by air.

***Do not make applications less than 7 days apart. Do not apply more than 24 fl. oz. (0.15 lb.) per acre per year.

For more information about Mustang Maxx insecticide, contact your FMC retailer or visit Ag.FMC.Com.

Mustang Maxx insecticide is a Restricted Use Pesticide. Always read and follow label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo and Mustang are trademarks of FMC Corporation or an affiliate. ©2023 FMC Corporation. All rights reserved. 23-FMC-1504 06/18

