

## POISON

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**ACTIVE CONSTITUENTS:**  
100 g/L FLUTRIAFOL  
4 g/L CYPERMETHRIN

**CONTENTS:**  
10 kg



GROUP	<b>3</b>	FUNGICIDE
GROUP	<b>3A</b>	INSECTICIDE

Seed dressing for treatment of wheat and barley as per the Directions for Use Table.

### SAFETY DIRECTIONS

Armour® C Seed Dressing is poisonous if swallowed. May irritate the eyes and skin. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Do not inhale dust. When preparing and using Armour® C Seed Dressing wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

### SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet that can be obtained from [www.fmccrop.com.au](http://www.fmccrop.com.au)

## GENERAL INSTRUCTIONS

GROUP	<b>3A</b>	INSECTICIDE
-------	-----------	-------------

### FUNGICIDE RESISTANCE WARNING

GROUP	<b>3</b>	FUNGICIDE
-------	----------	-----------

Armour® C Seed Dressing is a member of the DMI group of fungicides. For fungicide resistance management Armour® C Seed Dressing is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Armour® C Seed Dressing and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Armour® C Seed Dressing or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, FMC accepts no liability for any losses that may result from the failure of Armour® C Seed Dressing to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

For insecticide resistance management Armour® C Seed Dressing is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Armour® C Seed Dressing and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Armour® C Seed Dressing or other Group 3A insecticides are used repeatedly. The effectiveness of Armour® C Seed Dressing on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, FMC accepts no liability for any losses that may result from the failure of Armour® C Seed Dressing to control resistant insects.

Armour® C Seed Dressing may be subject to specific resistance management strategies. For further information contact your local supplier, FMC representative or local agricultural department agronomist.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Short coleoptile cultivars should not be sown too deep. In some situations trifluralin herbicides and triazole fungicides interact to suppress emergence, if possible avoid this contamination. Any seed not intended to be used for sowing should be destroyed.

### PROTECTION OF LIVESTOCK

Seed treated with Armour® C Seed Dressing must not be used for

animal consumption or be allowed to contaminate seed intended for animal consumption. Seed should not be left exposed on the ground (particularly during sowing).

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**DO NOT** feed or expose treated seed to wildlife and domestic animals, particularly birds.

**Harmful to fish. DO NOT** contaminate streams, rivers or waterways with the chemical or containers which have held the chemical or treated seed.

#### **STORAGE AND DISPOSAL**

Seed treated with Armour® C Seed Dressing must not be used for human consumption or be allowed to contaminate seed intended for human consumption. Treated seed when stored should be kept apart from other seed and the containers should be clearly marked to indicate the contents have been treated with Armour® C Seed Dressing. Bags which have held treated seed should not be used for any other purpose.

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food, foodstuffs, seed and fertilisers.

Shake empty bag into application equipment. **DO NOT** dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the bags below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## DIRECTIONS FOR USE

### RESTRAINT:

DO NOT apply 'Vydate' to seed pre-treated with Armour® C Seed Dressing.

Crop	Disease controlled and insect pestprotection	State	Rate per 100 kg seed	Critical Comments
Wheat	Speckled leaf blotch ( <i>Septoria tritici</i> )	NSW, Vic, Tas, SA, WA only	100 g	Seed must be treated to ensure even coverage of all grains, using correctly calibrated seed treatment equipment.  Reduced emergence may occur if seed treated with flutriafol is sown to a depth greater than 5 cm or in soil at a temperature less than 5°C or greater than 20°C.  Seed treatment will give suppression of leaf diseases for up to 8 weeks after sowing.
	Bunt ( <i>Tilletia foetida</i> , <i>T. caries</i> ) Loose Smut ( <i>Ustilago segetum</i> var. <i>tritici</i> ) Stripe/Yellow Rust ( <i>Puccinia striiformis</i> ) in young crops. Soilborne Flag Smut ( <i>Urocystis agropyri</i> ) Seedborne Flag Smut ( <i>Urocystis agropyri</i> )			
Barley	Barley Scald ( <i>Rhynchosporium secalis</i> ) Covered Smut ( <i>Ustilago segetum</i> var. <i>segetum</i> ) Loose Smut ( <i>Ustilago segetum</i> var. <i>tritici</i> ) Powdery Mildew ( <i>Erysiphe graminis</i> ) in young crops			
Wheat and Barley	Protection against insect pests of stored grains: Lesser grain borer ( <i>Rhyzopertha dominica</i> ) Granary weevil ( <i>Sitophilus granarius</i> ) Rice weevil ( <i>Sitophilus oryzae</i> ) Rust-red flour beetle ( <i>Tribolium castaneum</i> ) Saw toothed grain beetle ( <i>Oryzaephilus surinamensis</i> ) Tropical warehouse moth ( <i>Ephesia cautella</i> )			

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS

**DO NOT GRAZE PLANTS GROWN FROM TREATED SEED, OR CUT FOR STOCK FOOD, WITHIN 4 WEEKS OF SOWING.**

### NOTICE TO BUYER

To the extent permitted by the Competition and Consumer Act (2010) or any relevant legislation of any State or Territory (the "Legislation") all conditions and warranties and statutory or other rights of action, whether arising in contract or tort or whether due to the negligence of FMC or Seller, which buyer or any other user may have against FMC or Seller are hereby excluded provided however that any rights of the buyer pursuant to non-excludable conditions or warranties of the Legislation are expressly preserved. FMC hereby gives notice to buyer and other users that to the extent permitted by the Legislation it will not accept responsibility for any indirect or consequential loss of whatsoever nature arising from the storage, handling or use of this product. Where permitted by the Legislation FMC's liability shall in all circumstances be limited to the replacement of the product, or a refund of the purchase price paid therefor.

The product must be used and applied strictly in accordance with the label instructions and other directions for use. It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of FMC or the Seller. Buyer accepts these risks.

---

**IN A MEDICAL EMERGENCY CALL 1800 033 111 ALL HOURS**

---

The information set forth herein is furnished free of charge and is based on technical data that FMC believes to be reliable. It is intended for use by persons having technical skill at their own discretion and risk. FMC makes no warranties, expressed or implied, and assumes no liability in connection with any of this information. Nothing herein is to be taken as license to operate under or a recommendation to infringe on any patents.

FMC and Armour® C are trademarks of FMC Corporation or an affiliate.  
© 2021 FMC Corporation. All rights reserved.

\*Not an FMC trademark.

APVMA Approval No.: 30252/0906

FMC Australasia Pty Ltd.  
A.B.N. 45 095 326 891  
Level 2, Building B  
12 Julius Ave  
North Ryde NSW 2113  
Phone: 1800 066 355  
[www.fmccrop.com.au](http://www.fmccrop.com.au)

---

**SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111  
ALL HOURS – AUSTRALIA WIDE**

---