



POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT:
250 g/L LAMBDA-CYHALOTHRIN

CONTENTS:
1 L, 5 L



GROUP **3A** INSECTICIDE

For the control of certain insect pests in Barley, Cotton, Wheat and various field crops as per the Directions for Use.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Facial contact may cause temporary facial numbness. Avoid contact with eyes and skin. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If in eyes, hold open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet that can be obtained from www.fmccrop.com.au

GENERAL INSTRUCTIONS

RESISTANCE WARNING

GROUP **3A** INSECTICIDE

For insecticide resistance management, Cyhella® is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Cyhella® and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Cyhella® and other Group 3A insecticides are used repeatedly. The effectiveness of Cyhella® on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, FMC accepts no liability for any losses that may result from the failure of Cyhella® to control resistant insects.

Cyhella® may be subject to specific resistance management strategies. For further information contact your local supplier, FMC Australasia Pty Ltd representative or local agricultural department agronomist.

Helicoverpa armigera (Heliothis) resistance in Nth NSW and Qld: To help contain pyrethroid resistance in *H. armigera*, the Summer Crop Insecticide Strategy as developed by AIRAC, Qld Department of Primary Industries and the NSW

Department of Agriculture and Fisheries should be adhered to. Failure to observe the strategy may result in widespread resistance affecting the future viability of summer cropping.

MIXING: SHAKE WELL BEFORE USE

For ground or aircraft application with water: Cyhella® mixes readily with hard or soft water. Add the required quantity of product to water whilst under agitation to ensure thorough mixing. Agitate while spraying. It is not advisable to allow the mixed solution to stand longer than 24 hours before use. In extremely alkaline water (pH 9) spray immediately after mixing.

For ULV (Ultra Low Volume) application with oil: It is recommended that Cyhella® is mixed with a mineral spraying oil. See compatibility section for list of recommended mineral spraying oils. Add the required amount of product whilst under agitation to ensure thorough mixing. Agitate while spraying. It is not advisable to allow the mixed solution to stand longer than 24 hours before use.

APPLICATION

Good coverage is essential to ensure adequate control. The product may be applied by ground rig or aircraft.

Acceptable threshold values for eggs and larval numbers may vary according to the stage of the crop development and the pest management undertaken. Alternative higher thresholds

may be acceptable under certain circumstances.

Diluted with water: For ground rigs the volume of liquid applied should be 50 to 100 L/ha. Aerial application should be under conditions normally suitable for water based insecticides. Apply in at least 10 to 20 litres of water per hectare.

Mixed with oil: Apply the recommended rate of Cyhella® bulked with oil to total volume of 3 to 5 L/ha for cotton, sorghum and sunflowers. The total volume for all other crops should be 1.5 litres per hectare.

TIMING

This product is a contact and residual insecticide. Best results will be obtained if Cyhella® is applied as a protective treatment at regular intervals. However if spraying frequency is based on scouting, then for *Helicoverpa* spp application at egg hatch will give optimum results.

CROP CHECKING

Frequent and thorough checking of whole plants, terminals, squares, flowers, bolls or fruiting bodies as required, should be made over a random sample of plants, representative of the whole crop area.

Inspect crops after spraying to ensure a thorough kill has been obtained, however note that maximum kill may not be achieved until 48 hours after treatment. Then check at frequent intervals, not more than 2 days apart when insect pressure is heavy. Apply the recommended treatment as soon as crop check indicates spray is necessary.

COMPATIBILITY

This product, when applied as a water based spray is compatible with the following actives: pirimophos methyl, procymidone, fluazifop-P-butyl, paraquat, pirimicarb, paraquat plus diquat, and glyphosate.

PRECAUTION

Human flagging is not supported unless flaggers are protected by engineering controls such as vehicles with cabs.

RE-ENTRY PERIOD

DO NOT allow entry into treated fields/crops until the spray

has dried. If prior entry is necessary, wear cotton overalls and chemical resistant gloves.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Dangerous to fish and aquatic organisms. DO NOT contaminate streams, rivers or waterways with the product or the used containers. Tail waters which flow from treated areas should be prevented from entering river systems. A strategy to minimise spray drift should be employed at all times when aerially applying sprays near sensitive areas. Such a strategy is illustrated by the cotton's Best Management Practice Manual.

PROTECTION OF LIVESTOCK

Toxic to bees. DO NOT spray when bees are actively foraging. Risk is reduced by spraying in the early morning or late evening.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals onsite. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear protective equipment (see Safety Directions). Apply absorbent material such as earth, clay granules or cat clumping litter to the spill. Sweep up the material for disposal when absorption is completed and contain in a refuse vessel for disposal (see Storage and Disposal). If necessary, wash the spill area with an alkaline detergent and water and absorb, as above, the wash liquid for disposal.

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply if rain is expected within 6 hours.

For ULV application: Cyhella® can be bulked up with spraying oils for all uses except those indicated in the critical comments.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Barley, Wheat	Blackhead Pasture Cockchafer (<i>Aphodius tasmaniae</i>)	NSW, Vic, Tas, SA, WA only	20 or 40 mL/ha	14 days (H/G)	Treat as soon as possible after the autumn rains stimulate egg hatching and activity of existing larvae. This can be ascertained by monitoring soil populations in known areas. For best results spray when larvae have surfaced to feed after rain. Preferably use a boom spray delivering 70 to 100L water/ha. Use the lower rate until early June and the higher rate after mid-late June. DO NOT USE UVL APPLICATION FOR THIS PEST.
	Brown or Pink Cutworm (<i>Agrotis munda</i>)	All States	12 or 18 mL/ha		For best results apply at first sign of infestation before larvae are 10 mm long. If larvae are larger than 10 mm use the higher rate. Use a minimum 50 L water.
	Common Cutworm (<i>Agrotis infusa</i>)	NSW only			
	Pasture Webworm (<i>Hednota spp</i>)	NSW, Vic, Tas, SA, WA only	12 mL/ha		Pre-seeding: The product can be tank mixed with knockdown herbicides. Post-crop emergence: Inspect crop regularly from sowing. Spray at first sign of damage. Use a minimum 50 L water/ha. Apply at first sign infestation before larvae are 10 mm long.
	Redlegged Earthmite (<i>Halotydeus destructor</i>)		9 mL/ha		If mites are present on an establishing crop, apply at first sign of crop emergence. Monitor crop regularly for reinfestation and respray if necessary.
	Aphids (<i>Rospalosiphum spp</i>) (Barley Yellow Dwarf Virus vectors)		12 or 18 mL/ha		To control aphids spray should be applied at 4 and 8 weeks after emergence to reduce aphid colonisation and suppress Barley Yellow Dwarf Virus. Use the higher rate when greater than 15 aphids on 50% of tillers is expected during the season.
Broccoli, Brussels Sprouts, Cabbage, Cauliflowers, Forage Brassicas	Cabbage Cluster Caterpillar (<i>Crociodolomia parvonana</i>). Cabbage White Butterfly (<i>Pieris rapae</i>), Diamond Back Moth (<i>Plutella xylostella</i>)	All States	24 or 36 mL/ha plus non ionic spray adjuvant at 10 mL/100L spray volume.	2 days (HG)	Apply at first sign of infestation. For schedule spraying on a weekly basis, use the lower rate. For spraying as needed, use the higher rate for longer persistence. Use a minimum 500 L water/ha.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Canola	Cabbage White Butterfly (<i>Pieris rapae</i>), Cabbage Moth/Diamond Back Moth (<i>Plutella xylostella</i>)	All States	24 mL/ha	7 days (HG)	Apply as soon as larvae reach threshold numbers. Check with local officer of the Department of Agriculture for thresholds applicable to the particular growth stage of the crop.
	Grey Cluster Bug, Rutherglen Bug (<i>Nysius spp</i>)		36 mL/ha		Apply only near maturity when severe infestations are likely to downgrade yields.
	Native Budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, Tas, WA only	24 or 36 mL/ha		For best results, apply at hatching or soon after. Use higher rate if the crop is dense or the larvae are larger than 10 mm.
	Thrips (<i>Thrips tabaci</i>)	Qld, NSW, Vic, Tas, WA, NT	36 mL/ha		Apply only near maturity when severe infestations are likely to downgrade yields.
	Redlegged Earthmite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA, WA only	9 mL/ha		If mites are present on an establishing crop, apply at the first sign of crop emergence. Monitor the crop regularly for reinfestation and respray if necessary.
Chickpeas, Faba Beans, Lentils, Vetch	Native Budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, SA, WA only	24 or 36 mL/ha	7 days (H/G)	For best results apply at hatching or soon after. Use the higher rate if the crop is dense or the larvae are larger than 10 mm.
	Redlegged Earthmite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA, WA only	9 mL*/ha		If mites are present on an established crop, apply at first sign of crop emergence. Monitor crop regularly for reinfestation and respray if necessary. Control of Lucerne Flea will not be obtained with application.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Cotton	Apple Dimpling Bug (<i>Campylomma liebknechti</i>), Brokenbacked bug (<i>Taylorilygus pallidulus</i>), Brown Mirid (<i>C. pacificus</i>), Cottonseed Bug (<i>Oxycarenus luctuosus</i>), Green Mirid (<i>Creontiades dilutus</i>), Leafhoppers (<i>Aus-troasca viridigrisea</i> , <i>Amrasca terraereginae</i>), Pale Cotton Stainer (<i>Dysdercus sidae</i>)	Qld, NSW, WA, NT only	60 mL/ha	21 days (H)	Apply at recommended threshold levels as indicated by field checks.
	Cotton Bollworm (<i>Helicoverpa armigera</i>), Native Budworm (<i>Helicoverpa punctigera</i>)		60 mL/ha		Apply when egg laying is light, less than 25 eggs/100 terminals and no larvae are present.
			70 mL/ha		Apply when egg laying is moderate, greater than 25 eggs/100 terminals and/or when less than 12 newly hatched larvae/100 terminals are present.
			85 mL/ha		Apply when egg laying is heavy and continuous and/or when <i>H.punctigera</i> larvae are greater than 10 mm in length. For <i>H. amigera</i> , apply only to larvae less than 5 mm in length.
	Pink-Spotted Bollworm (<i>Pectinophora scutigera</i>)	Qld, NT only	70 mL/ha		Controlled with the <i>Helicoverpa spp</i> program when used at this rate. If the Pink-Spotted Bollworm is the only pest present, apply when more than 10 adult moths are caught in pheromone traps on 2 consecutive nights.
Field Peas	Native Budworm (<i>Helicoverpa punctigera</i>)	All States	24 or 26 mL/ha	7 days (H/G)	For best results, apply at hatching or soon after. Use higher rate if the crop is dense or the larvae are larger than 10 mm.
	Pea Weevil (<i>Bruchus pisorum</i>)	NSW, SA only	24 mL/ha		SA only: Follow State Department of Agriculture guidelines for controlling Pea Weevil. If these are unavailable, monitor the crops regularly once flowering commences and apply as soon as adult weevils are detected. Adults must be controlled before egg laying begins. Both Native Budworm and Pea Weevil populations can be easily monitored using a sweep net in the top section of the crop. WA only: Commence monitoring the crop for Pea Weevil presence using a sweep net prior to flowering. Spray when 1 weevil/100 sweeps is found for milling grade seed, or 1 weevil/25 sweeps for feed grade seed. Continue monitoring after spraying and respray if necessary. Use either a border spray (most cases) or whole crop spray, depending on Pea Weevil penetration of the crop.
		Vic, WA only	36 mL/ha		
Redlegged Earth Mite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA, WA only		9 mL*/ha		If mites are present on an establishing crop, apply at the first sign of crop emergence. Monitor crop regularly for reinfestation and respray if necessary. Control of Lucerne Flea will not be obtained with this application.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Lemons, Oranges	Fullers Rose Weevil (<i>Asynonychus cervinus</i>)	All States	300 mL/100L as a directed spray	4 weeks (H)	Firstly ensure that the trees are skirted and all weeds under the trees are removed. Apply 250 mL spray solution to the tree trunk at about 300 mm from the ground in a 100 mm band. Deliver the spray through a U shaped wand fitted with 4 nozzles evenly spaced around the tree. Trees must be treated in the early stages of adult weevils emerging from the ground.
Lucerne	Blackhead Pasture Cockchafer (<i>Aphodius tasmaniae</i>)	NSW, Vic, Tas, SA, WA only	20 or 40 mL/ha	14 days (H/G)	Treat as soon as possible after the autumn rains stimulate egg hatching and activity of existing larvae. This can be ascertained by monitoring soil populations in known areas. For best results spray when the larvae have surfaced to feed after rain. Preferably use a boom spray delivering 70 to 100 L water/ha. Use the lower rate until early June and the higher rate after mid-late June. DO NOT USE ULV APPLICATION FOR THIS PEST.
	Lucerne Leaf Roller (<i>Merophyas divulsana</i>)	All States	24 or 36 mL/ha		For best results apply at hatching or soon after. Use higher rate if the crop is dense or the larvae are larger than 10 mm. Apply the first spray when about 30% of the terminals are rolled.
	Native Budworm (<i>Helicoverpa punctigera</i>)				For best results apply at hatching or soon after. Use higher rate if the crop is dense or the larvae are larger than 10 mm.
	Pea Aphid				24 mL/ha
	Redlegged Earthmite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA, WA only	9 mL/ha		If mites are present on an establishing crop, apply at first sign of crop emergence. Monitor crop regularly for reinfestation and respray if necessary. Control of Lucerne Flea will not be obtained with this application.
Lupins	Brown Pasture Looper (<i>Ciampa arietaria</i>)	NSW, Vic, Tas, SA, WA only	12 mL/ha	14 days (H/G)	Once crop has emerged, inspect regularly and apply at the first sign of damage. use a minimum of 50 L water/ha. DO NOT USE ULV APPLICATION FOR THIS PEST.
	Native Budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, SA, WA only	24 mL/ha		For best results, apply at hatching or soon after when larvae are small. WA only: Environmental factors may cause populations of small caterpillars to decline, reducing damage potential. Spraying should commence once caterpillars are 12 mm in length.
	Redlegged Earthmite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA, WA only	9 mL/ha		If mites are present on an establishing crop, apply at the first sign of crop emergence. Monitor crop regularly for reinfestation and respray if necessary. Control of Lucerne Flea will not be obtained with this application.
Mung Beans, Navy Beans	Corn Earworm (<i>Helicoverpa armigera</i>) Native Budworm (<i>Helicoverpa punctigera</i>)	Qld, NSW, NT only	60 or 70 mL/ha	1 day (H/G) 14 days if dry harvest -ed	Apply when flower or pod feeding larvae reach populations of 1 to 2/m of row in navy beans and 1/m of row in mung beans. Use the higher rate if pest numbers are high or if larvae are larger than 10 mm. In Nth NSW and Qld where Corn Earworm has established resistance to pyrethroids DO NOT apply to Corn Earworm larvae larger than 5 mm.
Onions bulb	Onion Thrips	All States	40 mL/ha	14 days (H)	Apply when thrips first appear. Apply via ground equipment in a minimum 300 L water/ha. DO NOT exceed a maximum of 4 applications per crop with a minimum retreatment interval of 7 days between consecutive sprays.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Pasture	Blackheaded Pasture Cockchafer (<i>Aphodius tasmaniae</i>)	NSW, Vic, Tas, SA, WA only	20 or 40 mL/ha	14 days (H/G)	Treat as soon as possible after the autumn rains stimulate egg hatching and activity of existing larvae. This can be ascertained by monitoring soil populations in known areas. For best results spray when the larvae have surfaced to feed after rain. Preferably use a boom spray delivering 70 to 100 L water/ha. Use the lower rate until early June and the higher rate after mid-late June.
	Brown Pasture Looper (<i>Ciampa arietaria</i>)	All States	12 mL/ha		DO NOT USE ULV APPLICATION FOR THIS PEST.
	Brown or Pink Cutworm (<i>Agrotis munda</i>)		12 to 18 mL/ha		For best results apply at first sign of infestation before larvae are 10 mm long. If larvae are larger than 10 mm, use the higher rate. Use a minimum of 50 L of water.
	Common Cutworm (<i>Agrotis infusa</i>)	NSW only			
	Pasture Webworm (<i>Hednota</i> spp)	NSW, Vic, Tas, SA, WA only	12 mL/ha		Apply once larvae are present using adequate water to ensure good penetration.
	Redlegged Earth Mite (<i>Halotydeus destructor</i>)		9 mL/ha		If mites are present on an establishing crop, apply at first sign of crop emergence. Monitor crop regularly for reinfestation and respray if necessary. Control of Lucerne Flea will not be obtained with this application.
Potatoes	Vegetable Jassid (<i>Austroasca viridigrisea</i>)	All States	24 mL/ha	7 days (H)	Apply only when numbers are excessive.
Sorghum	Corn Earworm (<i>Helicoverpa armigera</i>)	Qld, NSW, NT only	60 or 70 mL/ha	14 days (H/G)	Apply when larval numbers reach 2/head. Use the higher rate if pest pressure is severe. Best results are achieved on small larvae.
	Sorghum Midge (<i>Corntarinia sorghicola</i>)		18 or 36 mL/ha		Apply when midge numbers reach 1 to 2/head. Use the higher rate for residual protection.
Soybeans	Corn Earworm (<i>Helicoverpa armigera</i>) Native Budworm (<i>Helicoverpa punctigera</i>)	Qld, NSW, Vic, NT only	60 or 70 mL/ha	21 days (H/G)	soybeans. Use the higher rate if pest numbers are high or if larvae are larger than 10 mm. In Nth NSW and Qld DO NOT apply to resistant <i>H. armigera</i> larvae larger than 5 mm length.
Sunflowers	Corn Earworm (<i>Helicoverpa armigera</i>) Native Budworm (<i>Helicoverpa punctigera</i>)	Qld, Nth NSW only	60 or 70 mL/ha	28 days (H)	Apply when an average of 2 to 3 larvae are present/head or when larvae are damaging plants. Use the higher rate if pest numbers are high and/or <i>H. punctigera</i> larvae are larger than 10 mm in length. In Nth NSW and Qld, DO NOT apply to resistant <i>H. armigera</i> larvae larger than 5 mm in length. General Comments: If flowering has started, application should be deferred until after flowering but before the heads turn down. If treatment is unavoidable during flowering and bees are actively foraging in the crop, there will be minimal effect in the colony if spraying occurs early morning or late afternoon.
	Grey Cluster Bug Rutherglen Bug (<i>Nysius</i> spp)	All States	36 mL/ha		Apply when numbers reach 10 to 15 adults per plant at budding in dry land crops or 20 to 25 in irrigated crops. If <i>Helicoverpa armigera</i> are also present in Nth NSW or Qld, use a minimum 60 mL product.
Tomatoes bush	Native Budworm (<i>Helicoverpa punctigera</i>)	All States	4 to 5 mL/100 L or 30 to 36 mL/ha	1 day (H)	Treat plants on a 7 to 14 day schedule. In Nth NSW and Qld, DO NOT apply to <i>H. armigera</i> larvae larger than 5 mm in length. In other areas for best results apply soon after egg lay. To help contain resistance, alternate sprays between different chemical groups. Check the crop every few days and follow the Summer Crop Insecticide Strategy. There may be phytotoxicity with some varieties, especially Floradade.
	Tomato Bug (<i>Helicoverpa armigera</i>)	Vic, Tas, SA, WA only			
		Qld, NSW, NT only	4 mL/100 L or 60 mL/ha		

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Tomatoes Trellis	Native Budworm (<i>Helicoverpa punctigera</i>) Tomato Bug (<i>Helicoverpa armigera</i>)	All States	4 to 5 mL/100 L	1 day (H)	

^ Blue Oat Mites often co-occur with Redlegged Earth Mites and the 9 mL/ha rates of Cyhella® may be less effective against Blue Oat Mites.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

HARVEST:

Mung Beans (if harvested green), Navy Beans (if harvested green), Tomatoes: **DO NOT** HARVEST FOR 1 DAY AFTER APPLICATION.

Cabbages, Cauliflowers, Broccoli, Brussels Sprouts: **DO NOT** HARVEST FOR 2 DAYS AFTER APPLICATION.

Canola, Chickpeas, Faba Beans, Field Peas, Lentils, Potatoes, Vetch: **DO NOT** HARVEST FOR 7 DAYS AFTER APPLICATION.

Barley, Lucerne, Lupins, Onions, Mung Beans (if harvested dry), Navy Beans (if harvested dry), Pasture, Sorghum, Wheat: **DO NOT** HARVEST FOR 14 DAYS AFTER APPLICATION.

Cotton, Soybeans: **DO NOT** HARVEST FOR 21 DAYS AFTER APPLICATION.

Lemons, Oranges, Sunflowers: **DO NOT** HARVEST FOR 4 WEEKS AFTER APPLICATION.

GRAZING:

Mung Beans (if harvested green), Navy Beans (if harvested green): **DO NOT** GRAZE OR CUT FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION.

Forage Brassicas: **DO NOT** GRAZE OR CUT FOR STOCK FOOD FOR 2 DAYS AFTER APPLICATION.

Canola, Chickpeas, Faba Beans, Field Peas, Lentils, Vetch: **DO NOT** GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

Barley, Lucerne, Lupins, Onions, Mung Beans (if harvested dry), Navy Beans (if harvested dry), Pasture, Sorghum, Wheat: **DO NOT** GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

Soybeans: **DO NOT** GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.

NOTICE TO BUYER

To the extent permitted by the Competition and Consumer Act (2010) or any relevant legislation of any State or Territory (the "Legislation") all conditions and warranties and statutory or other rights of action, whether arising in contract or tort or whether due to the negligence of FMC or Seller, which buyer or any other user may have against FMC or Seller are hereby excluded provided however that any rights of the buyer pursuant to non-excludable conditions or warranties of the Legislation are expressly preserved. FMC hereby gives notice to buyer and other users that to the extent permitted by the Legislation it will not accept responsibility for any indirect or consequential loss of whatsoever nature arising from the storage, handling or use of this product. Where permitted by the Legislation FMC's liability shall in all circumstances be limited to the replacement of the product, or a refund of the purchase price paid therefor.

The product must be used and applied strictly in accordance with the label instructions and other directions for use. It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of FMC or the Seller. Buyer accepts these risks.

IN A TRANSPORT EMERGENCY CALL 000 POLICE OR FIRE BRIGADE

The information set forth herein is furnished free of charge and is based on technical data that FMC believes to be reliable. It is intended for use by persons having technical skill at their own discretion and risk. FMC makes no warranties, expressed or implied, and assumes no liability in connection with any of this information. Nothing herein is to be taken as license to operate under or a recommendation to infringe on any patents.

FMC is a trademark of FMC Corporation or an affiliate. © 2021 FMC Corporation. All rights reserved.
Cyhella® is a trademark of LONZA.

*Not an FMC trademark.

APVMA Approval No: 65190/127156

FMC Australasia Pty Ltd.

A.B.N. 45 095 326 891

Level 2, Building B 12 Julius Ave

North Ryde NSW 2113

Phone: 1800 066 355

www.fmccrop.com.au

SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111
ALL HOURS – AUSTRALIA WIDE
