



Impact®

IN-FURROW AND FOLIAR FUNGICIDE

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SC

FORMULATION TYPE

Suspension
Concentrate

ACTIVE CONSTITUENT:

250 g/L FLUTRIAFOL

CONTENTS:

1000 L

GROUP **3** FUNGICIDE



For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use Table.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet that can be obtained from www.fmccrop.com.au

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

GROUP **3** FUNGICIDE

Impact® In-Furrow and Foliar Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Impact® In-Furrow and Foliar Fungicide is a Group C fungicide. Some naturally occurring individual fungi resistant to Impact® In-Furrow and Foliar Fungicide and other Group C fungicides may exist through normal genetic variability in any fungal population.

The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Impact® In-Furrow and Foliar Fungicide or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FMC accepts no liability for any losses that may result from the failure of Impact® In-Furrow and Foliar Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

FERTILISER APPLICATION

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. **DO NOT** mix seed with treated fertiliser.

FOLIAR APPLICATION

Check the crop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

Triple or preferable pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury containers in local authority landfill. If no landfill is available,

bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

DIRECTIONS FOR USE

A. IMPACT® APPLIED IN-FURROW AS A FERTILISER TREATMENT

| CROP | DISEASE | RATE PER HECTARE | WHP | CRITICAL COMMENTS | | |
|--------|---|------------------|--------------------|--|--|-------------|
| Canola | Blackleg (<i>Leptosphaeria maculans</i>) | 400 mL | 4 weeks Grazing | Treatment will reduce blackleg infection in both susceptible and tolerant Canola varieties. Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400 mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide. | | |
| | | | | <u>Fertiliser rate per hectare (kg)</u> | <u>Volume of product per 100 kg of fertiliser (mL)</u> | |
| | | | | 60 | 667 mL* | |
| | | | | 70 | 571 mL | |
| | | | | 80 | 500 mL | |
| | | | | 100 | 400 mL | |
| | | | | 200 | 333 mL | |
| Wheat | Stripe rust (<i>Puccinia striiformis</i>) | 200 or 400 mL | 4 weeks Grazing | For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe diseases are expected. Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specified rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide. | | |
| | Septoria tritici blotch (<i>S. tritici</i>) Take-all (<i>Geaumannomyces graminis</i>) | 400 mL | | | | |
| Barley | Powdery mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>) | 200 or 400 mL | | <u>Fertiliser rate per hectare (kg)</u> | <u>Volume of product per 100 kg of fertiliser (mL)</u> | |
| | | | | | 200 mL rate | 400 mL rate |
| | | | | 50 | 400 mL | 800 mL* |
| | | | | 60 | 333 mL | 667 mL* |
| | | | | 70 | 286 mL | 571 mL |
| | | | | 80 | 250 mL [^] | 500 mL |
| | | | | 100 | 200 mL [^] | 400 mL |
| | | | | 120 | 167 mL | 333 mL |
| | | | | *Application volumes over 600 mL per 100 kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. [^] Where application volume is low, product may be diluted with water to improve coverage. | | |

B. IMPACT® APPLIED AS A FOLIAR SPRAY

| CROP | DISEASE | RATE PER HECTARE | WHP | CRITICAL COMMENTS |
|--------|--|---|--------------------------------|---|
| Wheat | Stripe rust (<i>Puccinia striiformis</i>) | 250 mL – 500 mL Ground application Apply in 100 – 150 L of water plus 200 mL of spray solution | 7 weeks Harvest or Grazing | Apply when Stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate. |
| | Leaf rust (<i>P. recondita</i>) Septoria Nodorum blotch (<i>Septoria nodorum</i>) Septoria tritici blotch (<i>S. tritici</i>) | Aerial application Apply in a minimum of 15 L of water plus 200 mL 'BS 1000' per 100 L of spray solution. | | Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe disease persist during flowering, use a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate. |
| Barley | Powdery mildew (<i>Erysiphe graminis</i>) | | 10 weeks Harvest or Grazing | Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate, may be required. |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

A. APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY - **DO NOT** GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

B. APPLIED AS A FOLIAR SPRAY

DO NOT HARVEST EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY - 10 WEEKS

WHEAT - 7 WEEKS

DO NOT GRAZE OR CUT FOR STOCK FOOD EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY - 10 WEEKS

WHEAT - 7 WEEKS

NOTICE TO BUYER

To the extent permitted by the Competition and Consumer Act (2010) or any relevant legislation of any State or Territory (the "Legislation") all conditions and warranties and statutory or other rights of action, whether arising in contract or tort or whether due to the negligence of FMC or Seller, which buyer or any other user may have against FMC or Seller are hereby excluded provided however that any rights of the buyer pursuant to non-excludable conditions or warranties of the Legislation are expressly preserved. FMC hereby gives notice to buyer and other users that to the extent permitted by the Legislation it will not accept responsibility for any indirect or consequential loss of whatsoever nature arising from the storage, handling or use of this product. Where permitted by the Legislation FMC's liability shall in all circumstances be limited to the replacement of the product, or a refund of the purchase price paid therefor.

The product must be used and applied strictly in accordance with the label instructions and other directions for use. It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of FMC or the Seller. Buyer accepts these risks.

The information set forth herein is furnished free of charge and is based on technical data that FMC believes to be reliable. It is intended for use by persons having technical skill at their own discretion and risk. FMC makes no warranties, expressed or implied, and assumes no liability in connection with any of this information. Nothing herein is to be taken as license to operate under or a recommendation to infringe on any patents.

FMC and Impact® are trademarks of FMC Corporation or an affiliate. © 2021 FMC Corporation. All rights reserved.

APVMA Approval No.: 49781 / 0207

FMC Australasia Pty Ltd.
Level 2, Building B,
12 Julius Ave,
North Ryde NSW 2113
Australia
Phone: 1800 066 355
www.fmccrop.com.au

SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA WIDE
In a Transport Emergency Dial 000 Police or Fire Brigade
