



Rovral® Aquaflo
BROAD SPECTRUM FUNGICIDE

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SC

FORMULATION TYPE
Suspension
Concentrate

ACTIVE CONSTITUENT:

500 g/L IPRODIONE

CONTENTS:

1 L – 500 L



GROUP **2** FUNGICIDE

For the control of certain fungal diseases in various crops and situations as specified in the Directions For Use table.

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes and skin. When opening the container and using the product wear cotton overalls over normal clothing, buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves and face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26), New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet that can be obtained from www.fmccrop.com.au

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

GROUP **2** FUNGICIDE

Rovral® Aquaflo Fungicide is a member of the dicarboximide group of fungicides. For fungicides resistance management the product is a Group 2 Fungicide. Some naturally occurring individual fungi resistant to the product and other Group 2 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 2 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FMC Australasia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

RESISTANCE MANAGEMENT

Do rotate to use products with as many different modes of action as possible to reduce the possibility of resistance development where alternatives are available.

Where specific resistance management strategies are established these are detailed at the CropLife Australia website

(http://www.croplifeaustralia.org.au/default.asp?V_Doc_ID=1792) or from your local agronomist.

MIXING

Note: Rovral® Aquaflo may be unstable in conditions where the pH is 7 or higher. It is therefore essential to check the pH of the spray solution before adding Rovral® Aquaflo. A suitable registered buffering agent may have to be added to bring the pH down below 7.

Add the required amount of Rovral® Aquaflo to the spray tank containing half the required volume of water. Mix thoroughly and make up to the required volume with water.

APPLICATION

Good disease control requires even, thorough coverage of the target area. Application should be made using appropriate spray equipment and sufficient water to provide adequate penetration and coverage. Equipment settings and water volume may need to vary, depending on the growth stage of the crop.

Special instructions for Tree Crops/Vines

Dilute Spraying

- Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient spray solution to cover the crop to the point of run-off. Avoid excessive run-off.
- The required spray volume to achieve point of run off may

be determined by applying different test volumes, using different settings on the sprayer, or from industry guidelines or other expert advice.

- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume to achieve point of run-off will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies spray volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen spray volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1500 L/ha
 2. Your chosen concentrate spray volume: For example 500 L/ha
 3. The concentration factor in this example is: $3 \times$ (i.e. $1500 \text{ L} \div 500 \text{ L} = 3$).
 4. If the dilute label rate is 10 mL/100 L, then the concentrate rate becomes 3×10 , that is 30 mL of product per 100 L water for concentrate spraying.
- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
 - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

TURF NUTRITION

This product acts very rapidly and where the turf is of a low nutritional status, a slight yellowing may be seen. This will be corrected by nitrogenous fertiliser application. Where possible, it is recommended that the fertiliser programme should precede the fungicide application.

COMPATIBILITY

This product may be combined with:

*Aliette® WG (see NOTE below), Bugmaster® Flowable, calcium chloride, chlorpyrifos 500 g/L EC, copper oxychloride, dimethoate, Dithane® DF, Dithane M-45®, DPA, endosulfan (e.g. Thiodan® EC), fenitrothion, Kelthane® EC, maldison, metalaxyl, methomyl (e.g. Marlin®) or parathion-methyl.

When tank mixing products the order of mixing is determined by formulation type. As a guide the following mixing sequence is recommended:

1. Wettable powders
2. Suspension concentrates (e.g. Rovral® Aquaflo)
3. Water dispersible granules
4. Suspo-emulsions
5. Soluble powders
6. Solutions
7. Emulsifiable concentrates
8. Soluble concentrates
9. Wetting agents and oils

With any mixture, thoroughly agitate immediately before applying. It is not recommended to mix this product with more than one of the above chemicals in the tank. **DO NOT** tank mix this product with fertilisers.

NOTE: *Mixing Rovral® Aquaflo with Aliette® WG may result in some settling out.

As formulations of other manufacturers' products are beyond the control of FMC Australasia Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT graze treated areas or feed turf clippings from treated areas to animals including poultry.

REFER to INDEX on page 9 for the LISTING of each CROP and DISEASES

Tree Crops/Vines:

RATE					CRITICAL COMMENTS
CROP	DISEASE	STATE	RATE	WHP	
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Special Instructions for Tree Crops/Vines section.					For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Refer to the Special instructions for Tree Crops/Vines section.
Almonds	Blossom blight, brown rot (<i>Monilinia spp.</i> , <i>Sclerotinia spp.</i>)	All States	50 mL/100 L water	Nil	
Grapes	Grey mould (<i>Botrytis cinerea</i>)	NSW, Vic, WA only	100 mL / 100 L water	7 days	<p>Good crop hygiene will aid in the control of disease.</p> <p>This use is subject to a CropLife Australia fungicide resistance management strategy.</p> <p>The number of consecutive applications and the total number of applications of Group 2 Fungicides permitted is limited. Refer to CropLife Australia Resistance Management Guidelines. See the "General Instructions - Resistance Management" for details on where these guidelines can be obtained.</p> <p>Apply the spray to vines every 10 to 14 days ensuring that all fruit is thoroughly wet. Apply 3 applications at 10 to 14 day intervals from 10% bloom to petal fall for protection of flowers and young fruit. Apply a further 2 applications of Rovral® Aquaflor to control late season Botrytis.</p>
Kiwifruit	Botrytis blight (<i>Botrytis spp.</i>)				
Macadamias	Botrytis blight (<i>Botrytis spp.</i>)	All States	50 mL/100 L water	Nil	Apply as a thorough cover spray to flower racemes when they open. A follow up spray may be needed one week later if wet conditions persist during flowering. Remove nuts under trees prior to spraying.
Mandarins (non-bearing)	Alternaria leaf spot (brown spot)	Qld, WA, NT only	100 mL/100 L water		Apply to non-bearing trees of Murcott variety monthly from first flush in spring until flushing ceases in the autumn. Reduce intervals to fortnightly during periods of wet weather.
Passionfruit	Alternata spot (brown spot)	Qld, NSW, WA, NT only	100 mL/100 L water	1 day	<p>This use is subject to a CropLife Australia fungicide resistance management strategy:</p> <ol style="list-style-type: none"> Maintain a protective cover with protectant fungicide such as mancozeb. Limit the use of Rovral® Aquaflor to strategic periods, i.e. before, during and after extended wet periods. Always tank mix Rovral® Aquaflor with a protectant such as mancozeb. DO NOT apply more than four Rovral® Aquaflor (or other Group 2 fungicide) sprays in a season.

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Stone Fruit: Apricots, cherries, nectarines, peaches, plums	Orchard Spraying Blossom blight <i>(Monilinia fructicola,</i> <i>Monilinia laxa)</i> Brown rot <i>(Monilinia fructicola,</i> <i>Monilinia laxa)</i>	Qld, NSW, Vic, Tas, SA, WA only	50 to 75 mL/ 100 L water	Nil	<p>Critical timings for control of blossom blight, are 10% blossom, full bloom and petal/shuck fall and for control of subsequent brown rot in fruit, 3 weeks and 1 week pre-harvest. Use the higher rate under severe conditions of challenge, or for single applications of Rovral® Aquaflo in the spray program.</p> <p>This use is subject to a CropLife Australia fungicide resistance management strategy.</p> <p>The number of consecutive applications of Group 2 fungicides permitted is limited. Refer to the CropLife Australia Resistance Management Guidelines. See the "General Instructions - Resistance Management" for details on where these guidelines can be obtained.</p>

Fruit - Post-harvest dipping:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Pome fruit: Apples, pears	Post-harvest dipping Storage rots <i>(Penicillium spp.)</i> <i>(Botrytis spp.)</i> <i>(Gloeosporium spp.)</i>	All States	100 mL/100 L water	Nil	<p>To minimise the development of post-harvest rots handle fruit carefully to avoid fruit injury and dip promptly after harvest.</p> <p>Remove any infected fruit from the packing house immediately and destroy. When dipping, allow sufficient time to thoroughly wet the fruit. Top up dip with 100 mL Rovral® Aquaflo in 100 L of water.</p> <p>When using Rovral® as a post-harvest dip, DO NOT use a Group 2 fungicide as the final pre-harvest spray.</p> <p>Rotation between different modes of action in post-harvest dips is advised.</p> <p>This use is subject to a CropLife Australia fungicide resistance management strategy.</p> <p>See the "General Instructions - Resistance Management" for details on where these guidelines can be obtained.</p>
Stone Fruit: Apricots, cherries, nectarines, peaches, plums	Post-harvest dipping Brown rot <i>(Monilinia fructicola,</i> <i>Monilinia laxa)</i> * Transit rot <i>(Rhizopus spp.)</i>	Qld, NSW, Vic, Tas, SA, WA only			<p>To minimise the development of post-harvest rots handle fruit carefully to avoid fruit injury and dip promptly after harvest. Remove any infected fruit from the packing house immediately and destroy. When dipping, allow sufficient time to thoroughly wet the fruit. Top up dip with 200 mL Rovral® Aquaflo in 100 L of water. A non-ionic wetting agent should be added.</p> <p>*Transit rot is suppressed at this rate.</p> <p>When using Rovral® Aquatic as a post-harvest dip, DO NOT use a Group 2 fungicide as the final pre-harvest field spray.</p> <p>Rotation between different modes of action in post-harvest dips is advised.</p> <p>This use is subject to a CropLife Australia fungicide resistance management strategy. See the "General Instructions - Resistance Management" for details on where these guidelines can be obtained.</p>

Berries:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Strawberries	Grey mould (<i>Botrytis cinerea</i>)	All States	1.0 L/ha where spray volume is less than 1000 L/ha OR 100 mL/100 L water where spray volume equals or exceeds 1000 L/ha	1 day	This use is subject to a CropLife Australia fungicide resistance management strategy: 1. Apply a program of protectant fungicides during flowering. If conditions favour disease development during this period use Rovral® Aquaflo. 2. DO NOT apply more than two successive sprays of Rovral® Aquaflo (or other Group 2 Fungicide).
RATE In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Special Instructions for Tree Crops/Vines section.					CRITICAL COMMENTS For all use in the following table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Refer to the Special Instructions for Tree Crops/Vines section.
Blueberries	Grey mould (<i>Botrytis spp.</i>)	NSW, Qld, Tas, WA only	100 mL/100 L water		Apply every 10 to 14 days from flowering.
Raspberries	Grey mould (<i>Botrytis cinerea</i>)	All States			Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.
Youngberries	Grey mould (<i>Botrytis cinerea</i>)				Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.
Boysenberries	Grey mould (<i>Botrytis cinerea</i>)				Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.

Vegetables:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Celery	Sclerotinia rot (pink rot) (<i>Sclerotinia sclerotiorum</i>)	All States	1.0 L/ha where spray volume is less than 1000 L/ha	1 day	Commence spraying 1 to 2 weeks post-transplanting and then every 2 to 3 weeks. Use only five sprays.
Lettuces	Sclerotinia rot (drop) (<i>Sclerotinia sclerotiorum</i> , <i>Sclerotinia minor</i>)		OR 100 mL/100 L water where spray volume equals or exceeds 1000 L/ha	7 days	Spray should be directed to the stems at ground level and to the underside of lower leaves. This use is subject to a CropLife Australia fungicide resistance management strategy: 1. Apply Rovral® Aquaflo as a seedling drench soon after emergence. 2. Apply a protectant fungicide as a high volume foliar spray before planting out, then Rovral® Aquaflo immediately after planting. 3. Maintain cover with protectant fungicide sprays at 7-10 day intervals. 4. If weather conditions favour Botrytis infection, tank mix the protectant with Rovral® Aquaflo. 5. DO NOT apply Rovral® Aquaflo (or other Group 2 Fungicides) more than four
	Grey mould (<i>Botrytis spp.</i>)	Tas, WA only			

					times per season, irrespective of the target disease.
Potatoes	Sclerotinia rot (<i>Sclerotinia sclerotiorum</i>)	All States	500 ml to 1.0 L/ha where spray volume is less than 1000 L/ha OR 50 to 100 mL/100 L water where spray volume equals or exceeds 1000 L/ha	Nil	Apply 2 sprays, once immediately before and once immediately after hilling-up. For most effective treatment, concentrate the spray at the base of the stems and surrounding soil surface, where the fungus is active. Use the higher rate where disease is severe.
	Target spot, (early blight) (<i>Alternaria solani</i>)				Ensure thorough coverage to the whole plant. Treatment is generally not required until after flowering. Use the higher rate where disease is severe. Limit the use of Rovral® Aquaflor to periods when conditions favour disease development. This use is subject to a CropLife Australia fungicide resistance management strategy: The number of consecutive applications of Group 2 fungicides permitted is limited. Refer to the CropLife Australia Resistance Management Guidelines. See the "General Instructions - Resistance Management" for details on where these guidelines can be obtained.
	Hypocotyl rot (black scurf) (<i>Rhizoctonia solani</i>)		400 mL / tonne seed material		Rovral® Aquaflor will protect emerging shoots from hypocotyl rot, improving overall germination. Rovral® Aquaflor may also reduce occurrence of black scurf on the harvested potatoes. Ensure good coverage of seed material and planting furrow. This can be achieved by applying Rovral® Aquaflor as a fine spray to the seed at the time of planting using spray equipment mounted on the planter, and nozzles located at three points on each planter row to ensure uniform coating of the seed. DO NOT plant into waterlogged soil. A minimum water volume of 80 L/tonne seed should be used.
Tomatoes	Sclerotinia rot (<i>Sclerotinia sclerotiorum</i>)	Qld, NSW, Tas, SA, WA only	1.0 L/ha where spray volume is less than 1000 L/ha OR 100 mL/100 L water where spray volume equals or exceeds 1000 L/ha	7 days	Spray at 14-day intervals from transplanting and throughout the period of disease pressure.
	Grey mould (<i>Botrytis</i> spp.)	All States			Commence spraying 3 to 4 weeks after transplanting or at the onset of disease. Repeat treatment at 14-day intervals or when conditions favour spread of the disease, i.e. at trimming or deleafing. This use is subject to a CropLife Australia fungicide resistance management strategy: 1. Alternate or tank mix Rovral® Aquaflor with a protectant such as chlorothalonil. Avoid applying two Rovral® Aquaflor (or other Group 2 fungicide) sprays in succession, unless tank mixed with a protectant. 2. DO NOT apply more than four Rovral® Aquaflor (or other Group 2 fungicide) sprays in a season.

	Target spot (early blight) (<i>Alternaria solani</i>)	Qld, Tas, WA, NT only			Commence spraying 1 week post-transplanting. Use adequate water to give thorough coverage of the plants. Use high volume spray equipment. This use is subject to a CropLife Australia fungicide resistance management strategy: 1. Limit the use of Rovral® Aquaflor to periods when conditions favour disease development. 2. DO NOT apply more than four Rovral® Aquaflor (or other Group 2 fungicide) sprays in one season. Apply no more than two consecutive sprays of a Group 2 fungicide.
--	--	-----------------------	--	--	--

Field Crops:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Lucerne	Lucerne leaf spot (<i>Stemphylium botryosum</i>)	Qld, WA only	250 mL to 500 mL/ha where spray volume is less than 1000 L/ha OR 25 to 50 mL/100 L water where spray volume equals or exceeds 1000 L/ha	7 days	Spray every 10 to 14 days when cool, damp weather favours the disease. Use the higher rate under conditions of high disease pressure.
	Leptosphaerulina leaf spot (<i>Leptosphaerulina trifolii</i>)		Apply in at least 300 L water/ha every 10 to 14 days when cool, damp weather favours the disease. Use the higher rate under conditions of high disease pressure.		
Peanuts	Sclerotinia rot, (<i>Sclerotinia sclerotiorum</i> , <i>Sclerotinia minor</i>)		1.0 L/ha OR 220 mL/100 L water (spot application)	12 days	Apply when disease first appears. Repeat if necessary. Use a high water volume to ensure good coverage of foliage and stem at ground level. DO NOT mix Rovral® Aquaflor with a foliar fungicide due to the different target positions on the plant.
Soybeans	Black leaf blight (<i>Arkoala nigra</i>)	NSW, WA only	1.0 L/200 to 400 L water/ha	7 weeks	If disease is present on leaves apply an initial spray at early pod set (pods approximately 5 mm long). An additional spray 14 days later may be required if wet seasonal conditions prevail.

Ornamentals and Recreational Turf:

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Ornamentals	Botrytis blight (<i>Botrytis cinerea</i>)	All States	100 mL/100 L water	Spray at 14 day intervals commencing when the disease first becomes apparent and continuing until conditions no longer favour the disease. Spraying saintpaulia and poinsettia flowers may result in some petal scorch. Tepid water should be used and wet plants protected from direct sunlight. This use is subject to a CropLife Australia fungicide resistance management strategy: DO NOT apply more than two consecutive sprays of a Group 2 fungicide.

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Recreational turf	Brown patch (<i>Rhizoctonia solani</i>) Dollar spot (<i>Sclerotinia homoeocarpa</i>)	Qld, NSW, Tas, SA, WA only	60 to 90 mL/10 to 150 L of water per 100 m ²	Curative Programme: Repeat treatment after 14 days. Use the higher rate under conditions of severe disease pressure. For Preventative Treatment: Apply as a monthly spray program. This use is subject to an AFRAC anti-resistance strategy. To reduce the possibility of resistance build-up to Ippon Turf DO NOT apply more than 2 consecutive sprays of Rovral® Aquaflo or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Brown patch (<i>Curvularia spp.</i>), (<i>Dresch/era spp.</i>), (<i>Fusarium culmorum</i>)	Vic only	90 mL / 10 to 150 L water per 100 m ²	
	Fusarium patch (<i>Fusarium nivale</i>)	NSW, Vic, Tas, SA, WA only	90 mL / 10 to 150 L water per 100 m ²	Curative Programme: Repeat treatment after 14 days. Use the higher rate under conditions of severe disease pressure.
	Leaf spot (<i>Dreschlera spp.</i>)	Qld, NSW, Vic, SA, WA only	60 mL / 10 to 150 L water per 100 m ²	For Preventative Treatment: Apply as a monthly spray program. This use is subject to an AFRAC anti-resistance strategy. To reduce the possibility of resistance DO NOT apply more than two consecutive sprays of Rovral® Aquaflo or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Spring dead spot (<i>Leptosphaeria spp.</i>) [^]	NSW, SA only	65 mL / 10 to 150 L water per 100 m ²	For Preventative Treatment: Apply as a monthly soil drench throughout the year. Water in thoroughly after application.
	Spring dead spot (<i>Helminthosporium spp.</i>)	Vic, WA only	65 mL / 10 to 150 L water per 100 m ²	[^] When used in a preventative programme this product will limit the development of <i>Leptosphaeria</i> and improve the rate of turf recovery. To reduce the possibility of resistance DO NOT apply more than two consecutive sprays of Rovral® Aquaflo or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group
	Helminthosporium disease affecting leaf, crown and roots: (Black and white <i>Helminthosporium</i> spot: <i>Bipolaris spp.</i> <i>Dreschlera spp.</i>)	Qld, NSW, Vic, SA, WA only	45 mL / 10 to 150 L water per 100 m ²	For Curative Treatment: Apply as a foliar spray and repeat application after 7 to 14 days. For white spot symptoms on couch grass (<i>Cynodon dactylon</i>) only, use higher after volumes of 100 to 150 litres of water per 100 m ² . For Preventative Treatment: Apply as a high volume foliar spray. Repeat applications at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. To reduce the possibility of resistance DO NOT apply more than 2 consecutive sprays of Rovral® Aquaflo or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS (WHP):

Almonds, macadamias, mandarins, pome fruit, potatoes and stone fruit: NOT REQUIRED WHEN USED AS DIRECTED.

Boysenberries, celery, passionfruit, raspberries, strawberries, blueberries and youngberries: **DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.**

Grapes, kiwifruit, lettuce and tomatoes: **DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.**

Peanuts: **DO NOT HARVEST FOR 12 DAYS AFTER APPLICATION.**

Soybeans: **DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION.**

Lucerne: **DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 7 DAYS OF TREATMENT.**

Turf: **DO NOT** GRAZE TREATED AREAS OR FEED TURF CLIPPINGS FROM TREATED AREAS TO ANIMALS INCLUDING POULTRY.

EXPORT OF TREATED PRODUCE

Growers should note that MRLs or import tolerances may not exist in all markets for produce treated with Rovral® Aquaflor.

If you are growing produce for export, please check with FMC Australasia Pty Ltd, for the latest information on MRLs and import tolerances BEFORE using Rovral® Aquaflor.

INDEX

CROP	DISEASE (common name)	DISEASE (scientific name)	Page No.
Almonds	Blossom blight, brown rot	<i>Monilinia</i> spp., <i>Sclerotinia</i> spp.	3
Apricots	Blossom blight, brown rot	<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>	4
	Transit rot	<i>Rhizopus</i> spp.	4
Apples	Storage rots	<i>Penicillium</i> spp., <i>Botrytis</i> spp., <i>Gloeosporium</i> spp.	4
Blueberries	Grey mould	<i>Botrytis</i> spp.	5
Boysenberries	Grey mould	<i>Botrytis cinerea</i>	5
Celery	Sclerotinia rot (pink rot)	<i>Sclerotinia sclerotiorum</i>	5
Cherries	Blossom blight, brown rot	<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>	4
	Transit rot	<i>Rhizopus</i> spp.	4
Grapes	Grey mould	<i>Botrytis cinerea</i>	3
Kiwifruit	Botrytis blight	<i>Botrytis</i> spp.	3
Lettuces	Sclerotinia rot {drop}	<i>Sclerotinia sclerotiorum</i> , <i>Sclerotinia minor</i>	5
	Grey mould	<i>Botrytis</i> spp.	5
Lucerne	Lucerne leaf spot	<i>Stemphylium botryosum</i>	7
	Leptosphaerulina leaf spot	<i>Leptosphaerulina trifolii</i>	7
Macadamias	Botrytis blight	<i>Botrytis</i> spp.	3
Mandarins	Alternaria: leaf spot (brown spot)	<i>Alternaria alternata</i>	3
Nectarines	Blossom blight, brown rot	<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>	4
	Transit rot	<i>Rhizopus</i> spp.	4
Ornamentals	Botrytis blight	<i>Botrytis cinerea</i>	7
Passionfruit	Alternata spot (brown spot)	<i>Alternaria</i> spp., <i>Alternaria passiflorae</i>	3
Peaches	Blossom blight, brown rot	<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>	4
	Transit rot	<i>Rhizopus</i> spp.	4
Peanuts	Sclerotinia rot	<i>Sclerotinia sclerotiorum</i> , <i>Sclerotinia minor</i>	7
Pears	Storage rots	<i>Penicillium</i> spp., <i>Botrytis</i> spp., <i>Gloeosporium</i> spp.	4
Plums	Blossom blight, brown rot	<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>	4
	Transit rot	<i>Rhizopus</i> spp.	4
Potatoes	Sclerotinia rot	<i>Sclerotinia sclerotiorum</i>	6
	Target spot (early blight)	<i>Alternaria solani</i>	6
	Hypocotyl rot (black scurf)	<i>Rhizoctonia solani</i>	6
Raspberries	Grey mould	<i>Botrytis cinerea</i>	5
Soybeans	Black leaf blight	<i>Arkoola nigra</i>	7
Strawberries	Grey mould	<i>Botrytis cinerea</i>	5
Tomatoes	Sclerotinia rot	<i>Sclerotinia sclerotiorum</i>	6
	Grey mould	<i>Botrytis cinerea</i>	6
	Target spot (early blight)	<i>Alternaria solani</i>	6
Young berries	Grey mould	<i>Botrytis cinerea</i>	5

NOTICE TO BUYER

To the extent permitted by the Competition and Consumer Act (2010) or any relevant legislation of any State or Territory (the "Legislation") all conditions and warranties and statutory or other rights of action, whether arising in contract or tort or whether due to the negligence of FMC or Seller, which buyer or any other user may have against FMC or Seller are hereby excluded provided however that any rights of the buyer pursuant to non-excludable conditions or warranties of the Legislation are expressly preserved. FMC hereby gives notice to buyer and other users that to the extent permitted by the Legislation it will not accept responsibility for any indirect or consequential loss of whatsoever nature arising from the storage, handling or use of this product. Where permitted by the Legislation FMC's liability shall in all circumstances be limited to the replacement of the product, or a refund of the purchase price paid therefor.

The product must be used and applied strictly in accordance with the label instructions and other directions for use. It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of FMC or the Seller. Buyer accepts these risks.

The information set forth herein is furnished free of charge and is based on technical data that FMC believes to be reliable. It is intended for use by persons having technical skill at their own discretion and risk. FMC makes no warranties, expressed or implied, and assumes no liability in connection with any of this information. Nothing herein is to be taken as license to operate under or a recommendation to infringe on any patents.

FMC and Rovral® are trademarks of FMC Corporation or an affiliate.

© 2021 FMC Corporation. All rights reserved.

*Non FMC Trademark

APVMA Approval Number: 45725/132407

FMC Australasia Pty Ltd.

A.B.N. 45 095 326 891

Level 2, Building B

12 Julius Ave, North Ryde NSW 2113

Phone: 1800 066 355

www.fmccrop.com.au



SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111

ALL HOURS – AUSTRALIA WIDE
