

PREDICADE[®]

HERBICIDE

The all-in-one solution for grassy and broadleaf weeds in wheat

What is Predicade[™] herbicide?

Predicade[™] herbicide provides broad-spectrum weed control in spring, durum and winter wheat, with convenient handling and five active ingredients for exceptional resistance management.

How does Predicade[™] herbicide help my farm business?

Excellent broad-spectrum performance. Predicade[™] herbicide targets more than 30 of Western Canada's key grassy and broadleaf weeds and excels on the toughest of these, including Group 1-resistant wild oats. Many growers who missed their spring burnoff depend on Predicade[™] herbicide to get their wheat clean.

Next-level resistance management. With five active ingredients from two herbicide Groups, you can be confident you're doing your utmost to manage the threat of resistance and keep your future herbicide options open.

Convenience in mixing and application. One case of Predicade[™] herbicide provides everything you need for strong grassy and broadleaf weed control. One case treats 40 acres. It's that simple, which helps minimize the risk of mixing and application errors.

INCROP

Quick facts:

Crop staging:

3-leaf to 6-leaf with 3 tillers, prior to jointing

Water volume:

Minimum 5 gal/ac (50 L/ha)

Packaging:

One (1) case treats 40 acres

Rainfastness:

Two (2) hours

Predicade[™] herbicide provides exceptional weed control.



Sprayer miss

Source: Photo taken 21 days after application.
Oakville, MB.

When and how is Predicade™ herbicide applied?

Predicade™ herbicide can be applied from the 3-leaf stage to 6-leaf with 3 tillers, but prior to jointing (presence of first node).

How does Predicade™ herbicide affect my crop rotation?

Re-cropping flexibility is a key benefit. The following crops can be seeded the year after applying Predicade™ herbicide: alfalfa, barley, canola, field corn, flax, dry beans, lentils, mustard, oats, peas, soybeans, spring and durum wheat or sunflowers.

Control of Shepherd's Purse.



Source: Photo taken 21 days after application.
Oakville, MB.

Crops

- Wheat (durum, spring, winter)

Weeds

Broadleaf weeds

- Annual smartweed (green smartweed, lady's-thumb)
- Canada thistle¹ (top growth control)
- Cleavers (1-6 whorls)
- Common chickweed (1-6 leaf)
- Cow cockle
- Dandelion (< 15 cm in diameter)
- Flixweed
- Hemp-nettle
- Kochia (including Groups 2 & 9-resistant biotypes; seedling to 8-leaf)
- Lamb's-quarters
- Narrow-leaved hawk's-beard
- Night flowering catchfly
- Pale smartweed (1-6 leaf)
- Redroot pigweed
- Round-leaved mallow (1-5 leaf)
- Russian thistle
- Scentless chamomile (< 10 cm)
- Shepherd's-purse (1-6 leaf)
- Sow thistle, perennial
- Stinkweed
- Stork's-bill (1-6 leaf)
- Volunteer canola (2-4 leaf) including Clearfield® varieties
- Volunteer flax (< 12 cm)
- White cockle (< 10 cm)
- Wild buckwheat
- Wild mustard

Grassy weeds

- Barnyard grass
- Green foxtail
- Japanese brome² (1-6 leaf)
- Persian dandel¹
- Volunteer canary seed
- Wild oats
- Yellow foxtail¹

¹ Suppression

² Control of spring-germinated Japanese brome. Suppression of overwintered Japanese brome. Best results are obtained after a pre-seed or burnoff application with a glyphosate herbicide.

Questions? Ask your retailer about Predicade™ herbicide today.

Always read and follow label instructions.
Member of CropLife Canada.

FMC and Predicade are trademarks or service marks of FMC Corporation and/or an affiliate.
©2019 FMC Corporation. All rights reserved. 64602 - 2/19



@FMCAgCanada

FMCcrop.ca | 1-833-362-7722