



Aim[®] EC herbicide can turn harvest conditions in your favour.

Add Aim EC herbicide to your pre-harvest application to control weeds and dry-down plant material to make harvest faster and easier.

How can Aim EC herbicide make my harvest better?

Some years, growing conditions give weeds and overgrown plant material the upper hand, making harvest tougher and more time-consuming than it needs to be. An application of Aim EC herbicide can turn harvest conditions in your favour.

Fast-acting. Adding Aim EC herbicide to your pre-harvest application helps dry down plant material and provides better, faster control of a wide range of troublesome broadleaf weeds for an easier harvest.

Powerful Group 14 broadleaf activity. Aim EC herbicide is effective on a range of troublesome broadleaf weeds that can hinder harvest, including many that are resistant to other chemistries.

Convenient. Aim EC herbicide features an easy-to-handle formulation and low use rates, to get you out there quicker and done sooner.

Use with or without Glyphosate. Harvest aids, like glyphosate are under increased scrutiny from buyers limiting growers choice to help with crop and weed dry down.

Now available in 1.2 and 4.8 L jug sizes for easier use.

Aim EC herbicide vs Heat[®]

Application: September 5. Photo taken: 8 days after application.



No harvest aid

Aim EC herbicide

Heat[®]

Source: Josephburg, Alberta, 2017

Aim EC herbicide 47mL/ac +
1 REL Weathermax + 0.25% v/v Ag-Surf™

Heat[®] herbicide 30 ac/jug +
1 REL Weathermax + 1% merge

**HARVEST AID
GROUP
14**

Quick facts:

Formulation:
EC – emulsifiable concentrate

Rates:
Apply at 47 mL per acre
1.2 L jug = 25 ac/jug
2.0 L jug = 42 ac/jug
4.8 L jug = 102 ac/jug

Water volume:
10 gal/ac (100 L/ha)

Rainfastness:
1 hour

Timing:
Pre-seed burndown
Chemfallow

Harvest aid
Post-harvest

Crops:
Barley
Beans
Oat
Peas
Potatoes
Soybeans
Wheat

* See label for potato rate.

When do I apply Aim® EC herbicide in a pre-harvest application with glyphosate?

Pre-Harvest Interval

Crop/Crop Group/Crop Subgroup	PHI (Days Before Harvest)
Sorghum (harvest aid)	3
Barley (harvest aid)	3
Millet (harvest aid)	3
Oats (harvest aid)	3
Triticale (harvest aid)	3
Wheat (harvest aid)	3
Soybean (harvest aid)	3
Potato (harvest aid)	7
Dried shelled peas and beans (harvest aid)	3

Wheat, barley and oats. Apply when grain moisture is less than 30% (hard dough stage; thumbnail impression remains on seed).

Peas and beans. Apply when grain moisture is less than 30%. Apply to dry beans when 80% to 90% of bean leaves have fallen and pods are mature (yellow to brown in color). Apply to dry pea when majority (75% to 80%) of the pods are brown, the bottom pods are ripe and the pea seeds have detached from the pods.

Soybeans. Apply when crop is less than 30% moisture, has lost 80% to 90% of leaves and 80% of pods are brown.

What if the crop and weeds are extra thick?

For situations where canopies are thicker, making weed coverage more challenging, FMC recommends using an adjuvant with your Aim EC herbicide plus glyphosate application.

- Agral® 90 or Ag-Surf™ at 0.25% v/v (0.25 litres per 100 litres of spray solution)
- Merge at 1% v/v (1 litre per 100 litres of spray solution)



Untreated Dry Beans



Aim EC herbicide 47mL/ac 6 DAA

Source: Plum Coulee, MB, 2019



FMC Canada will donate 10 cents for every acre of our post-harvest products sold between July 15 and October 31, 2020. That includes not only Aim EC herbicide, but also new Intruvix™ herbicide, Focus® herbicide, Express® brand herbicides and our fall-applied PrecisionPac® blends.

Questions? Ask your retailer about Aim EC herbicide today.

Terms and Conditions Apply. Contact your local retailer for details.
Always read and follow label directions. Member of CropLife Canada.
FMC, the FMC logo and Aim are trademarks of FMC Corporation or an affiliate.
© 2020 FMC Corporation. All rights reserved. 72501 - 7/20
Agral® (Syngenta); Ag-Surf™ (IPCO); Heat® (BASF)

    @FMCAgCanada

FMCcrop.ca | 1-833-362-7722