



BARRICADE[®]

II

HERBICIDE

Consistently powerful performance on your toughest broadleaf weeds

What is Barricade[®] II herbicide?

Barricade II herbicide controls more than 20 broadleaf weeds in wheat, barley and oats, with three active ingredients from two herbicide Groups for sound resistance management.

How does Barricade II herbicide help my farm business?

Outstanding control of the toughest weeds. Control of narrow-leaved hawk's beard, cleavers, kochia, shepherd's-purse, wild buckwheat and more.

Excellent risk management tool. This herbicide contains three active ingredients from two Groups, making it a sound component of a long-term resistance management strategy.

Application and rotational flexibility. Barricade II herbicide can be sprayed as early as the 2-leaf stage and as late as flag leaf. It has a wide range of registered and supported grass tank-mix partners. Most crops can be planted the following year.

Tough on weeds, gentle on crops. More than 99% of growers who use Barricade II herbicide report they're satisfied with the product's effectiveness. More importantly, crop safety is not compromised.

Proven formulation. Barricade II herbicide is powered by Solumax[®] soluble granules for effective and consistent weed control. Because sprayer cleanout is easy, there's less risk to subsequently sprayed crops like peas or canola.

IN-CROP

Quick facts:

Application timing:
2-leaf to flag leaf stage
(spring wheat, durum, barley)
3-tiller to before the flag leaf
(winter wheat)

Water volume:
Minimum 5 gal/ac (50 L/ha)

Packaging:
One (1) case treats 40 acres
Registered surfactant required
at 0.2% v/v

Rainfastness: 1 hour

Efficacy on Narrow-leaved hawk's-beard



Day of application



47 days after application

Efficacy on Cleavers



Day of application



29 days after application

Source: Vegreville, AB, 2014

Which grass herbicides can be tank-mixed with Barricade® II herbicide?

- Assert®
- Axial®
- Clodinafop
- Everest® 3.0
- GP Herbicide
- MCPA
- Puma® Advance
- Simplicity™ brands
- Traxos®
- Varro®
- 2,4-D Ester



Crops

- Spring wheat
- Durum wheat
- Winter wheat
- Spring barley
- Oats*

Weeds

- Canada thistle¹
- Cleavers (1 to 6 whorls)
- Common chickweed
- Cow cockle
- Dandelion² (spring and fall rosettes up to 15 cm)
- Flixweed
- Hemp-nettle
- Kochia
- Lamb's-quarters
- Narrow-leaved hawk's-beard (<10cm)
- Night-flowering catchfly (<10cm)
- Redroot pigweed
- Russian thistle
- Round-leaved mallow (1-5 leaf)
- Scentsless chamomile²
- Shepherd's purse (up to 20 cm)
- Smartweed (Lady's-thumb)
- Sow thistle (perennial)
- Stinkweed
- Stork's-bill
- Volunteer canola²
- Volunteer flax (12 cm in height)
- White cockle²
- Wild buckwheat
- Wild mustard

¹ Suppression.

² Controlled when Barricade II herbicide is tank-mixed with MCPA.

*when tank-mixed with MCPA

Questions? Ask your retailer about Barricade II herbicide today.

Always read and follow label instructions.
Member of CropLife Canada.

FMC, the FMC logo and Barricade are trademarks of FMC Corporation or an affiliate. Assert is a trademark of Nufarm. Axial, Sierra, and Traxos are trademarks of Syngenta. Everest is a trademark of Arysta LifeScience. Puma and Varro are trademarks of Bayer CropScience. Simplicity is a trademark of Corteva.
© 2020 FMC Corporation. All rights reserved. 72186 6/20





@FMCAgCanada
FMCcrop.ca | 1-833-362-7722