


FALL IS THE TIME FOR WEED CONTROL

3 great reasons why fall weed control is the right choice for an easier spring.

1. Using pre-harvest weed control not only helps with harvest, but it leaves your fields clean in early fall.
2. Fall temperatures trigger nutrient movement to the roots of perennials. A systemic herbicide applied post-harvest takes advantage of this movement into the root system, allowing for better control.
3. Late fall applications of a herbicide that creates a soil barrier after moisture activation improves the consistency of early germinating weed control next spring.








THINKING
FALL WEED
CONTROL?
THINK FMC.

A TIMELY FALL APPLICATION SETS YOU UP FOR A QUICK START AND A PROFITABLE CROP THE FOLLOWING YEAR. HERE'S A LOOK AT THE BENEFITS OF FMC'S FALL WEED CONTROL PORTFOLIO.


MORE WAYS TO NAIL YOUR FALL WEED CONTROL

AIM[®] EC *Turn harvest conditions in your favour with quicker dry down.*
HERBICIDE

-  Pre-harvest. Apply when crop moisture is <30%
-  Barley, beans, oats, peas, potatoes, soybeans & wheat
-  Resistant kochia, cleavers, vol. canola, waterhemp, Russian thistle, shepherd's purse, round-leaved mallow
-  47 mL/ac
-  Can be used with or without glyphosate
Check label for crop specific pre-harvest intervals








Untreated Dry Beans




Aim[®] EC herbicide 47mL/ac 6 DAA

Source: Plum Coulee, MB, 2019

FOCUS[®] *NOW registered for fall application.*
HERBICIDE

-  Post-harvest
-  Field corn, lentils, spring and winter wheat, soybeans
NEW field peas and sunflowers
-  Emerged winter annuals such as stinkweed and mustard
Early season control of wild oats the following spring
-  33 ac/jug at 3-7% organic matter
40 ac/jug at 1-3% organic matter
-  Faster burnoff when added to glyphosate



Source: FMC Global Research Station, Hanley, SK


EXPRESS[®] BRAND *Powerful burnoff from shoots to roots.*
HERBICIDES

Which one best suits your post-harvest weed control needs?


Express[®] FX herbicide for resistance management and control of resistant kochia, dandelion and more

Express[®] PRO herbicide for extended control of dandelion, vol. canola, narrow-leaved hawk's-beard and cleavers

Express[®] SG herbicide for maximum rotational flexibility

 Must be tank-mixed with glyphosate

 80 ac/jug



Source: Controlled growth room environment, 25 days after application, Dr. François Tardif, Peter Smith, University of Guelph, Plant Agriculture Department, June, 2013.



October, 2013, Manitoba. Express SG herbicide + glyphosate 19 days after application.

2 more reasons to choose FMC herbicides this fall.

1. Once again, we are teaming up with local retailers to donate \$0.10 per acre from our 2021 fall herbicide sales, July 15 to October 31, 2021, to organizations that serve local communities. It could be to support sports, arts, seniors or something else – if they help people, we'll help them.
2. **The ACE Resistant Weeds program runs until October 31st, 2021.** Any unmatched purchases of Aim[®]EC, Command[®] 360 ME or Express[®] brands will be eligible for up to a \$5/acre rebate on your fall weed control.

CLEAN FIELDS *Community Yields*

ACE
RESISTANT WEEDS