



# Raising the bar even higher on broadleaf weed control

## What is Barricade® M herbicide?

Barricade® M herbicide provides consistently powerful performance against broadleaf weeds in wheat (spring, winter, durum) and barley, with four active ingredients to manage resistance and outstanding re-cropping flexibility the following year.

## How does Barricade® M herbicide help my farm business?

**Outstanding control of the toughest weeds.** Barricade® M herbicide provides excellent control of hard-to-kill weeds including perennial sow thistle, wild buckwheat, lamb's-quarters, pigweeds, dandelion and Canada thistle<sup>1</sup> by working systemically to control the entire weed.

**Excellent resistance management tool.** With four active ingredients from two different herbicide Groups (2 and 4), Barricade® M herbicide takes on the toughest broadleaves you face and helps manage weed resistance.

**Application flexibility.** Barricade® M herbicide can be sprayed as early as 3-leaf and as late as just before the flag leaf, with full crop safety.

**Consistent performance.** Powered by Solumax® soluble granules, Barricade® M herbicide provides effective and consistent weed control. Once you're done, Solumax® ensures easy sprayer cleanout to reduce risk to subsequently sprayed crops like canola, corn or soybeans.

**Keeps your cropping options open.** The following crops can be seeded the year after applying Barricade® M herbicide: alfalfa, barley, canola, corn, dry beans, flax, forage grasses, lentils, mustard, oats, peas, potatoes, rye, soybeans, sugar beets, sunflowers and wheat.

GROUPS  
2 | 4

### Quick facts:

**Application timing:**  
3-leaf to just before flag leaf

**Water volume:**  
10 gal/ac (100 L/Ha)

**Packaging:**  
One (1) case treats 40 acres

**Rainfastness:**  
1 hour

**Surfactant is recommended**

### Efficacy on Dandelions



16 Days after application



47 Days after application

### Efficacy on Sow-Thistle



16 Days after application



47 Days after application

## When and how is Barricade® M herbicide applied?

The window of application for Barricade® M herbicide is from the 3-leaf stage to just before flag leaf. One case of Barricade® M herbicide treats 40 acres. Each case contains a dry herbicide in a 486-gram jug and two liquid herbicide jugs. Add a registered non-ionic surfactant (NIS) at 2 L per 1000 L of spray solution (0.2% v/v).



### Crops

- Spring wheat
- Durum wheat
- Winter wheat
- Spring barley

### Weeds

- Canada thistle<sup>1</sup>
- Cleavers (1 to 6 whorls)
- Common chickweed
- Cow cockle
- Dandelion (spring and fall rosettes up to 15 cm)
- Flixweed
- Hemp-nettle
- Kochia
- Lamb's-quarters
- Narrow-leaved hawk's-beard (<10cm)
- Night-flowering catchfly (<10cm)
- Redroot pigweed
- Russian thistle
- Round-leaved mallow (1-5 leaf)
- Scentless chamomile
- Smartweed (Lady's-thumb)
- Stinkweed
- Stork's-bill
- Volunteer canola
- Volunteer flax (12 cm in height)
- White cockle
- Wild buckwheat
- Wild mustard

<sup>1</sup> Suppression.

**Questions? Ask your retailer about Barricade® M herbicide today.**

Always read and follow label instructions. Member of CropLife Canada.

FMC, Barricade and Solumax are trademarks or service marks of FMC Corporation and/or an affiliate.  
© 2019 FMC Corporation. All rights reserved. 5/19 - 64334

  @FMCAgCanada

FMCcrop.ca | 1-833-362-7722