



# Aim<sup>®</sup> EC herbicide can turn harvest conditions in your favour.

Add Aim EC herbicide to your pre-harvest glyphosate application to control weeds and dry-down plant material to make harvest faster and easier.

## How can Aim EC herbicide improve my harvest?

Some years, growing conditions give weeds the upper hand, making harvest tougher and more time-consuming than it needs to be. An application of Aim EC herbicide can turn harvest conditions in your favour.

**Fast-acting.** Adding Aim EC herbicide to your pre-harvest application helps dry down plant material and provides better, faster control of a wide range of troublesome broadleaf weeds for an easier harvest.

**Powerful Group 14 broadleaf activity.** Aim EC herbicide is effective on a range of difficult broadleaf weeds that can hinder harvest, including many that are resistant to other chemistries.

**Convenient.** Aim EC herbicide features an easy-to-handle formulation and low use rates, to get you out there quicker and done sooner.

**Use with or without Glyphosate.** Harvest aids, like glyphosate, are under increased scrutiny from buyers limiting growers choice to help with crop and weed dry down. When applied without glyphosate, ensure that a registered surfactant is used. When Aim EC herbicide is applied with glyphosate, a registered surfactant is recommended where crop canopies are thicker.

Now available in 1.2 and 4.8 L jug sizes for easier use.

## Aim EC herbicide vs Heat<sup>®</sup> LQ

Application: September 5. Photo taken: 8 days after application.



Source: Josephburg, Alberta, 2017

Aim EC herbicide 47mL/ac +  
1 REL Weathermax + 0.25% v/v Ag-Surf<sup>™</sup>

Heat<sup>®</sup> LQ herbicide 30 ac/jug +  
1 REL Weathermax + 1% merge

**HARVEST AID  
GROUP  
14**

### Quick facts:

**Formulation:**

EC – emulsifiable concentrate

**Rates:**

Apply at 47 mL per acre

1.2 L jug = 25 ac/jug

4.8 L jug = 102 ac/jug

**Water volume:**

Minimum of 10 gal/ac (100 L/ha)

**Rainfastness:**

1 hour

**Timing:**

Pre-seed burndown

Chemfallow

**Harvest aid**

**Key Crops:**

Barley

Dry beans

Oat

Peas

Potatoes\*

Soybeans

Wheat

\* See label for potato rate and application information.

**Registered Surfactants:**

Agrial<sup>®</sup> 90 or Ag-Surf<sup>™</sup> at 0.25% v/v  
(0.25 litres per 100 litres of spray solution)

Merge at 1% v/v (1 litre per 100 litres of spray solution)

When Aim EC herbicide is applied with glyphosate, a registered surfactant is recommended where crop canopies are thicker. When applied without glyphosate, ensure that a registered surfactant is used.

## When do I apply Aim® EC herbicide in a pre-harvest application with glyphosate?

**Wheat, barley and oats.** Apply when grain moisture is less than 30% (hard dough stage; thumbnail impression remains on seed).

**Peas and beans.** Apply when grain moisture is less than 30%. Apply to dry beans when 80% to 90% of bean leaves have fallen and pods are mature (yellow to brown in color). Apply to dry pea when majority (75% to 80%) of the pods are brown, the bottom pods are ripe and the pea seeds have detached from the pods.

**Soybeans.** Apply when crop is less than 30% moisture, has lost 80% to 90% of leaves and 80% of pods are brown.

### Pre-Harvest Interval

When using Aim EC with glyphosate, refer to the glyphosate label for specific pre-harvest interval recommendations.

**When using Aim EC alone, without glyphosate, the pre-harvest intervals to the right apply.**

The use of Aim EC Herbicide will not speed up the maturity of green crops.

Always check with your grain buyer and/or grain contracts prior to using pre-harvest herbicides.

For more information see: [keepitclean.ca](http://keepitclean.ca)

Crop/Crop Group/Crop Subgroup	PHI (Days Before Harvest)
Sorghum (harvest aid)	3
Barley (harvest aid)	3
Millet (harvest aid)	3
Oats (harvest aid)	3
Triticale (harvest aid)	3
Wheat (harvest aid)	3
Soybean (harvest aid)	3
Potato (harvest aid)	7
Dried shelled peas and beans (harvest aid)	3

## What if the crop and weeds are extra thick?

For situations where canopies are thicker, making weed coverage more challenging, FMC recommends using an adjuvant with your Aim EC herbicide plus glyphosate application.

- Agral® 90 or Ag-Surf™ at 0.25% v/v (0.25 litres per 100 litres of spray solution)
- Merge at 1% v/v (1 litre per 100 litres of spray solution)

### Aim EC Harvest Aid on Dry Beans



Untreated Dry Beans



Aim EC herbicide 47mL/ac 6 DAA

Spray Date: August 29, 2020. Photos taken: September 4, 2020.



FMC Canada will donate 10 cents for every acre of our post-harvest products sold between July 15 and October 31, 2021. That includes not only Aim EC herbicide but also Express® brand herbicides, Focus® herbicide and our fall-applied PrecisionPac® blends.

## Questions? Ask your retailer about Aim EC herbicide today.

Terms and Conditions Apply. Contact your local retailer for details.  
Always read and follow label directions. Member of CropLife Canada.  
FMC, the FMC logo, Aim, Express, Focus and PrecisionPac are trademarks of FMC Corporation or an affiliate.  
© 2021 FMC Corporation. All rights reserved. 78749 - 7/21  
Agral® (Syngenta); Ag-Surf® (IPCO); Heat® (BASF)



[ag.FMC.com/ca](http://ag.FMC.com/ca) | 1-833-362-7722