

ONE PASS. ONE SOLUTION.

PrecisionPac® CS-100-12 herbicide controls a wide range of broadleaf and grassy weeds including exceptional wild oat control in spring, durum and winter wheat. PrecisionPac® CS-100-12 herbicide is a cost-effective option for growers with lighter broadleaf weed pressure.

Why PrecisionPac® herbicides? Dispensed in easy-to-handle, premeasured pouches for your exact weed spectrum and sprayer size, you buy only the amount you need. That means customized, top-performing weed control, great value and no leftovers to store!

The flexibility of PrecisionPac® herbicides gives you the power to fully customize your weed control.

The end result is:

-  High Performing Weed Control
-  Completely Customized Weed Control
-  Easy to Use
-  Right Product, Right Rate, Right Acre
-  Multiple Modes of Action
-  Pick Length Of Herbicide Activity
-  Reduced Waste
-  Constantly Evolving



    @FMCAgCanada
ag.FMC.com/ca | 1-833-362-7722

Always read and follow label directions. Member of CropLife Canada.

FMC, the FMC logo, Perimeter and PrecisionPac are trademarks of FMC Corporation or an affiliate. © 2021 FMC Corporation. All rights reserved. 79737 - 10/21

Liberate and LI 700 are trademarks of Loveland Products, Inc. Sentry is a trademark of Nufarm Agriculture, Inc. Agral is a trademark of a Syngenta Group Company. AgSurf is a trademark of Interprovincial Cooperative Limited. Enhance and Pro-Surf is a trademark of Norac Concepts Inc. HIActivate is a trademark of WinField Solutions, LLC.

FMC An Agricultural Sciences Company



PRECISIONPAC
APPLICATION INNOVATION

CS-100-12
Customizable,
flexible in-crop grass
and broadleaf control.

PRECISIONPAC® CS-100-12 Herbicide

Customizable control and flexible
application timing for in-crop grass
and broadleaf weed control.

Grassy weeds controlled

- Barnyard grass (1-5 leaf)
- Downy brome¹ (2-6 leaf, 4-tiller)
- Green foxtail¹ (1-5 leaf)
- Japanese brome (1-6 leaf)
- Persian darnel (1 leaf to 4 leaf, 2 tiller)¹
- Wild oats (up to 4-leaf, 2 tiller)
- Yellow foxtail (1-5 leaf)

Broadleaf weeds controlled

- Canada thistle¹ (up to 30 cm tall, pre-bud)
- Cleavers (up to 6 whorl)
- Common chickweed
- Corn spurry (up to 2 whorl stage, less than 10 cm height)
- Cow cockle (up to 8 leaf)
- Dandelion¹ (seedlings and overwintered rosettes less than or equal to 20 cm)
- Flixweed
- Green smartweed
- Hemp-nettle (1-8 leaf)
- Lamb's-quarters
- Narrow-leaved hawk's beard (rosette)
- Redroot pigweed (1-8 leaf)
- Round-leaved mallow (up to 6-leaf, less than 10 cm)
- Russian thistle
- Shepherd's purse (up to 30 cm tall)
- Smartweed (lady's-thumb, 1-5 leaf)
- Stinkweed (up to 30 cm tall)
- Volunteer canola¹ (1-6 leaf)
- White cockle¹ (up to first flower, less than 20 cm stage)
- Wild buckwheat (1-6 leaf)
- Wild mustard



Crops:

Wheat (spring, durum and winter)



Timing:

Spring and durum wheat: 3 leaf to before flag leaf stage

Winter wheat: Apply in the spring when winter wheat is in the 2-7 leaf, 4 tiller stage



Crop rotation:

10 months: Field corn, sunflower, potatoes

11 months: Barley, brown mustard, canola, dry bean (of the species *Phaseolus vulgaris*), flax, canola quality Brassica juncea, lentils, oats, field peas, chickpea, spring wheat, soybean and yellow mustard or fields can be summerfallowed.



Application information:

Rates: 40.5 g/ac

Acres per grower bag:

Minimum: 20 acres; Maximum: 200 acres

Surfactant: Add a registered non-ionic surfactant (NIS) such as: Agral® 90, Liberate® Adjuvant, Ag-Surf® II, HiActivate® Non-ionic Liquid Spreader/Activator, Super Spreader Adjuvant, Enhance®, Pro-Surf® II or Sentry® at 2.5 L per 1000 L of spray solution.

Rainfastness: Two (2) hours

Water volume: 5-10 imp gal/ac

Mixing Instructions: Ensure that PrecisionPac® CS-100-12 herbicide is fully dissolved before adding tank-mix partners.

CS-100-12 herbicide is also registered for aerial application.

Grazing/Harvest Restrictions: Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.



Barnyard grass



Japanese brome



Yellow Foxtail



Wild oats



Narrow-leaved hawk's beard



Hemp nettle



Flix Weed



Lamb's quarters

¹ Suppression.