

ONE PASS. ONE SOLUTION.

PrecisionPac® CS-100-23235 herbicide combines our pyroxsulam and PP-23235 herbicide for an all-in-one combination that controls broadleaf and grassy weeds in spring and durum wheat. PrecisionPac® CS-100-23235 herbicide is an ideal product for the dark brown, black and grey soil zones.

Dispensed in easy-to-handle, premeasured pouches for your exact weed spectrum and sprayer size, you buy only the amount you need. That means customized, top-performing weed control, great value and no leftovers to store!

The flexibility of PrecisionPac® herbicides gives you the power to fully customize your weed control.

The end result is:

-  High Performing Weed Control
-  Completely Customized Weed Control
-  Easy to Use
-  Right Product, Right Rate, Right Acre
-  Multiple Modes of Action
-  Pick Length Of Herbicide Activity
-  Reduced Waste
-  Constantly Evolving



    @FMCAgCanada
ag.FMC.com/ca | 1-833-362-7722

Always read and follow label directions. Member of CropLife Canada.

FMC, the FMC logo, Perimeter and PrecisionPac are trademarks of FMC Corporation or an affiliate. © 2021 FMC Corporation. All rights reserved. 79737 - 10/21
Liberate and LI 700 are trademarks of Loveland Products, Inc. Sentry is a trademark of Nufarm Agriculture, Inc. Agral is a trademark of a Syngenta Group Company. AgSurf is a trademark of Interprovincial Cooperative Limited. Enhance and Pro-Surf are trademarks of Norac Concepts Inc. HIActivate is a trademark of WinField Solutions, LLC.

FMC | An Agricultural Sciences Company



PRECISIONPAC
APPLICATION INNOVATION

CS-100-23235
gives you
tailor-made,
one-pass control.

PRECISIONPAC[®] CS-100-23235 Herbicide

PrecisionPac[®] CS-100-23235 herbicide delivers premium grassy and broadleaf weed control for the most challenging fields.

Grassy Weeds Controlled

- Barnyard grass (1-5 leaf)
- Downy brome¹ (2-6 leaf, 4 tillers)
- Green foxtail¹ (1-5 leaf)
- Japanese brome (1-6 leaf)
- Persian darnel¹ (1-4 leaf, 2 tiller)
- Wild oats (up to 4 leaf, 2 tiller)
- Yellow foxtail (1-5 leaf)

Broadleaf Weeds Controlled

- Annual smartweed (green smartweed, lady's-thumb)
- Annual sow thistle³
- Canada thistle¹
- Cleavers (1-6 whorl)
- Common chickweed (up to 10 cm)
- Corn spurry (up to 2 whorl stage, <10 cm in height)
- Cow cockle
- Dandelion (spring or fall rosettes up to 20 cm in height)
- Flixweed
- Hemp-nettle
- Kochia⁴
- Lamb's-quarters
- Narrow-leaved hawk's-beard (up to 20 cm)
- Night flowering catchfly
- Perennial sow thistle
- Redroot pigweed
- Round-leaved mallow (1-6 leaf, <10 cm in size)
- Russian thistle
- Scentless chamomile
- Shepherd's purse (up to 30 cm)
- Stinkweed
- Stork's bill
- Volunteer canola² (1-6 leaf)
- Volunteer flax⁴
- White cockle
- Wild buckwheat
- Wild mustard

¹Suppression

²Including Group 2 herbicide tolerant canola varieties when tank-mixed with MCPA Ester.

³When tank-mixed with MCPA Ester

⁴When tank-mixed with Perimeter[®] II herbicide



Crops:

Wheat (spring and durum)



Tank-mix Options:

Perimeter[®] II: For added broadleaf control including volunteer flax and kochia

2,4-D Ester: For additional kochia control, including Group 2 biotypes

MCPA Ester: For control of volunteer Group 2 herbicide tolerant canola and annual sow thistle



Timing:

Spring and durum wheat (3-leaf to before the flag leaf stage)



Crop Rotation:

10 Months: Field Corn

11 Months: Barley, canola, dry beans, flax, lentils, oats, peas, soybeans and spring wheat.

Application Information:

Rates: 41.15 g/ac

Acres per grower bag: Minimum: 50 acres; Maximum: 200 acres

Water Volume: 5-10 imp gal/ac

Rainfastness: Two (2) hours

Surfactant: Add a registered non-ionic surfactant (NIS) such as Agral[®] 90, Liberate[®] Adjuvant, Ag-Surf[®] II, HiActivate[®] Non-ionic Liquid Spreader/Activator, Super Spreader Adjuvant, Enhance[®], Pro-Surf[®] II or Sentry[®] at 2.5 L per 1000 L of spray solution.

Mixing Instructions: Ensure that PrecisionPac[®] CS-100-23235 herbicide is fully dissolved before adding tank-mix partners.

Grazing/Harvest Restrictions: Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.

Night-flowering Catchfly



Volunteer Canola Control

