

ONE PASS. ONE SOLUTION.

PrecisionPac® CS-100-2525 herbicide combines our pyroxsulam and PP-2525 herbicide for an all-in-one combination that controls broadleaf and grassy weeds in spring and durum wheat.

Dispensed in easy-to-handle, premeasured pouches for your exact weed spectrum and sprayer size, you buy only the amount you need.

That means customized, top-performing weed control, great value and no leftovers to store!

The flexibility of PrecisionPac® herbicides gives you the power to fully customize your weed control.

The end result is:

- | | | | |
|---|---|--|--|
| 
High Performing Weed Control | 
Completely Customized Weed Control | 
Easy to Use | 
Right Product, Right Rate, Right Acre |
| 
Multiple Modes of Action | 
Pick Length Of Herbicide Activity | 
Reduced Waste | 
Constantly Evolving |



    @FMCAgCanada
ag.FMC.com/ca | 1-833-362-7722

Always read and follow label directions. Member of CropLife Canada.

FMC, the FMC logo, Perimeter and PrecisionPac are trademarks of FMC Corporation or an affiliate. © 2021 FMC Corporation. All rights reserved. 79737 - 10/21
Liberate and LI 700 are trademarks of Loveland Products, Inc. Sentry is a trademark of Nufarm Agriculture, Inc. Agral is a trademark of a Syngenta Group Company. AgSurf is a trademark of Interprovincial Cooperative Limited. Enhance and Pro-Surf are trademarks of Norac Concepts Inc. HIActivate is a trademark of WinField Solutions, LLC.

FMC An Agricultural Sciences Company



PRECISIONPAC
APPLICATION INNOVATION

CS-100-2525 for tailor-made, one-pass control.

PRECISIONPAC® CS-100-2525 Herbicide

PrecisionPac CS-100-2525 herbicide delivers outstanding control of your toughest grassy and broadleaf weeds.

Grassy Weeds Controlled

- Barnyard grass (1-5 leaf)
- Downy brome¹ (2-6 leaf, 4 tiller)
- Green foxtail¹ (1-5 leaf)
- Japanese brome (1-6 leaf)
- Persian darnel¹ (1 - 4 leaf, 2 tiller)
- Wild oats (up to 4 leaf, 2 tiller)
- Yellow foxtail (1-5 leaf)

Broadleaf Weeds Controlled

- Annual smartweed (green smartweed, lady's-thumb)
- Canada thistle¹ (up to 30 cm tall, pre-bud)
- Cleavers (up to 6 whorl)
- Common chickweed (up to 10 cm)
- Corn spurry (up to 2 whorl stage, <10 cm in height)
- Cow cockle
- Dandelion¹ (seedlings and overwintered rosettes less than or equal to 20 cm)
- Flixweed
- Hemp-nettle
- Kochia³
- Lamb's-quarters
- Narrow-leaved hawk's-beard
- Night flowering catchfly
- Perennial sow thistle
- Redroot pigweed
- Round-leaved mallow (up to 6 leaf stage, <10 cm in size)
- Russian thistle
- Shepherd's purse (up to 30 cm tall)
- Stinkweed (up to 30 cm tall)
- Stork's bill³ (1-6 leaf)
- Volunteer flax³ (up to 12 cm in height)
- Volunteer tolerant canola² (1-6 leaf)
- Wild buckwheat
- Wild mustard

¹Suppression

²Tank-mix with MCPA Ester to control Group 2 tolerant canola.

³For control tank-mix with Perimeter® II herbicide



Crops:

Wheat (spring and durum)



Tank-mix Options:

Perimeter® II: For added broadleaf control including kochia (including Group 2 resistant biotypes; seedling to 8-leaf stage), volunteer flax (up to 12 cm in height) and stork's bill (1-6 leaf)

MCPA Ester: For control of volunteer Group 2 herbicide tolerant canola, dandelion, scentless chamomile and white cockle

2,4-D Ester: For added control of kochia (including Group 2 resistant biotypes)



Timing:

Spring and durum wheat: 3 leaf to before the flag leaf stage



Crop Rotation:

10 months: Potato, sunflower

11 months: Barley, brown mustard, canola, dry bean (of the species *Phaseolus vulgaris*), flax, canola quality Brassica juncea, lentils, oats, field peas, chickpea, spring wheat, soybean and yellow mustard or fields can be summerfallowed

Excellent Wild Buckwheat Control



Source: Carman, MB, 2016. 29 DAA

Outstanding Grassy and Broadleaf Weed Control in Seeding Miss



PrecisionPac CS-100-2525 herbicide. 13 DAA

Source: Carman, MB, 2016



Untreated Check

Application Information:

Rates: 40.5 g/ac

Acres per grower bag: Minimum: 20 acres;
Maximum: 200 acres

Water Volume: 5-10 imp. gal/ac

Rainfastness: Two (2) hours

Surfactant: Add Agral® 90, Liberate® Adjuvant, Ag-Surf® II, HiActivate® Non-ionic Liquid Spreader/Activator, Super Spreader Adjuvant, Enhance®, Pro-Surf® II or Sentry® at 2.5 L per 1000 L of spray solution

Mixing Instructions: Ensure that PrecisionPac CS-100-2525 herbicide is fully dissolved before adding tank-mix partners. PrecisionPac CS-100-2525 herbicide is also registered for aerial application.

Grazing/Harvest Restrictions: Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.