

ONE PASS. ONE SOLUTION.

PrecisionPac® CS-75-12 herbicide is a broadleaf weed specialist. This powerful blend delivers cost-effective control of key weeds including chickweed, lamb's-quarters and wild buckwheat. If wild oat pressure is low, PrecisionPac® CS-75-12 herbicide is an effective, economical choice to protect your spring, durum and winter wheat.

Why PrecisionPac® herbicides? Dispensed in easy-to-handle, premeasured pouches for your exact weed spectrum and sprayer size, you buy only the amount you need. That means customized, top-performing weed control, great value and no leftovers to store!

The flexibility of PrecisionPac® herbicides gives you the power to fully customize your weed control.

The end result is:



    @FMCAgCanada
ag.FMC.com/ca | 1-833-362-7722

Always read and follow label directions. Member of CropLife Canada.

FMC, the FMC logo, Perimeter and PrecisionPac are trademarks of FMC Corporation or an affiliate.

© 2021 FMC Corporation. All rights reserved. 79737 - 10/21

Liberate and LI 700 are trademarks of Loveland Products, Inc. Sentry is a trademark of Nufarm Agriculture, Inc. Agral is a trademark of a Syngenta Group Company. AgSurf is a trademark of Interprovincial Cooperative Limited. Enhance and Pro-Surf is a trademark of Norac Concepts Inc. HiActivate is a trademark of WinField Solutions, LLC.

FMC An Agricultural Sciences Company



PRECISIONPAC
APPLICATION INNOVATION

CS-75-12
Cost effective
broadleaf and grass
control in your spring
and durum wheat.

PRECISIONPAC® CS-75-12 Herbicide

When wild oat pressure is low,
this PrecisionPac blend is an ideal and economical
choice for in-crop broadleaf weed control.

Grassy weeds controlled

- Wild oats (up to 4 leaf, 2 tiller) for control of wild populations <75 plants/m²
- Japanese brome (1 leaf to 4 leaf, 2 tiller; under good growing conditions)

Broadleaf weeds controlled

- Common chickweed (1-6 leaf)
- Corn spurry
- Cow cockle
- Green smartweed
- Hemp-nettle
- Lamb's-quarters
- Narrow-leaved hawk's beard (seedlings and overwintered rosettes prior to bolting)¹
- Redroot pigweed
- Russian thistle
- Smartweed (Lady's thumb)
- Stinkweed
- Wild mustard
- Wild buckwheat (1-3 leaf)



Crops:

Wheat (spring, durum and winter)



Timing:

Spring and durum wheat: 3 leaf to before flag leaf stage

Winter wheat: Apply in the spring when winter wheat is in the 2-7 leaf, 4 tiller stage



Crop rotation:

10 months: Field corn, sunflower, potatoes

11 months: Barley, brown mustard, canola, dry bean (of the species *Phaseolus vulgaris*), flax, canola quality Brassica juncea, lentils, oats, field peas, chickpea, spring wheat, soybean and yellow mustard or fields can be summerfallowed.



Application information:

Rates: 33 g/ac

Acres per grower bag:

Minimum: 20 acres; Maximum: 220 acres

Surfactant: Add a registered non-ionic surfactant (NIS) such as: Agral® 90, Liberate® Adjuvant, Ag-Surf® II, HiActivate® Non-ionic Liquid Spreader/Activator, Super Spreader Adjuvant, Enhance®, Pro-Surf® II or Sentry® at 2.5 L per 1000 L of spray solution.

Rainfastness: Two (2) hours

Water volume: 5-10 imp gal/ac

Mixing Instructions: Ensure that PrecisionPac® CS-75-12 herbicide is fully dissolved before adding tank-mix partners. CS-75-12 is also registered for aerial application.

Grazing/Harvest Restrictions: Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.



Wild oats



Wild buckwheat

¹ Suppression.