

## GREAT WEED CONTROL. EXCEPTIONAL VALUE.

PrecisionPac® CS-75-2525 herbicide offers a cost-effective solution for controlling low populations of wild oats. It combines a reduced rate of our pyroxulam and PP-2525 herbicide for an all-in-one combination that controls low populations of wild oats and exceptional broadleaf weed control in spring and durum wheat. Great weed control for a great price.

Dispensed in easy-to-handle, premeasured pouches for your exact weed spectrum and sprayer size, you buy only the amount you need. That means customized, top-performing weed control, great value and no leftovers to store!

The flexibility of PrecisionPac® herbicides gives you the power to fully customize your weed control.

The end result is:



    @FMCAgCanada  
ag.FMC.com/ca | 1-833-362-7722

Always read and follow label directions. Member of CropLife Canada.

FMC, the FMC logo, Perimeter and PrecisionPac are trademarks of FMC Corporation or an affiliate.

© 2021 FMC Corporation. All rights reserved. 79737 - 10/21

Liberate and LI 700 are trademarks of Loveland Products, Inc. Sentry is a trademark of Nufarm Agriculture, Inc. Agral is a trademark of a Syngenta Group Company. AgSurf is a trademark of Interprovincial Cooperative Limited. Enhance and Pro-Surf is a trademark of Norac Concepts Inc. HiActivate is a trademark of WinField Solutions, LLC.

**FMC** An Agricultural  
Sciences Company



**PRECISIONPAC**  
APPLICATION INNOVATION

**CS-75-2525**  
provides  
tailor-made,  
one-pass control.

# PRECISIONPAC<sup>®</sup> CS-75-2525 Herbicide

Offers a solution for controlling low populations of wild oats with the same powerful broadleaf performance of PrecisionPac<sup>®</sup> PP-2525 herbicide.

## Grassy Weeds Controlled

- Wild oats\* (up to 4 leaf, 2 tiller)
- Japanese brome (1 leaf to 4 leaf, 2 tiller stage; under good growing conditions)

## Broadleaf Weeds Controlled

- Annual smartweed (green smartweed, lady's thumb)
- Canada thistle<sup>1</sup>
- Cleavers<sup>4</sup> (1-6 whorl)
- Common chickweed (1-6 whorl)
- Cow cockle
- Dandelion<sup>3</sup> (spring and fall rosettes, up to 15 cm in diameter)
- Flixweed
- Hemp-nettle
- Kochia<sup>4</sup>
- Lamb's-quarters
- Narrow-leaved hawk's-beard
- Night flowering catchfly
- Perennial sow thistle
- Redroot pigweed
- Russian thistle
- Scentless chamomile<sup>3</sup>
- Stinkweed
- Stork's bill<sup>4</sup> (1-6 leaf)
- Volunteer canola<sup>2</sup> (1-6 leaf)
- Volunteer flax<sup>4</sup> (up to 12 cm in height)
- White cockle<sup>3</sup>
- Wild buckwheat
- Wild mustard

\* For control of low wild oat populations (<75 plants/m<sup>2</sup>) only

<sup>1</sup> Suppression

<sup>2</sup> Tank-mix with MCPA Ester to control Group 2 herbicide tolerant canola

<sup>3</sup> When tank-mixed with MCPA

<sup>4</sup> For control tank-mix with Perimeter<sup>®</sup> II herbicide



## Crops:

Wheat (spring and durum)



## Tank-mix Options:

**Perimeter<sup>®</sup> II:** For added broadleaf control including cleavers, kochia, stork's-bill and volunteer flax

**2,4-D Ester:** For added control of kochia (including Group 2 resistant biotypes)

**MCPA:** For control of volunteer Group 2 herbicide tolerant canola, dandelion, scentless chamomile and white cockle



## Timing:

Spring and durum wheat: 3 leaf to before the flag leaf stage



## Crop Rotation:

**10 months:** Field corn, potato, sunflower

**11 months:** Barley, brown mustard, canola, dry bean (of the species *Phaseolus vulgaris*), flax, canola quality Brassica juncea, lentils, oats, field peas, chickpea, spring wheat, soybean and yellow mustard or fields can be summerfallowed

## Excellent Wild Buckwheat Control



## Application Information:

**Rates:** 33.14 g/ac

**Acres per grower bag:** Minimum: 20 acres;

Maximum: 220 acres

**Water Volume:** 5-10 gallons/acre

**Rainfastness:** Two (2) hours

**Surfactant:** Add a registered non-ionic surfactant (NIS)

such as: Agral<sup>®</sup> 90, Liberate<sup>®</sup> Adjuvant, Ag-Surf<sup>®</sup> II, HiActivate<sup>®</sup> Non-ionic Liquid Spreader/Activator, Super Spreader Adjuvant, Enhance<sup>®</sup>, Pro-Surf<sup>®</sup> II or Sentry<sup>®</sup> at 2.5 L per 1000 L of spray solution

**Mixing Instructions:** Ensure that PrecisionPac<sup>®</sup>

CS-75-2525 herbicide is fully dissolved before adding tank-mix partners

PrecisionPac CS-75-2525 herbicide is also registered for aerial application.

**Grazing/Harvest Restrictions:** Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.