



Aim® EC herbicide can turn harvest conditions in your favour

Add Aim® EC herbicide to your pre-harvest glyphosate application to control weeds and dry-down plant material to make harvest faster and easier.

How can Aim® EC herbicide improve my harvest?

Some years, growing conditions give weeds the upper hand, making harvest tougher and more time-consuming than it needs to be. An application of Aim® EC herbicide can turn harvest conditions in your favour.

Fast-acting. Adding Aim® EC herbicide to your pre-harvest application helps dry-down plant material and provides better, faster control of a wide range of troublesome broadleaf weeds for an easier harvest.

Powerful Group 14 broadleaf activity. Aim® EC herbicide is effective on a range of difficult broadleaf weeds that can hinder harvest, including many that are resistant to other chemistries.

Convenient. Aim® EC herbicide features an easy-to-handle formulation and low use rates, to get you out there quicker and done sooner.

Use with or without Glyphosate. Harvest aids, like glyphosate, are under increased scrutiny from buyers, limiting growers' choices to help with crop and weed dry-down. When applied without glyphosate, ensure that a registered surfactant is used.

Now available in 1.2 and 4.8 L jug sizes for easier use.

HARVEST AID GROUP 14

Quick facts:

Formulation:

EC – emulsifiable concentrate

Pre-harvest Rates:

Apply at 47 mL per acre

1.2 L jug = 25 ac/jug

4.8 L jug = 102 ac/jug

Post-harvest Rates:

1.2 L jug = 40 - 80 ac/jug

4.8 L jug = 160 - 320 ac/jug

Water volume:

Minimum of 10 gal/ac (100 L/ha)

Rainfastness:

1 hour

Timing:

Pre-seed burnoff

Chemfallow

Harvest aid

Post-harvest

Key Crops:

Barley

Dry beans

Oat

Peas

Potatoes*

Soybeans

Wheat

* See label for potato rate and application information.

Registered Surfactants:

Non-ionic surfactant at 0.25% v/v (0.25 litres per 100 litres of spray solution)

Destination™ MSO at 1% v/v (1 litre per 100 litres of spray solution)

Journey™ HSOC at 0.5% v/v (0.5 litre per 100 litres of spray solution) Merge® at 1% v/v (1 litre per 100 litres of spray solution)

When applied without glyphosate, ensure that a registered surfactant is used.

For harvest aid use, a surfactant is recommended with or without glyphosate.

Aim® EC herbicide vs Heat® LQ

Application: September 5. Photo taken: 8 days after application.



No harvest aid

Aim® EC herbicide

Heat® LQ

Source: Josephburg, Alberta, 2017

Aim® EC herbicide 47mL/ac +
1 REL WeatherMAX + 0.25% v/v Ag-Surf™

Heat® LQ herbicide 30 ac/jug +
1 REL WeatherMAX + 1% Merge®

When do I apply Aim® EC herbicide in a pre-harvest application with glyphosate?

Wheat, barley and oats: Apply when grain moisture is less than 30% (hard dough stage; thumbnail impression remains on seed).

Field peas: Apply when grain moisture is less than 30%. Apply to dry field pea when majority (75% to 80%) of the pods are brown, the bottom pods are ripe and the pea seeds have detached from the pods.

Chickpeas: Apply when grain moisture is less than 30%. Apply to chickpeas when 80% to 90% of bean leaves have fallen and pods are mature (yellow to brown in color) and 80% to 90% leaf drop (original leaves) has occurred.

Faba beans: Apply when grain moisture is less than 30%. At this stage, the majority of the faba bean plants will be ripe and dry. Pods will be fully filled and the bottom pods will be tan or black in color.

Dry beans: Apply when grain moisture is less than 30%. Apply to dry beans when 80% to 90% of bean leaves have fallen and pods are mature (yellow to brown in color).

Soybeans: Apply when grain moisture is less than 30%. Apply to soybeans when crop has lost 80% to 90% of leaves and 80% of pods are brown.

Pre-Harvest Interval

When using Aim® EC herbicide alone, without glyphosate, the pre-harvest intervals below apply.

Crop/Crop Group/Crop Subgroup	PHI (Days Before Harvest)
Barley (harvest aid)	3
Oats (harvest aid)	3
Wheat (harvest aid)	3
Soybean (harvest aid)	3
Potato (harvest aid)	7
Dried shelled peas and beans (harvest aid)	3

When using Aim® EC herbicide with glyphosate, refer to the glyphosate label for specific pre-harvest interval recommendations. The use of Aim® EC herbicide will not speed up the maturity of green crops.

Always check with your grain buyer and/or grain contracts prior to using pre-harvest herbicides.

For more information see: keepitclean.ca

Aim® EC herbicide Harvest Aid on Dry Beans



Untreated Dry Beans



Aim® EC herbicide 47mL/ac 6 DAA

Spray Date: August 29, 2020. Photos taken: September 4, 2020.

Questions? Ask your retailer about Aim® EC herbicide today.

* Terms and Conditions Apply. Contact your local retailer for details.
Always read and follow label instructions. Member of CropLife Canada.
FMC, the FMC logo, and Aim are trademarks of FMC Corporation or an affiliate.
Heat and Merge are trademarks of BASF, WeatherMAX is a trademark of Bayer Group, Ag-Surf is a trademark of IPSCO,
Destination MSO and Journey HSOC are trademarks of WinField United,
© 2023 FMC Corporation. All rights reserved. 5719 - 7/23





 @FMCAGCanada
ag.FMC.com/ca | 1-833-362-7722