



Intruvix® II herbicide consistently provides quick and lasting burnoff control of tough weeds like kochia, narrow-leaved hawk's-beard and volunteer canola in front of cereals

What is Intruvix® II herbicide?

Adding Intruvix® II herbicide to your glyphosate provides an exceptional burnoff before planting cereals. It delivers fast-acting, systemic action on many broadleaf weeds and uses 4 modes of action when tank-mixed with glyphosate, to prevent resistant weed escapes.

How does Intruvix® II herbicide help my farm business?

Top Quality Performance

- Intruvix® II herbicide is fast acting with the power of systemic action for unparalleled weed control.
- Over 50 labelled weeds including kochia, cleavers, volunteer canola, wild buckwheat and narrow-leaved hawk's-beard.

Manage Weed Resistance

- Resistant weeds are a growing problem for Canadian farmers.
- Intruvix® II herbicide, plus glyphosate, uses 4 modes of action to tackle resistant weeds.

Flexibility

- Suitable for all soil types from brown to black to grey.
- For spring burnoff and chemfallow applications, Intruvix® II herbicide offers wide-open recrop flexibility next spring.

GROUPS
2, 4, 14 & 9*

Quick facts:

Application timing:

- Pre-seed
- Chemfallow


Water volume:

10 US gal/ac (100 L/ha)

Packaging:

One case treats 80 acres
40 acres per jug.
2 x 2.58 kg jugs per case.

*When tank-mixed with glyphosate



When and how is Intruvix® II herbicide applied?

Intruvix® II herbicide can be applied pre-seed, and as a chemfallow application. Rainfall soon after application could reduce efficacy.

What's in the box?

Intruvix® II herbicide is an all-in-one formulation. One (1) case treats 80 acres.


Which herbicides can be tank-mixed with Intruvix® II herbicide?

Must be tank-mixed with glyphosate.


No additional surfactant is required when tank-mixed with 0.5 REL glyphosate per acre.

The following mixing order is recommended:

When tank mixing, add Intruvix® II herbicide first and then add glyphosate.



Source: 2023 FMC Internal Trials. Lethbridge, AB *Rate 0.5 REL



Source: 2023 FMC Internal Trials. Lethbridge, AB *Rate 0.5 REL/ac

Crops

- Wheat (spring, durum, winter)
- Barley
- Oats

Weeds

(Intruvix® II herbicide plus 0.5 L/ac glyphosate equivalent)

- Canada fleabane (up to 8 cm)
- Canada thistle (rosette)¹
- Carpetweed
- Cleavers (up to 15 cm)
- Cocklebur
- Common ragweed (up to 8 cm)
- Cow cockle (up to 3 leaf)
- Dandelion (up to 15 cm)
- Downy brome (up to 15 cm)
- Flixweed (up to 15 cm)
- Foxtail (giant, green up to 15 cm)
- Hemp-nettle (up to 15 cm)
- Jimsonweed
- Kochia (up to 10 cm)
(Incl. Group 2 & 9-resistant biotypes)
- Lady's thumb (up to 15 cm)
- Lamb's-quarters (up to 15 cm)
- Morning glory (up to 10 cm)
- Narrow-leaved hawk's-beard (up to 8 cm)
- Nightshade (Eastern black, black, hairy) (up to 10 cm)
- Pennsylvania smartweed (seedling)
- Persian darnel (up to 15 cm)
- Pigweed (prostrate, smooth, tumble)
- Purslane
- Round-leaved mallow
- Redroot pigweed (up to 15 cm)
- Russian thistle (up to 15 cm)
- Scentless chamomile (up to 8 cm)¹
- Shepherd's purse
- Stinkweed (up to 15 cm)
- Tansy mustard
- Velvetleaf (up to 10 cm)
- Volunteer flax (up to 15 cm)
- Volunteer barley (up to 15 cm)
- Volunteer canola (up to 15 cm)
- Volunteer wheat (up to 15 cm)
- Waterhemp (tall, common up to 10 cm)
- Wild oats (up to 15 cm)
- Wild buckwheat (up to 8 leaf)
- Wild mustard (up to 15 cm)
- White cockle (rosette)¹

¹Suppression

Questions? Ask your retailer about Intruvix® II herbicide today.

Always read and follow label instructions. Member of CropLife Canada.

FMC, the FMC logo and Intruvix are trademarks of FMC Corporation or an affiliate.
© 2025 FMC Corporation. All rights reserved. 23364 - 10/25