

GROUP	2	HERBICIDE
-------	----------	-----------

SPITFIRE® 50 SC Herbicide

Suspension

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY

SPITFIRE® 50 SC Herbicide is a selective herbicide for postemergent control or suppression of annual broadleaf weeds including cleavers, wild buckwheat and chickweed in spring wheat (including durum), spring barley, and oats (tank mixes only).

ACTIVE INGREDIENT: Florasulam 50 g/L
Contains 1,2-benzisothiazoline-3-one at 0.02% as a preservative

REGISTRATION NO. **31252**
PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

MAY CAUSE SLIGHT EYE IRRITATION



IN CASE OF EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

NET CONTENTS: 0.8 L – Bulk

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

PRECAUTIONS

- **MAY CAUSE SLIGHT EYE IRRITATION**
- **KEEP OUT OF REACH OF CHILDREN**
- **DO NOT APPLY BY AIR**
- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application with a closed cab.
- **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
- **At completion of spraying or end of the day:** Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.
- Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient. Treat symptomatically.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

- **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.

- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store in original containers in a secure, dry heated storage. SPITFIRE® 50 SC Herbicide will freeze at -10°C. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Store this product away from food and feed. Do not contaminate food, feedstuffs or domestic water supplies. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

DO NOT reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Returnable Containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label.

SPITFIRE® is a trademark of FMC Corporation or an affiliate.

PMRA Approved Label – M.A.
2023-2424 2024-10-29

GROUP	2	HERBICIDE
-------	----------	-----------

SPITFIRE® 50 SC Herbicide

Suspension

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY

SPITFIRE® 50 SC Herbicide is a selective herbicide for postemergent control or suppression of annual broadleaf weeds including cleavers, wild buckwheat and chickweed in spring wheat (including durum), spring barley, and oats (tank mixes only).

ACTIVE INGREDIENT: Florasulam 50 g/L
Contains 1,2-benzisothiazoline-3-one at 0.02% as a preservative

REGISTRATION NO. **31252**
PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

MAY CAUSE SLIGHT EYE IRRITATION

CAUTION



POISON

IN CASE OF EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

PRECAUTIONS

- **MAY CAUSE SLIGHT EYE IRRITATION**
- **KEEP OUT OF REACH OF CHILDREN**
- **DO NOT APPLY BY AIR**

- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application with a closed cab.
- DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

- **At completion of spraying or end of the day:** Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.
- Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient. Treat symptomatically.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

- TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store in original containers in a secure, dry heated storage. SPITFIRE® 50 SC Herbicide will freeze at -10°C. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Store this product away from food and feed. Do not contaminate food, feedstuffs or domestic water supplies. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

DO NOT reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Returnable Containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

SPITFIRE® 50 SC Herbicide is a selective postemergence herbicide for the control or suppression of hard-to-kill annual broadleaf weeds such as chickweed, wild buckwheat and cleavers in spring wheat (including durum), spring barley, and oats (tank mixes only) not underseeded with legumes. SPITFIRE® 50 SC Herbicide is mixed with water and applied as a uniform broadcast spray. It is non-corrosive, nonflammable, and nonvolatile.

SPITFIRE® 50 SC Herbicide must be applied early postemergence, to the main flush of actively growing broadleaf weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of SPITFIRE® 50 SC Herbicide by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately

PMRA Approved Label – M.A.
2023-2424 2024-10-29

controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds. See “DIRECTIONS FOR USE” section of this label for complete use details.

SPITFIRE® 50 SC Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

MODE OF ACTION

SPITFIRE® 50 SC Herbicide inhibits the production of the ALS enzyme in plants. This enzyme is essential for the production of certain amino acids which are essential for plant growth. SPITFIRE® 50 SC Herbicide is a Group 2 mode of action herbicide.

GENERAL USE PRECAUTIONS

Sensitive Plants

Do not apply SPITFIRE® 50 SC Herbicide directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

Non-Target Sites

Do not apply where proximity of susceptible crops (e.g., canola and legumes) or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Precautions section of the label.

Crop Rotation

Fields previously treated with SPITFIRE® 50 SC Herbicide can be seeded the following year to barley, canola, peas, wheat or fields can be summerfallowed.

Do not use in successive years at the same site.

This product has potential to leach. Do not apply excessive irrigation.

Tank Mixtures:

This product may be tank mixed with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact FMC of Canada Limited at 1-833-362-7722 for information before applying any tank mix that is not specifically recommended on this label.

Spray Equipment Precaution

Do not apply through any type of irrigation system.

DO NOT APPLY BY AIR.

To Reduce Spray Drift:

1. Use nozzles delivering higher volumes and coarser droplets.
2. Use low pressures (200 to 275 kPa).
3. Use 100 L/ha of spray solution.
4. Spray when the wind velocity is 15 km/hr or less.
5. Spot treatments should only be applied with a calibrated boom to prevent over-application.

Sprayer Clean-Out Instructions

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:
 - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
 - Agitate and circulate for 15 minutes, and flush through booms and hoses.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.
3. Second rinse:
 - Fill the tank with clean water.
 - Add All Clear Spray Tank Decontaminator, or Clean-Out Spray Tank Cleaner, or 1 L of household ammonia (containing a minimum of 3 % ammonia) per 100 L of water, or similar tank cleaning agent as per manufacturer's recommendations while filling the tank with clean water.
 - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
 - After flushing the boom and hoses, drain tank completely.
 - Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
4. Third rinse:
 - Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.

- Drain tank completely.

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty odour which may cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE.

DO NOT APPLY TO CROPS UNDERSEEDDED WITH LEGUMES.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

SPRAY BUFFER ZONES

A spray buffer zone is **NOT** required for:

- uses with hand-held application equipment permitted on this label

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Crop	Method of application	Spray Buffer Zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat:
		Less than 1 m	Greater than 1 m	
All labelled crops	Field sprayer (medium)	1	0	3

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product that are larger than 1m can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticide portion of the Canada.ca web site.

SPITFIRE® 50 SC HERBICIDE ALONE

Crops Registered

Spring wheat (including durum) and spring barley

Application Directions

Apply 100 mL of SPITFIRE® 50 SC Herbicide per hectare in a minimum of 100 L per hectare of water. Add Agral® 90 at 0.2% v/v. See weeds species controlled under “Weeds Controlled or Suppressed by SPITFIRE® 50 SC Herbicide Alone at 100 mL/ha.” Apply to actively growing wheat and barley from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Application Timing

Apply to actively growing weeds in the 2-4 leaf stage. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Only weeds which are emerged at the time of application will be affected. If foliage is wet at the time of application control may be decreased. Under conditions of low crop and high weed density, control may be reduced.

Weeds Controlled or Suppressed by SPITFIRE® 50 SC Herbicide Alone at 100mL/ha

Weeds Controlled:	Weeds Suppressed:
Buckwheat, wild Canola, volunteer ¹ Chickweed, common Cleavers Mustard, wild Shepherd's purse Smartweed Stinkweed,	Hempnettle, Pigweed, redroot Sowthistle, annual Sowthistle, perennial ²

¹Including herbicide-tolerant canola varieties except Clearfield Canola.

²Applications made at advanced leaf stages will reduce product effectiveness.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of SPITFIRE® 50 SC Herbicide
4. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
5. Add Agral® 90 as the last ingredient at the rate of 0.2% v/v (2 L per 1000 L of spray volume).

PMRA Approved Label – M.A.
2023-2424 2024-10-29

6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g., canola and legumes).
7. Follow sprayer clean-up directions.

Preharvest/Grazing Intervals

Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.

TANK MIX COMBINATIONS:

SPITFIRE® 50 SC HERBICIDE + MCPA LV 500 HERBICIDE

Crops Registered

Spring wheat (including durum), spring barley, and oats

Application Directions

For control of a wide spectrum of broadleaf weeds apply SPITFIRE® 50 SC Herbicide at 100 mL/ha tank mixed with MCPA LV 500 Herbicide at 0.70 L/ha (350 g ai/ha) in 100 L per hectare of water. Apply to actively growing wheat, barley, and oats from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Weeds Controlled or Suppressed by the Tank Mixture of SPITFIRE® 50 SC Herbicide + MCPA LV 500 Herbicide

Weeds Controlled	Weeds Suppressed
Buckwheat, wild	Dandelions (seedlings and overwintering rosettes less than 15 cm) ³
Burdock ¹	Plantain (top growth)
Canola, volunteer ²	Sowthistle, annual
Chickweed, common	Sowthistle, perennial
Cleavers	Stork's bill ³
Flixweed (spring rosettes only)	Thistle, Canada ³
Hempnettle ³	
Kochia ¹	
Lamb's-quarters	
Mustard, ball	
Mustard, wild	
Pigweed, redroot ³	
Pigweed, Russian ¹	
Prickly lettuce ¹	
Ragweed, common	
Shepherd's purse	
Smartweed	
Stinkweed	
Sunflower, annual ¹	

¹Up to the 4-leaf stage

²Including herbicide-tolerant canola varieties

³Optimum control can be achieved with the addition of 140 mL/ha (70 g ai/ha) of MCPA LV 500 Herbicide

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of SPITFIRE® 50 SC Herbicide
4. Add the correct amount of MCPA LV 500 Herbicide and continue to agitate.
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g., canola and legumes).
7. Follow sprayer clean-up directions.

Note: Do not add a surfactant to this tank mixture.

Pre-Harvest/Grazing Intervals (SPITFIRE® 50 SC Herbicide + MCPA LV 500 Herbicide)

1. Do not cut the treated crop for hay or graze treated crop within 7 days after application.
2. Do not harvest the treated crop within 60 days after application.

TANK MIX COMBINATIONS:

SPITFIRE® 50 SC HERBICIDE+ CURTAIL™ M HERBICIDE

Crops Registered

Spring wheat (including durum), spring barley, and oats

Application Directions

For control of a wide spectrum of annual broadleaf weeds plus Canada thistle and perennial sowthistle, apply SPITFIRE® 50 SC Herbicide at 100 mL/ha tank mixed with 1.5 L/ha of Curtail M Herbicide in 100 L per hectare of water. Apply to actively growing wheat, barley, and oats from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Weeds Controlled or Suppressed by the Tank Mixture of SPITFIRE® 50 SC Herbicide + Curtail™ M Herbicide

Weeds Controlled	Weeds Suppressed
Buckwheat, wild Canola, volunteer ¹ Chickweed, common Cleavers Flixweed (spring rosettes only) Hempnettle Lamb's-quarters Mustard, wild Pigweed, redroot Shepherd's-purse Sowthistle, perennial (top growth only) ² Sowthistle, annual Stinkweed Stork's-bill Thistle, Canada	Dandelion (overwintered rosettes) Plantain (top growth)

¹Including herbicide-tolerant canola varieties

²Control not observed until a minimum of 40 days after treatment

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of SPITFIRE® 50 SC Herbicide
4. Add the correct amount of Curtail M Herbicide and continue to agitate.
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g., canola and legumes).
7. Follow sprayer clean-up directions.

Note: Do not add a surfactant to this tank mixture.

Pre-Harvest/Grazing Intervals (SPITFIRE® 50 SC Herbicide + Curtail M Herbicide)

1. Do not cut the treated crop for hay or graze treated crop within 7 days after application.
2. Do not harvest the treated crop within 60 days after application.

TANK-MIX COMBINATIONS:

SPITFIRE® 50 SC HERBICIDE + 2,4-D LV 600 HERBICIDE

Crops Registered

Spring wheat (including durum)

Application Directions

For control of a wide spectrum of broadleaf weeds apply SPITFIRE® 50 SC Herbicide at 100 mL/ha tank-mixed with 2,4-D LV 600 Herbicide at 1.0 L/ha. Apply when wheat is in the 3 leaf expanded stage to the 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be affected. Best results are obtained from application made to seedling weeds.

Weeds Controlled or Suppressed by the Tank Mixture of SPITFIRE® 50 SC Herbicide + 2,4-D LV 600 Herbicide

Weeds Controlled	Weeds Suppressed
Bluebur	Hempnettle
Burdock	Sowthistle, perennial (top growth control)
Buckwheat, tartary	Thistle, Canada (top growth control)
Buckwheat, wild	
Canola, volunteer ¹	
Chickweed, common	
Cleavers	
Cocklebur	
Dandelions (seedlings and overwintered rosettes)	
Flixweed	
Kochia ²	
Lady's-thumb	
Lamb's-quarters	
Mustard, ball	
Mustard, wild	

Weeds Controlled	Weeds Suppressed
Pigweed, redroot Plantain Prickly lettuce Radish, wild Ragweed, common Shepherd's-purse Smartweed Sowthistle, annual Stinkweed Sunflower, annual Thistle, Russian Vetch	

¹Including herbicide-tolerant canola varieties

²Up to 5 cm in height.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add SPITFIRE® 50 SC Herbicide
4. Add 2,4-D LV 600 Herbicide and continue to agitate.
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g., canola and legumes).
7. Follow sprayer clean-up directions.

Pre-Harvest/Grazing Intervals (SPITFIRE® 50 SC Herbicide + 2,4-D LV 600 Herbicide)

1. Do not permit lactating dairy animals to graze fields within 7 days after application.
2. Do not harvest forage or cut hay within 30 days after application.
3. Withdraw meat animals from treated fields at least 3 days before slaughter.
4. Do not harvest the treated mature crop within 60 days after application.

TANK MIX COMBINATIONS:

SPITFIRE® 50 SC HERBICIDE + GLYPHOSATE PRESENT AS ISOPROPYLAMINE SALT OR DIMETHYLAMINE SALT

Crops Registered

Spring barley, Oats, Spring wheat (including durum)

Application Directions

SPITFIRE® 50 SC Herbicide may be tank mixed with glyphosate present as isopropylamine salt or dimethylamine salt to control volunteer glyphosate-tolerant canola, wild buckwheat (up to 5 leaf) and top growth control of dandelion (up to 6 leaf stage), in addition to the labelled weeds controlled by SPITFIRE® 50 SC Herbicide or glyphosate.

Apply SPITFIRE® 50 SC Herbicide at 100 mL/ha with glyphosate herbicide (present as isopropylamine salt or dimethylamine salt) at 450 g a.e./ha in 100 L water per hectare.

Application Timing

Apply in the spring prior to planting barley, oats and wheat, or as an initial summerfallow treatment, or in the fall prior to planting barley, oats and wheat the following spring. For annual weed control in the fall, only the weeds at labelled growth stage will be controlled. Heavy frost prior to application in the fall may decrease control.

Do not use in successive years at the same site.

Apply only one of the following treatments to the same site in the same year: a tank-mix of SPITFIRE® 50 SC Herbicide and glyphosate (present as isopropylamine salt or dimethylamine salt, prior to planting or as the initial treatment in summerfallow), or any post emergent herbicide containing florsulam (Spectrum™ Herbicide Tank-Mix).

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, SPITFIRE® 50 SC Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to SPITFIRE® 50 SC Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of SPITFIRE® 50 SC Herbicide or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information, related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates, precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area, if possible, by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact FMC of Canada Ltd. at 1-833-362-7722.

PMRA Approved Label – M.A.
2023-2424 2024-10-29

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label.

SPITFIRE® is a trademark of FMC Corporation or an affiliate.

Agral® is a trademark of a Syngenta Group Company.

Ag-Surf® is a trademark of Interprovincial Cooperative Ltd.

Curtail™, and Spectrum™ are trademarks of Corteva Agriscience Canada Company

Assert® is a registered trademark of Nufarm Agriculture Inc.

Everest® is a registered trademark of Bayer Corp. used under licence by Bayer Inc.

Solupak® is a registered trademark of Bayer Inc.