

Wirkungsspektrum der Getreideherbizide.

	Omnera® LQM®	Pointer® Plus	Pointer® SX®	Artus®	Aurora®
Wintergetreide	WG, WR, T, WW	WG, WR, T, WW	WG, WR, T, WW	WG, WR, T, WW	WG, WR, T, WW
Aufwandmenge	1,0 l/ha	50 g/ha	45 g/ha	50 g/ha	50 g/ha
Sommergetreide	SG, SW	SG, SW, Hafer	SG, SW, Hafer	SG, SW, Hafer	SG, SW, Hafer
Aufwandmenge	1,0 l/ha	50 g/ha	45 g/ha	50 g/ha	50 g/ha
Ackerfrauenmantel, Gemeiner	●	●	●	●	–
Acker-Gänsedistel	●	●	●	●	–
Acker-Gauchheil	●	●	●	●	–
Acker-Hellerkraut	●	●	●	●	●
Acker-Kratzdistel	●	●	●	●	●
Acker-Krummhals	●	●	●	●	–
Acker-Lichtnelke	●	●	●	●	–
Acker-Minze	●	●	●	●	–
Acker-Senf	●	●	●	●	●
Acker-Spörgel	●	●	●	●	–
Acker-Steinsame*	●	●	●	●	–
Acker-Winde*	●	●	●	●	–
Amarant-Arten	●	●	●	●	●
Ampfer-Arten	●	●	●	●	–
Besenrauke, Gemeine	●	●	●	●	–
Bingelkraut, Einjähriges	●	●	●	●	●
Ehrenpreis*, Acker- und Feld-	●	●	●	●	●
Ehrenpreis*, Persischer	●	●	●	●	●
Erdrauch*, Gemeiner	●	●	●	●	●
Franzosenkraut-Arten	●	●	●	●	–
Gänsefuß-Arten*	●	●	●	●	●
Hahnenfuß-Arten	●	●	●	●	●
Hederich, Gemeiner	●	●	●	●	●
Hirtentäschel, Gemeines	●	●	●	●	●
Hohlzahn, Gemeiner	●	●	●	●	●
Hundskerbel	●	●	●	●	–
Hundspetersilie	●	●	●	●	–
Kamille-Arten	●	●	●	●	●
Klatsch-Mohn	●	●	●	●	●
Kletten-Labkraut	●	●	●	●	●*
Knöterich*, Ampfer-	●	●	●	●	●

	Omnera® LQM®	Pointer® Plus	Pointer® SX®	Artus®	Aurora®
Wintergetreide	WG, WR, T, WW	WG, WR, T, WW	WG, WR, T, WW	WG, WR, T, WW	WG, WR, T, WW
Aufwandmenge	1,0 l/ha	50 g/ha	45 g/ha	50 g/ha	50 g/ha
Sommergetreide	SG, SW	SG, SW, Hafer	SG, SW, Hafer	SG, SW, Hafer	SG, SW, Hafer
Aufwandmenge	1,0 l/ha	50 g/ha	45 g/ha	50 g/ha	50 g/ha
Knöterich*, Floh-	●	●	●	●	●
Knöterich*, Vogel-	●	●	●	●	●
Knöterich*, Winden-	●	●	●	●	●
Kohl-Gänsedistel	●	●	●	●	–
Kornblume*	●	●	●	●	●
Kreuzkraut, Gemeines	●	●	●	●	–
Löwenzahn	●	●	●	●	–
Melde*, Gemeine	●	●	●	●	–
Möhre, Wilde	●	●	●	●	–
Nachtschatten, Schwarzer	●	●	●	●	●
Ölrettich, Ausfall-	●	●	●	●	–
Phacelia, Ausfall-	●	●	●	●	–
Rainkohl, Gemeiner	●	●	●	●	–
Raps**, Ausfall-	●	●	●	●	●
Rauke, Weg-	●	●	●	●	●
Reiherschnabel	●	●	●	●	–
Saatwucherblume*	●	●	●	●	–
Schierling, Gefleckter	●	●	●	●	–
Sonnenblumen***, Ausfall-	●	●	●	●	–
Stiefmütterchen-Arten*	●	●	●	●	●
Storchschnabel-Arten	●	●	●	●	–
Taubnessel-Arten	●	●	●	●	●
Vergissmeinnicht, Acker-	●	●	●	●	●
Vogelmiere	●	●	●	●	●
Wicken-Arten	●	●	●	●	–
Zweizahn, Dreigeteilter	●	●	●	●	–

● sehr gute Wirkung ● gute Wirkung ● Teilwirkung ● nicht ausreichende Wirkung – keine Information

* In frühen Entwicklungsstadien bis max. BBCH 14–16 bzw. 2.–3. Quirl
** Keine ausreichende Wirkung auf Clearfield®-Ausfallraps
*** Keine ausreichende Wirkung auf Tribenuron-methyl -resistente Ausfallsonnenblumen

Quelle: FMC Ergebnisse