

Steward® EC Insecticide



ALFALFA

Effectively combat alfalfa weevil larvae and other pests.

Alfalfa growers count on Steward® EC insecticide to help keep key insects from taking a bite out of profits by delivering fast, broad-spectrum protection from alfalfa weevil larvae, beet armyworm and many Lepidopteran pests commonly resistant to other insecticides. Studies have shown that Steward EC insecticide works fast, targeting insects through ingestion. It provides rapid feeding cessation of adults and up to two weeks of residual control, as demonstrated in trial results from multiple locations.

In addition to these benefits, Steward EC insecticide has a low impact on most predatory insects and mites when used in accordance with the label. It is also easy to mix, handle and clean up for all around optimal use in the field.

QUICK FACTS:



- Controls key alfalfa pests such as alfalfa weevil larvae, beet armyworm, cabbage looper, grasshopper and Western yellowstriped armyworm.
- Fast acting and long-lasting with up to 14 days of residual control.
- Rainfast within two hours.
- Short preharvest interval of 7 days.
- Excellent fit in integrated pest management (IPM) and insect resistance management (IRM) programs.
- Low impact on most predatory insects and mites.

Label Information

- **Active Ingredient:** Indoxacarb
- **Signal Word:** Caution
- **REI:** 12 hrs
- **Last Application:** 7 days to harvest

Precautions/Restrictions

- Do not apply more than 45 fl. oz./A or 0.44 lb. ai/A of indoxacarb-containing products per calendar year.
- Apply no more than 11.3 fl. oz./A or 0.11 lb. ai/A of indoxacarb-containing products per cutting.



The EPA label for Steward EC contains the following statements: This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area.

Stallion brand insecticide is a Restricted Use Pesticide. Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo, Stallion brand and Steward are trademarks of FMC Corporation or an affiliate. ©2022 FMC Corporation. All rights reserved. 22-FMC-2351 10/22

Application Information

- May be applied by ground, air or overhead sprinkler chemigation.
- Most effective through pest ingestion of treated plant tissue so thorough spray coverage is essential for optimum control

Crop	PESTS	APPLICATION RATE (fl. oz./A)
Alfalfa (including alfalfa grown for seed)	Cabbage looper (except CA) Grasshoppers	4.6 - 11.3
	Alfalfa caterpillar Alfalfa weevil larvae Beet armyworm Cabbage looper (CA only) Egyptian alfalfa weevil larvae Granulate cutworm	6.7 - 11.3
	Potato leafhopper* (except California) Lygus bugs (Western U.S.) * Western yellowstriped armyworm	9.2 - 11.3

Note: Apply lower listed rates for light to moderate infestations. Use intermediate to high rates within the listed rate range on heavier infestations or when later instar larvae exist. Use the highest listed rate for controlling severe infestations or when longer residual control is desired.

When Steward EC insecticide is used on alfalfa grown for seed, the seed may not be used for sprouts or livestock feed.

*Suppression only.

TO LEARN MORE ABOUT STEWARD EC INSECTICIDE, CONTACT YOUR FMC RETAILER OR VISIT AG.FMC.COM.

