
APPLICATION BEST PRACTICES FOR RHYME® FUNGICIDE IN TREE NUTS



This guide can be used as an application best practices guide for Rhyme® fungicide chemigation based on your soil type, recent rainfall or irrigation.



DISEASES

Ceratocystis

- Botryosphaeriaceae
- Cytospora
- Eutypa
- Phomopsis

Rhyme fungicide drip fall root growth

APPLICATION RECOMMENDATIONS THROUGH A PRESSURIZED IRRIGATION SYSTEM.

To help achieve maximum performance, make four applications at 7 oz./A for the year

Bearing Almond:

First application - Apply 7 fl. oz./A at the beginning of spring root flush.

Second application - Reapply at the beginning of fall root flush.

Non- bearing almond:

First application - Apply 7 fl. oz./A at the beginning of spring root flush.

Second application - Reapply within 30 day intervals.

Third application - Apply 7 fl. oz./A at the beginning of fall root flush.

Walnut:

First application - Apply 7 fl. oz./A at the beginning of spring root flush.

Second application - Reapply within 30 day intervals.

Third application - Apply 7 fl. oz./A at the beginning of fall root flush.

In California, this product may only be applied to fields that are not irrigated or are irrigated using pressurized irrigation methods such as macro-sprinkler, micro-sprinkler, or drip system.

This Rhyme fungicide recommendation is made as permitted under FIFRA Section 2(ee) for the management of wood disease complex on grapes in California. This recommendation has not been submitted to or approved by the EPA. The 2(ee) expiration date is 12/05/2024.

Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo and Rhyme are trademarks of FMC Corporation or an affiliate. ©2023 FMC Corporation. All rights reserved. 23-FMC-3200 01/23

