Tebuconazole 50% + Trifloxystrobin 25% WG

Gezeko®

Systemic Fungicide

Tebuconazole 50% + Trifloxystrobin 25% WG formulation contains 500g of Tebuconazole and 250 g of Trifloxystrobin active ingredient per kg of formulation. Tebuconzaole 50% + Trifloxystrobin 25% WG is a fungicide recommended for use as a foliar spray for the control of Shealth blight, Leaf blast & Neck blast, Glume discoloration (dirty panicle), False smut and Brown leaf spot diseases in Rice, Early blight in Tomato, Powdery Mildew in Grape; Powdery Mildew, Anthracnose & Alternaria leaf spot in Chilli, Yellow rust, Powdery mildew in Wheat; Powdery Mildew, Anthracnose in Mango, Alternaria leaf spot in Cotton, Sigatoka leaf spot in Banana and Premature leaf fall, Powdery Mildew in Apple, Alternaria leaf blight/ Leaf spot in Cabbage, Leaf blight in Maize and Powdery mildew disease in Gherkin, Cercospora leaf spot in Blackgram ,Blister Blight Tea and Tikka leaf spot in disease in Groundnut.

Chemical Composition :

Tebuconazole a.i.	:	50.00%m/m
Trifloxystrobin a.i.	:	25.00%m/m
Formaldehyde, reaction products with Sulfonated 1,1'-oxybis (methylbenzene), sodium salt	:	15.00%m/m
Naphthalenesulfonic acids, branched and linear butyl derivatives, sodium salt	:	1.00%m/m
Urea/formaldehyde condensate	:	3.00%m/m
Kaolin	:	Balance
Total	:	100.00%m/m

RECOMMENDATION FOR USE :

Crop	Common name of the disease	Dosage/per hectare			Interval between last application
		a.i (g)	Formulation in (g)	Dilution in water (Litres)	
Rice	Sheath blight, Leaf blast & Neck blast. Glume discolouration (dirty panicle)	100+50	200	375-500	21
	False smut, Brown leaf spot	175+87.5 to 200+100	350-400	500	35
Grape	Powdery Mildew	87.5+43.75	175	1000	34
Chilli	Powdery Mildew, Anthracnose & Alternaria leaf spot	125+62.5	250	500	5
Wheat	Yellow rust, Powdery Mildew	150+75	300	300-500	40
Tomato	Early blight	175+ 87.5	350	500	3
Mango	Powdery Mildew, Anthracnose	0.056%-0.075%	0.075%-0.1% (75-100 g/ 100 lit water)	Spray fluid as required depending on size of tree	15
Cotton	Alternaria leaf spot	150+75	300	500	28
Banana	Sigatoka leaf spot	175+ 87.5	350	750	20
Apple	Premature leaf fall, Powdery mildew	0.03%	0.04% (40 g/ 100 lit. water)	Spray fluid as required depending on size of tree	30
Cabbage	Alternaria Leaf blight/ leaf spot	150+75	300	500	5
Maize	Leaf blight	175+ 87.5	350	500	15
Gherkin	Powdery mildew	150+75	300	500	5
Black gram	Cercospora leaf spot	150+75	300	500	19
Tea	Blister Blight	31.25+ 62.5	125	500-foliar spraying with knapsack sprayer 80-mist blower	7
Ground nut	Tikka leaf spot	175 + 87.5	350	500	31

Caution : Not to be used on crops other than specified on this label/leaflet.

CAUTION : The product is not permitted for use near aquaculture. This product is highly toxic to aquatic organisms. Strobilurin compound is prone to resistance build up and hence continuous use should be discouraged.

RESISTANCE MANAGEMENT : Do not use fungicides with same mode of action on consecutive generation of fungal diseases. Alternate the sprays with effective fungicides having different mode of action. Do not use less / more than the recommended label rates of the product.

Tebuconazole:	Group	3	Fungicide
Trifloxystrobin:	Group	11	Fungicide

Time of application and application equipment : Tebuconazole 50% + Trifloxystrobin 25% WG is recommended to be applied with knapsack sprayer fitted with hollow cone nozzle at early appearance of disease. Depending upon the severity of disease 1-2 more spray may be given for disease management.

PRECAUTIONS : Keep locked away and out of reach of children. Avoid contact with spilled product or contaminated surfaces. Wash exposed parts of the body with soap and water after handling and before eating drinking and smoking. DO NOT eat, drink or smoke when using this product .Read label and leaflet before use and follow instructions carefully. If medical advice is needed have product container / label and leaflet at hand. DO NOT re - use this container for any other purpose . Wear personal protective equipment while working with the product, Use of Nitrile gloves, face mask and ensure protection of eyes. Wash the spray equipment's after use.

Symptoms of poisoning : Weakness, Fatigue, Dyspnea.

First aid : Get the victim into the fresh air immediately. If there is a risk of unconsciousness, position and transport in stable sideways position. Remove soiled or soaked clothing immediately and dispose of safely. In case of contact with eyes, rinse the eyes thoroughly with water. In case of contact with skin wash immediately with copious water and soap. In case of ingestion, do not induce vomiting,wash out mouth with water.Call a physician immediately.

Antidote : No specific antidote is known. Give symptomatic treatment.

Storage : (1) The packages containing fungicide shall be stored in separate room or premises away from the rooms or premises used for storing other articles or shall be kept in separate rooms / almirahs under lock and key depending upon the nature and quantity of the fungicide. (2) The rooms or premises meant for storing the fungicide shall be well - built, well lit, dry, well ventilated and of sufficient dimensions to avoid contamination with vapor / air.

Disposal of empty container : (1) The empty container shall be broken and burried away from the habitation or incinerated. (2) The used packages shall not be left outside to prevent their re-use. (3) Packages or surplus material and washing should be disposed off in safe manner so as to prevent environment and water pollution.

टेबुकोनाजोल ५०% + ट्राईफ्लोक्सिस्ट्रोबीन २५% घुलनशील कणक

गैज़को®

अन्तरप्रवाही फफूँदनाशक

टेबुकोनाजोल ५०%+ ट्राईफ्लोक्सिस्ट्रोबीन २५% घुलनशील कणक में ५०० ग्राम टेबुकोनाजोल और २५० ग्राम ट्राईफ्लोक्सिस्ट्रोबीन सक्रिय तत्व प्रति किलो फार्मुलेशन में समाविष्ट है। टेबुकोनाजोल ५०%+ ट्राईफ्लोक्सिस्ट्रोबीन २५% घुलनशील कणक धान में पर्णच्छेद अंगमारी, पत्ता झुलसा व गरदन तोड झुलसा,भद्दे सिट्टे,भूरा पर्ण धब्बा और झूठा कंडुवा रोग,टमाटर में लगने वाले अगेती अंगमारी रोग,अंगुर में लगने वाले चूर्णी फफूँदी, मिर्च में लगने वाले चूर्णी फफूँदी,पर्ण चित्ती रोग, अलटरनेरिया पत्ती धब्बा रोग, गेहूं में लगने वाले पीला गेरूआ ,चूर्णी फफूँदी, आम में लगने वाले चूर्णी फफूँदी,पर्णचित्ती, कपास में लगने वाले अलटरनेरिया पत्ती दब्बा रोग, केले की फसल में लगने वाले सिगाटोका पत्ती धब्बा रोग, सेब में समय से पूर्व पत्तों का झड़ना,चूर्णी फफूँदी रोग,गोबी में लगने वाले अल्टरनेरिया पत्ते के अंगमारी/पर्ण चित्ती,मक्के में लगने वाले पत्ते की अंगमारी, घरकिन में चूर्णी फफूँदी रोग, उडीद में सरकोस्पोरा पत्ती धब्बा रोग,चाय में फफोला अंगमारी रोग तथा मूंगफली में टिका पत्ती धब्बा रोग की रोकथाम के लिए सिफारिश किया जाता है।

रासायनिक संरचना :

टेब्युकोनाझोल ५०% + ट्राइफ्लोक्सिस्ट्रोबीन २५% डब्ल्यूजी

गैझको®

आंतरप्रवाही बुरशीनाशक

टेब्युकोनाझोल ५०% + ट्राइफ्लोक्सिस्ट्रोबीन २५% डब्ल्यूजी मधे ५०० ग्रॅम टेब्युकोनाझोल आणि २५० ग्रॅम ट्राइफ्लोक्सिस्ट्रोबीन सक्रिय तत्व प्रति किलो फॉर्म्युलेशन मधे समाविष्ट आहे. टेब्युकोनाझोल ५०% + ट्राइफ्लोक्सिस्ट्रोबीन २५% डब्ल्यू जी एक बुरशीनाशक आहे, ज्याचा वापर भातावर लागणाऱ्या लोंबी करपा, पानाचा करपा आणि नेक ब्लास्ट आणि ग्लूम डिसकलरशन (खराब पॅनिकल) टोमॅटोचा आरंभीचा करपा, (आंब्यावरील भूरी रोग आणि एन्थ्राकनोज), गव्हावरील पिवळा तांबेरा व भूरी रोग, द्राक्षा वरील भूरी रोग, मिरची वरील भूरी रोग, एन्थ्राकनाज अल्टरनेरीया सफरचंद वरील वेळेच्या आधी पान गळण व भूरी रोग, कापूस वरील अल्टरनेरिया पानावरील ठिपके, केळी वरील सिगाटोका पानावरील ठिपके, कोबी वरील अल्टरनेरिया पानावरील ठिपके मका वरील पानावरील करपा,काकडी भूरी रोग (पावडरी मिल्डयू), उडीद सरकपस्पोरा पानावरील ठिबके, चहा ब्लिस्टर ब्लाइट,भूईमुग टिक्का पानावरील ठिबके च्यानियंत्रणासाठी केला जातो.

रासायनिक संरचना :

ટેબ્યુકોનાઝોલ ૫૦% + ટ્રાયફલોક્સીસ્ટ્રોબિન ૨૫% WG

ગૈઝકો®

આંતરપ્રવાહી ફૂગનાશક

ટેબ્યુકોનાઝોલ ૫૦% + ટ્રાયફલોક્સીસ્ટ્રોબિન ૨૫% WG ફૉર્મ્યુલેશન, પ્રતિ ૧ કિ.ગ્રા., ૫૦૦ ગ્રામ ટેબ્યુકોનાઝોલ અને ૨૫૦ ગ્રામ ટ્રાયફલોક્સીસ્ટ્રોબિન ધરાવે છે. ટેબ્યુકોનાઝોલ ૫૦% + ટ્રાયફલોક્સીસ્ટ્રોબિન ૨૫% WG એક ભલામણ કરેલ ફૂગનાશક છે, જેથી ડાંગરના પાકમાં શીથબ્લાઈટ, પાનનો સુકારો તથા નેક બ્લાસ્ટ અને અંગારિયો (ફૂગનાં ગંદા ડાઘ) ટામેટામાં વહેલો સુકારો, કેરીમાં ભૂકી છારો, ફ્રુક્ષરોગ, ઘઉંમાં ભૂકી છારો, પીળો ગેરૂ, દ્રાક્ષ ભૂકી છારો, મરચાં ભૂકી છારો, ફ્રુક્ષરોગ, પાનનાં ટપકાના રોગને, ગરકીન ભુકી છારો, અડદ સર્કોસ્પોરા લીફ સ્પૉટ,ચા પાનનો સુકારો, મગફળી ટિક્કા લીફ સ્પૉટ નિયંત્રણ માં રાખવાની સફરજનમાં પૂર્ણ વિકાસ પામ્યા પહેલા પાંદડાનું ખરવું તથા સફેદ છારી, કપાસમાં પાનના ટપકાંનો રોગ કેળામાં પાનના સીગાટોકાનો રોગ, મકાઈમાં પાનનો સુકારો, કોબીમાં પાનના ટપકાંનો રોગ ભલામણ કરવામાં આવે છે.

રાસાયણિક સંયોજન :

టెబ్యుకొనాజోల్ 50% + ట్రైఫ్లాక్సీస్ట్రోబిన్ 25% డబ్ల్యుజి

గైజకో®

అంతర్వాహిక శిలీంధ్రనాశిని

டெபுகோனசோல் 50% + ட்ரைஃப்ளோக்ஸிஸ்ட்ரோபின் 25% டபிள்யூஜி

கைஜ்கோ®

ஊடுருவிச் செயல்படும் பூஞ்சாணக் கொல்லி

ٹیبوکونازول50%+ٹریفلوکسیٹروبین25%ڈبلیوجی

گیزکو®

سسٹمِک پھپھوندکش

PESTICIDE FOR AGRICULTURAL USE

Tebuconazole 50% + Trifloxystrobin 25% WG

Gezeko®

Systemic Fungicide

टेबुकोनाजोल ५०% + ट्राईफ्लोक्सिस्ट्रोबीन २५% घुलनशील कणक

गैज़को®

अर्न्तप्रवाही फफूँदनाशक

ٹیبوکونازول50%+ٹریفلوکسیٹروبین25%ڈبلیوجی

گیزکو®

سسٹمِک پھپھوندکش

IB867147787A

FMC

Marketed by: FMC India Private Limited,
TCG Financial Centre, 2nd Floor, C-53, Bandra Kurla Complex, Bandra (E), Bandra Suburban, Mumbai – 400098, Maharashtra, India.

विक्रेता : एफएमसी इंडिया प्राईवेट लिमिटेड,
टीसीजी फाइनेंशियल सेंटर, २ री मंजिल, सी–५३, बांद्रा कुर्ला कॉम्पलेक्स, बान्द्रा (पूर्व), बांद्रा सबर्बन, मुंबई – ४०००९८, महाराष्ट्रा, भारत.

વિક્રેતાઃ એફ એમ સી ઈંડિયા પ્રાઈવેટ લિમિટેડ,
ટીસીજી ફાઈનેંશિયલ સેંટર, 2 જા માળે, સી -53, બાંદ્રા કુર્લા કોમ્પલેક્સ, બાંદ્રા (પૂર્વ), બાંદ્રા સબર્બન, મુંબઈ - 400098, મહારાષ્ટ્રા, ભારત.

Manufactured by : **Bayer AG, Germany**
Imported and repacked by :
Bayer CropScience Limited,
Factory : Plot No. 203/1, G I D C, Estate, Panoli - 394116, Dist : Bharuch, Gujarat.
Regd. Office: Bayer House, Central Avenue, Hiranandani Estate, Thane (West) - 400 607.

निर्माता : **बायर एजी, जर्मनी**
आयतकर्ता तथा पुन: पैकिंगकर्ता : **बायर क्रॉपसाईन्स लिमिटेड,**
फैक्टरी : प्लॉट नं. २०३/१, जी आई डी सी इस्टेट,
पनोली – ३९४११६, जिला: भरूच, गुजरात.
पंजीकृत कार्यालय: बायर हाउस, सेंट्रल एवेन्यू,
हीरानंदानी एस्टेट, ठाणे (पश्चिम) – ४०० ६०७.

ઉત્પાદનકર્તા **: બાયર એજી, જર્મની**
આયતકર્તા અને પૂનઃ પૅકીંગકરનારઃ
બાયર ક્રોપસાયન્સ લિમિટેડ,
ફેક્ટરી : પ્લૉટ નં. ૨૦૩/૧, જી આઈ ડી સી એસ્ટેટ,
પનોલી - ૩૯૪૧૧૬, જિલ્લો : ભરુચ, ગુજરાત.
રજિ. આફિસ : બાયર હાઉસ, સેન્ટ્રલ એવેન્યુ,
હીરાનંદાની એસ્ટેટ, ઠાણે (પશ્ચિમ)- ૪૦૦ ૬૦૭.

In case of customer complaints,
please contact customer care executive at :
Phone: +91-22-6704 5404
Email : customer.care@fmc.com;
FMC India Private Limited, TCG Financial Centre, 2nd Floor, C-53,
Bandra Kurla Complex, Bandra (E),
Bandra Suburban,
Mumbai – 400098, Maharashtra, India.