



कियालो®
फ्लूप्रियम 17.7% w/w + टेबुकाज़ोल 17.7% w/w का एक विद्यमान
सर्वव्यापी फंगिसाइड (सक्रिय घटक) 17.70% w/w, फ्लूप्रियम (सक्रिय घटक) 17.70% w/w, टोबुकाज़ोल (सक्रिय घटक) 17.70% w/w, एक्सथान गम (Thickener) 0.12% w/w, डोनारे 2-methyl-2-propanol (Dispersant) 2.21% w/w, Naphthalene sulfonates condensed with formaldehyde (Dispersant) 0.09% w/w, Ethoxylated polyurethane (Dispersant) 0.09% w/w, D-glucopyranose, oligomeric, decyl octyl glycolides (Dispersant) 5.73% w/w, Poly (oxy-1,2-ethanediyl), α-isobutyryl-ω-hydroxy-Dispersant) 2.21% w/w, Water, demineralized (Solvent) 0.5. Total 100.00% w/w. Density at 20°C: 1.130 g/ml.

Net Content : 1 Litre
Systemic Fungicide

Chemical Composition : Flupyrim (Active Ingredient) 17.70% w/w, Tebuconazole (Active Ingredient) 17.70% w/w, Xanthan gum (Thickener) 0.12% w/w, Donarex 2-methyl-2-propanol (Dispersant) 2.21% w/w, Naphthalene sulfonates condensed with formaldehyde (Dispersant) 0.09% w/w, Ethoxylated polyurethane in propylene glycol and water (Dispersant) 3.54% w/w, Polyisobutylene, α-Me (Anti foam) 0.35% w/w, 1,2-benzisothiazol-3 (2 H-one) (Preservative) 0.09% w/w, 1,2-propanediol (Antifreeze) 2.21% w/w, D-glucopyranose, oligomeric, decyl octyl glycolides (Dispersant) 5.73% w/w, Poly (oxy-1,2-ethanediyl), α-isobutyryl-ω-hydroxy-Dispersant) 2.21% w/w, Water, demineralized (Solvent) 0.5. Total 100.00% w/w. Density at 20°C: 1.130 g/ml.

Use: Flupyrim 17.7% w/w + Tebuconazole 17.7% w/w SC is a systemic fungicide recommended for the control of Powdery mildew and Anthracnose diseases in Grapes and Chili. **Caution:** Do not spray near water bodies/areas where Pesticulture/Inter-cropping pisciculture is practiced. The product is toxic to fish. **Precautions:** Keep away from food, drink & animal feeding stuff and also out of reach of children. Avoid contact with skin, eyes and clothing. Wash hand thoroughly with soap and water after handling and before eating, chewing gum or tobacco. Wear personal protective equipments, while working with the product. Remove soiled clothing immediately and wash before reuse. **Symptoms of Poisoning:** Headache, Nausea, Vomiting, Dizziness, Weakness, Fatigue, Disposition, Convulsions. **First Aid Measures:** Remove contaminated clothing immediately and dispose of safely. **Inhalation:** In case of inhalation move to the fresh air and keep patient warm and rest. Call a physician immediately. **Skin Contact:** In case of skin contact, remove clothes, wash off skin with soap and plenty of water. Call a physician. **Eye Contact:** In case of contact with eye, rinse immediately with plenty of water and seek medical advice. **Ingestion:** When swallowed accidentally, do not induce vomiting. Rinse out mouth and give water in small sips to drink Get medical help. **Antidote:** No specific antidote is known. Treat symptomatically and supportively. **Storage:** 1. The packages containing fungicide shall be stored in separate room or premises, away from the rooms or premises used for storing other articles or shall be kept in separate rooms / almirahs under lock and key depending upon the nature and quantity of the fungicide. 2. The rooms or premises meant for storing the fungicide shall be well built, well lit, dry well ventilated and of sufficient dimensions to avoid contamination with vapour / air. **Disposal of Empty Container:** 1. The empty container shall be broken and buried away from the habitation or incinerated. 2. The used packages shall not be left outside to prevent their re-use. 3. Packages or surplus material and washings should be disposed off in safe manner so as to prevent environment and water pollution. **Caution:** Not to be used on crops other than specified on this label / leaflet.

KIVALO®



FUNGICIDE

FMC

1 Litre

Kivalo®
Fluopyrim 17.7% w/w + Tebuconazole 17.7% w/w SC

Net Content : 1 Litre
Systemic Fungicide

Chemical Composition : Flupyrim (Active Ingredient) 17.70% w/w, Tebuconazole (Active Ingredient) 17.70% w/w, Xanthan gum (Thickener) 0.12% w/w, Donarex 2-methyl-2-propanol (Dispersant) 2.21% w/w, Naphthalene sulfonates condensed with formaldehyde (Dispersant) 0.09% w/w, Ethoxylated polyurethane in propylene glycol and water (Dispersant) 3.54% w/w, Polyisobutylene, α-Me (Anti foam) 0.35% w/w, 1,2-benzisothiazol-3 (2 H-one) (Preservative) 0.09% w/w, 1,2-propanediol (Antifreeze) 2.21% w/w, D-glucopyranose, oligomeric, decyl octyl glycolides (Dispersant) 5.73% w/w, Poly (oxy-1,2-ethanediyl), α-isobutyryl-ω-hydroxy-Dispersant) 2.21% w/w, Water, demineralized (Solvent) 0.5. Total 100.00% w/w. Density at 20°C: 1.130 g/ml.

Use: Flupyrim 17.7% w/w + Tebuconazole 17.7% w/w SC is a systemic fungicide recommended for the control of Powdery mildew and Anthracnose diseases in Grapes and Chili. **Caution:** Do not spray near water bodies/areas where Pesticulture/Inter-cropping pisciculture is practiced. The product is toxic to fish. **Precautions:** Keep away from food, drink & animal feeding stuff and also out of reach of children. Avoid contact with skin, eyes and clothing. Wash hand thoroughly with soap and water after handling and before eating, chewing gum or tobacco. Wear personal protective equipments, while working with the product. Remove soiled clothing immediately and wash before reuse. **Symptoms of Poisoning:** Headache, Nausea, Vomiting, Dizziness, Weakness, Fatigue, Disposition, Convulsions. **First Aid Measures:** Remove contaminated clothing immediately and dispose of safely. **Inhalation:** In case of inhalation move to the fresh air and keep patient warm and rest. Call a physician immediately. **Skin Contact:** In case of skin contact, remove clothes, wash off skin with soap and plenty of water. Call a physician. **Eye Contact:** In case of contact with eye, rinse immediately with plenty of water and seek medical advice. **Ingestion:** When swallowed accidentally, do not induce vomiting. Rinse out mouth and give water in small sips to drink Get medical help. **Antidote:** No specific antidote is known. Treat symptomatically and supportively. **Storage:** 1. The packages containing fungicide shall be stored in separate room or premises, away from the rooms or premises used for storing other articles or shall be kept in separate rooms / almirahs under lock and key depending upon the nature and quantity of the fungicide. 2. The rooms or premises meant for storing the fungicide shall be well built, well lit, dry well ventilated and of sufficient dimensions to avoid contamination with vapour / air. **Disposal of Empty Container:** 1. The empty container shall be broken and buried away from the habitation or incinerated. 2. The used packages shall not be left outside to prevent their re-use. 3. Packages or surplus material and washings should be disposed off in safe manner so as to prevent environment and water pollution. **Caution:** Not to be used on crops other than specified on this label / leaflet.

KEEP OUT OF THE REACH OF CHILDREN
आणक़ोरी
दूर राहो.



वर्षों की पहुँच से दूर रखिये
1866743728A

Fluopyrim 17.7% w/w + Tebuconazole 17.7% w/w SC
 ® Registered Trade Mark of Bayer Group
 Manufactured by: **Bayer CropScience Limited**,
 Factory: Plot No. 66/1 - 75/2, GIDD Estate, Himmatnagar - 383001, Dist. Sabarkantha, Gujarat.
Regd. Office : Bayer House, Central Avenue, Hiranandani Estate, Thane (West) - 400 607

निर्माता: बायर क्रॉपसाइंस लिमिटेड,
 कारखाना: प्लॉट नं. 66/1-75/2, जीडिड एस्टेट, हिमतनगर - 383001, डिस्ट. सबारकन्था, गुजरात.
रजि. कार्यालय: बायर हाउस, सेंट्रल एवेंयू, हिरानंदानी एस्टेट, थाने (पश्चिम) - 400 607

दोस्तानेदारता से बाजार में उपलब्ध कराया जाएगा।
 भारतीय निर्यात: 17.70% w/w, फ्लूप्रियम (सक्रिय घटक) 17.70% w/w, टोबुकाज़ोल (सक्रिय घटक) 17.70% w/w, एक्सथान गम (Thickener) 0.12% w/w, डोनारे 2-methyl-2-propanol (Dispersant) 2.21% w/w, Naphthalene sulfonates condensed with formaldehyde (Dispersant) 0.09% w/w, Ethoxylated polyurethane (Dispersant) 0.09% w/w, D-glucopyranose, oligomeric, decyl octyl glycolides (Dispersant) 5.73% w/w, Poly (oxy-1,2-ethanediyl), α-isobutyryl-ω-hydroxy-Dispersant) 2.21% w/w, Water, demineralized (Solvent) 0.5. Total 100.00% w/w. Density at 20°C: 1.130 g/ml.

Marketed by: FMC India Private Limited, TCG Financial Centre,
 2nd Floor, C-53, Bandra Kuria Complex, Bandra (E),
 Bandra Suburban, Mumbai - 400089, Maharashtra, India.

कियालो® फ्लूप्रियम 17.7% w/w + टेबुकाज़ोल 17.7% w/w SC
 2 वें मंजिल, सी-53, बान्द्रा कुरिया कॉम्प्लेक्स, बान्द्रा (पूर्व), मुंबई, महाराष्ट्र, भारत.
 मुंबई - 400089, महाराष्ट्र, भारत.

कियालो® से ही कियालो® डिस्ट्रीब्यूटर्स, डीलरों को उपलब्ध कराया जाएगा।
 2 वें मंजिल, सी-53, बान्द्रा कुरिया कॉम्प्लेक्स, बान्द्रा (पूर्व), मुंबई, महाराष्ट्र, भारत.
 मुंबई - 400089, महाराष्ट्र, भारत.

In case of customer, please contact customer care executive at:
 Phone: **+91 22-6704 5404**
 Email: **Customer.care@fmc.com;**
 FMC India Private Limited, TCG Financial Centre, 2nd Floor, C-53, Bandra Kuria Complex, Bandra (E), Bandra Suburban, Mumbai - 400089, Maharashtra, India.

Regn. No. CIR-1773/2015(357)-Fluopyrim + Tebuconazole(SC)-01
 रजि. नं. 357/15

Net Content :
1 Litre

Batch No.
 बैच नं. / बैच नं.

Mfg. Date
 निर्माण तिथि / उत्पादन तारीख

Expiry Date
 अंतिम तिथि / संचयन समाप्ति तिथि

MRP: Rs. (incl. of all taxes)
 अधिकतम बिक्री मूल्य (सर्व करों सहित)
 43mm x 34mm

PDP - 161.87Sq.cm=2.5mm

*Dotted Line and dimensions not to be printed. This area should not be varnished.