HOCUSIA®
Fluopicolide 5.56% w/w + Propamocarb hydrochloride 55.6% w/w SC
(Protectant and Curative Fungicide)

Pesticide for agricultural use only

Fluopicolide 5.56% w/w + Propamocarb hydrochloride 55.6% w/w SC is a protectant and curative fungicide, recommended for use as foliar spray for the control late blight in Potato. It contains 55.6 g of Fluopicolide and 556 g of Propamocarb hydrochloride active ingredients per kg of formulation which is equivalent to 62.5 g of Fluopicolide and 625 g of Propamocarb hydrochloride active ingredients per litre of formulation.

Chemical Composition:

Component	Contents (w/w)
Fluopicolide a.i.	5.56%
Propamocarb hydrochloride a.i.	55.60%
Treated Attapulgite clay (Thickening agent)	3.91%
Comb co polymer of methylmetacrylate with polyethyleneglycol in propylene glycol + water (Emulsifier)	1.39%
Sulfonated aromatic polymer, Sodium salt (90-100%) + Sodium sulfate (5-10%) (Dispersing agent)	0.694%
Copolymer block of ethylene oxide & propylene oxide (Dispersing agent)	0.694%
Silicone oil emulsion (Antifoaming agent)	0.514%
Water (Solvent)	Q.S.
TOTAL	100.00%

Density at 20°C - 1.125 g/ml

Recommendation for Use:

Name of Crop	Common name of disease	Doses per hectare			Waiting Period between last spray to harvest (days)
		Active ingredient (gram)	Formulation (Litre)	Dilution in Water (Litre)	
Potato	Late Blight	Fluopicolide 78.13 + Propamocarb Hydrochloride 781.3 To Fluopicolide 93.8 + Propamocarb Hydrochloride 937.5	1.25 to 1.5	500	16

Caution: Not to be used on crops other than Specified on this Label leaflet.

Time of Application and Application Equipment: Give 1st spray as soon as the signs of diseases are noticed on the leaves and subsequent one or two sprays at 10-15 days interval depending on the disease severity. Spray fungicide with high volume knapsack sprayer.

Precautions: Avoid unnecessary contact with product. Wear personal protective equipment, while working with the product. Keep away from source of ignition. Keep away from food, drink & animal feeding stuff and also out of reach of children. Remove soiled clothing immediately and clean thoroughly before using again. Avoid contact with skin, eyes and clothing. Wash hand thoroughly with soap and water immediately after work.

Symptoms of Poisoning: Lethargy, ataxia, spasm

First Aid Measures: Remove contaminated clothing immediately and dispose of safely. **Inhalation:** In case of inhalation move to the fresh air and keep patient warm and rest. Call a physician immediately. **Skin Contact:** In case of skin contact, remove wash off with soap and plenty of water. Call a physician. **Eye Contact:** In case of contact with eye, rinse immediately with plenty of water and seek medical advice. **Ingestion:** When swallowed accidentally, do not induce vomiting. Get medical help.

Antidote: Treat symptomatically and supportively.

Storage: 1. The packages containing fungicide shall be stored in separate room or premises, away from the rooms or premises used for storing other articles or shall be kept in separate rooms / almirahs under lock and key depending upon the nature and quantity of the fungicide. 2. The rooms or premises meant for storing the fungicide shall be well built, well lit, dry, well ventilated and of sufficient dimensions to avoid contamination with vapour/ air.

Disposal of empty Container: 1. The empty container shall be broken and buried away from the habitation or incinerated. 2. The used packages shall not be left outside to prevent their re-use. 3. Packages or surplus material and washings should be disposed-off in safe manner so as to prevent environment and water pollution.

Manufactured by: **Bayer CropScience Limited.** Factory: Plot No. 661 - 702, GIDC Estate, Himatnagar - 383001, Dist. Sabarkantha, Gujarat. **Regd. Office:** Bayer House, Central Avenue, Hiranandani Estate, Thane (West) - 400 607. **Marketed by : FMC India Private Limited.** TCG Financial Centre, 2nd Floor, C-53, Bandra Kurla Complex, Bandra (E), Bandra Suburban, Mumbai – 400098, Maharashtra, India.

होकुसिया®
फ्लुओपिकोलाईड ५.५६% भा/भा+ प्रोपामोकार्व हाईड्रोक्लोराइड ५५.६% भा/भा विलेय सान्द्रण
(रक्षात्मक एवं साध्य फफूँदनाशक)

केवल कृषि उपयोग के लिए

[illegible]

होकुसिया®
फ्लुओपिकोलाईड ५.५६% भा/भा + प्रोपॅमोकार्ब हायड्रोक्लोराईड ५५.६% भा/भा एससी
(संरक्षक आणि उपचारात्मक बुरशीनाशक)

फक्त शेतीय उपयोगाकरिता

[illegible]

હોકુસિયા®
ફ્લુપિકોલાઇડ ૫.૫૬% ભાર/ભાર + પ્રોપામોકાર્બ હાઈડ્રોક્લોરાઈડ ૫૫.૬% ભાર/ભાર એસસી
(સંરક્ષક અને રોગહર ફુગનાશક)

ફક્ત ખેતીવાડીના ઉપયોગ માટે

[illegible]

హొకుసియా®

[illegible]

ہوکسیہ®
فلوپیکولائیڈ 5.56% وزن/وزن+ پروپاموکارب ہائیڈروکلورائیڈ 55.6% وزن/وزن ایس سی
(محافظ اور شفابخش فطر کش)

صرف زرعی استعمال کے لیے

[illegible]

INSTRUCTIONS FOR USE

PESTICIDE FOR AGRICULTURAL USE ONLY

HOCUSIA®
Fluopicolide 5.56% w/w + Propamocarb hydrochloride 55.6% w/w SC
(Protectant and Curative Fungicide)

Pesticide for agricultural use only

होकुसिया®
फ्लुओपिकोलाईड ५.५६% भा/भा+ प्रोपामोकार्व हाईड्रोक्लोराइड ५५.६% भा/भा विलेय सान्द्रण
(रक्षात्मक एवं साध्य फफूँदनाशक)

केवल कृषि उपयोग के लिए

હોકુસિયા®
ફ્લુપિકોલાઇડ ૫.૫૬% ભાર/ભાર + પ્રોપામોકાર્બ હાઈડ્રોક્લોરાઈડ ૫૫.૬% ભાર/ભાર એસસી
(સંરક્ષક અને રોગહર ફુગનાશક)

ફક્ત ખેતીવાડીના ઉપયોગ માટે

FMC

IB873109738

Manufactured by: **Bayer CropScience Limited.** Factory: Plot No. 661 - 702, GIDC Estate, Himatnagar - 383001, Dist. Sabarkantha, Gujarat. **Regd. Office:** Bayer House, Central Avenue, Hiranandani Estate, Thane (West) - 400 607. **Marketed by : FMC India Private Limited.** TCG Financial Centre, 2nd Floor, C-53, Bandra Kurla Complex, Bandra (E), Bandra Suburban, Mumbai – 400098, Maharashtra, India.

[illegible]

[illegible]

In case of customer complaints, please contact customer care executive at :

Phone 1800 102 6545

Email Ask@fmc.com

FMC India Private Limited, TCG Financial Centre, 2nd Floor, C-53, Bandra Kurla Complex, Bandra (E), Bandra Suburban, Mumbai – 400098, Maharashtra, India.

Size : 385mm (w) x 240mm (h)

ಹೊಕುಸಿಯಾ®

ಫ್ಲುಪಿಕೊಲೈಡ್ 5.56% ಡಬ್ಲ್ಯು/ಡಬ್ಲ್ಯು +
ಪ್ರೊಪಮೊಕಾರ್ಬ್ ಹೈಡ್ರೊಕ್ಲೋರೈಡ್ 55.6% ಡಬ್ಲ್ಯು/ಡಬ್ಲ್ಯು ಎಸ್‌ಸಿ
(ಸಂರಕ್ಷಕ ಮತ್ತು ಪರಿಹಾರಕ ಶಿಲೀಂಧ್ರನಾಶಕ)
ಕೃಷಿಯ ಉಪಯೋಗಕ್ಕಾಗಿ ಮಾತ್ರ

[illegible]

ஹோகுஸியா®

ஃப்ளூபிகோலைட் 5.56% w/w +
ப்ரோபமோகார்ப் ஹைட்ரோக்ளோரைட் 55.6% w/w SC
(ப்ரோடெக்டென்ட் மற்றும் க்யூரேடிவ் ஃபங்கிஸைட்)
விவசாய பயன்பாட்டிற்கு மட்டுமான

[illegible]

ഹോകുസിയാ®

ഫ്ലൂവോപ്പിക്കോളൈഡ് 5.56% ഡബ്ലിയു/ഡബ്ലിയു +
പ്രൊപ്പമോകാർബ് ഹൈഡ്രോക്ലോറൈഡ്
55.6% ഡബ്ലിയു/ഡബ്ലിയു എസ് സി
(പ്രതിരോധവും പ്രതിവിധിയുമായ കുമിൾനാശിനി)
കാർഷികാവശ്യത്തിനു മാത്രമുള്ള

[illegible]

ਹੋਕੁਸਿਯਾ®

ਫਲੂਓਪਿਕੋਲਾਇਡ ੫.੫੬% ਡਬ/ਡਬ +
ਪ੍ਰੋਪਾਮੋਕਾਰਬ ਹਾਈਡ੍ਰੋਕਲੋਰਾਈਡ ੫੫.੬% ਡਬ/ਡਬ ਐਸਸੀ
(ਰੱਖਿਆਤਮਕ ਅਤੇ ਅਸਰ ਕਰਨ ਵਾਲਾ ਉੱਲੀਨਾਸ਼ਕ)
ਸਿਰਫ਼ ਖੇਤੀਬਾੜੀ ਵਿਚ ਵਰਤਣ ਲਈ

[illegible]

হোকুসিয়া®

ফ্লুওপিকোলাইড ৫.৫৬% ডব্লিউ / ডব্লিউ +
প্রোপামোকার্ব হাইড্রোক্লোরাইড
৫৫.৬% ডব্লিউ / ডব্লিউ এসসি
(ছত্রাক প্রতিরোধক এবং প্রতিষেধক ঔষধ)
শুধুমাত্র কৃষিকাজে ব্যবহারের জন্যে

[illegible]

হোকুচিয়া®

ফ্লুপিকোলাইড ৫.৫৬% ডব্লিউ / ডব্লিউ +
প্ৰ'পামোকাৰ্ব হাইড্ৰ'ক্লোৰাইড ৫৫.৬% ডব্লিউ / ডব্লিউ এচচি
(সুৰক্ষিত আৰু প্ৰতিকাৰক ভেঁকুৰনাশক)
কেৱল কৃষিভিত্তিক ব্যৱহাৰৰ বাবে

[illegible]

ହୋକୁସିଆ®

ଫ୍ଲୁଓପିକୋଲାଇଡ ୫.୫୬% ଡବ୍ଲୁ/ଡବ୍ଲୁ +
ପ୍ରୋପାମୋକାର୍ବ ହାଇଡ୍ରୋକ୍ଲୋରାଇଡ ୫୫.୬% ଡବ୍ଲୁ/ଡବ୍ଲୁ ଏସ୍‌ସି
[illegible]
[illegible]

[illegible]

Size : 385mm (w) x 240mm (h)