

Molybdenum 250

A foliar applied source of plant available molybdenum

CONTAINS

Molvbdenum (Mo): 250 a/l

RATES OF USE AND TIMING

Beans and peas: Apply 0.25 I/ha when plants are 5 to 15 cm tall. For moderate to severe deficiency, repeat at 10 to 14 day intervals from the first

Broccoli, cabbage, calabrese: Apply 0.25 |/ha from 4 to 6 leaf stage. For moderate to severe deficiency, repeat at 10 to 14 day intervals.

Oilseed rape: Apply 0.25 I/ha at the 4 to 6 stage followed by 0.25 I/ha at onset of stem extension. A further application can be made at 10 to 14 days later for a severe deficiency. Avoid application during flowering.

Cucurbits: Apply 0.25 I/ha at the end of the vegetative growth before the flowering; repeat if needed after the fruit set.

Fruit trees (avocado, macadamias, cashew, etc...), coffee, cotton: Apply 0.25 I/ha before flowering.

Onions: Apply 0.25 I/ha as soon as there is sufficient leaf cover to intercept the spray. Repeat two weeks later if necessary. In case of moderate to severe deficiency, repeat at 10 to 14 day intervals.

Water volume: Apply in at least 200 litres of water per hectare.

Best results will be obtained when Molybdenum 250 is applied during the early morning or evening when moisture is present in the plant. Do not apply in extremes of temperature, very bright sunlight or when the crop is under drought or other stress.

APPLICATION

Shake the container well before opening and use.

Spray equipment should be clean.

Half fill the spray tank with water, begin agitation and add the required quantity of Molybdenum 250. Rinse the container thoroughly, add the rinsings to the spray-tank, complete the filling and apply without delay. Maintain agitation throughout.

Clean sprayer thoroughly after use.

COMPATIBILITY

Molybdenum 250 is physically and chemically compatible with a wide range of agrochemical products. Consult your supplier for the latest information. Up-to-date information is available at www.fmc-agro.co.uk. Always read the label of the partner product carefully before mixing with Molvbdenum 250.

STORAGE

Protect from frost. Store above 5°C.

Keep in cool, well ventilated suitable store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs.

SAFETY PRECAUTIONS

H315: Causes skin irritation.

H319: Causes serious eye irritation.

Warning

P264: Wash hands thoroughly after handling.

P337+313: If eve irritation persists: Get medical advice.

P280: Wear protective gloves/protective clothing/eve protection/face protection.

P302+352: IF ON SKIN: Wash with plenty of soap and water.

P305+351+338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

P332+313: If skin irritation occurs: Get medical advice/attention.

Made in UK

Batch No and DoM: See neck of the bottle

Expiry date: 3 years after the manufacturing date.

Imported and Distributed by:



Hygrotech (EA) Ltd, P.O. Box 41446, 00100, Nairobi, Kenya Naivasha: +254 51 221 2184 Limuru: +254 20 205 3916



TRIACHEM TANZANIA LIMITED PLOT NO 49AA, P.O.BOX 2750. ESSO ROAD, INDUSTRIAL AREA. ARUSHA- TANZANIA PHONE: +255272547485/8774(FAX) WEB: WWW.TRIACHEM.COM

TFRA REG NO: 0146

FMC

270 x 130 - 11001875 22056443 LBL SL-F 1L MOLYBDENUM 250 TRI TZ KE 1 SINGLE FRONT.indd 1

21/11/2022 15:09









FILL IN DIMENSIONS (WXH) KENYA 270 x 130 mm 11001875 Base for Booklet MOLYBDENUM 250 1L TZ KE TRI-HYG Booklet 22056443 LBL_SL-F_1L_MOLYBDENUM 250_TRI TZ KE_1 Carton label Preprinted Material