



DANGER

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT:

Contains 69 g/L fenoxaprop-P-ethyl in the form of an oil in water emulsion.

Also contains 34.5 g/L cloquintocet-mexyl as a safener and 390 g/L hydrocarbon liquid.

CONTENTS:

10 L

GROUP A HERBICIDE

For selective grass weed control in autumn and spring sown Barley, Wheat and Perennial Ryegrass and in Pasture.

FIRST AID

If medical advice is needed, have product container or label at hand. If swallowed immediately call the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. **DO NOT** induce vomiting. If skin contact occurs, get medical advice/attention. If on skin wash with plenty of soap and water. If skin irritation or rash occurs get medical advice/attention. Get medical advice/attention if you feel unwell.

PLEASE READ THE LABEL COMPLETELY BEFORE USE

Approved pursuant to the HSNO Act 1996, Approval Code HSR100112
See www.epa.govt.nz for approval conditions.
Registered pursuant to the ACVM Act 1997, No. P008030
See www.foodsafety.govt.nz for registration conditions.



HSNO Classifications: 6.1E, 6.3B, 6.5B, 6.9B, 9.1C, 9.2D

WARNING

DANGER: May be harmful if swallowed and enters airways. Causes mild skin irritation. May cause an allergic skin reaction. May cause damage to organs through prolonged or repeated exposure.

Harmful to aquatic life with long lasting effects. Harmful to the soil environment. Avoid release to the environment.

DO NOT apply directly into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

HANDLING PRECAUTIONS

Keep out of reach of children. Read label before use. **This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.**

PERSONAL PROTECTION

DO NOT breathe vapours. May be harmful if swallowed, inhaled or absorbed through the skin. Avoid skin contact or inhalation. Avoid contact with the eyes. **DO NOT** eat, drink or smoke while using. When preparing spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat and elbow-length PVC gloves.

Remove protective clothing and wash hands and face thoroughly before meals and after work. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

STORAGE:

Store in original container tightly closed and in a locked, dry, cool, well ventilated area, away from feed, seeds and foodstuffs. Store in accordance with NZS 8409 Management of Agrichemicals. Stores containing more than 1000 L require secondary containment, are subject to signage, and require emergency response plans. When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of this product if it is older than this.

SPILLS AND DISPOSAL

When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

Disposal: Container disposal: Triple rinse empty container adding rinsate to the spray tank. Recycling rinsed container through Agrecovery (0800 247 326, www.agrecovery.co.nz).

Otherwise crush or puncture and bury in an approved landfill. **DO NOT** reuse container for any other purpose. Dispose of this product only by using according to this label. Always follow local Authority requirements.

GENERAL INFORMATION

WEEDS CONTROLLED: Wild oats, Lesser canary grass (*Phalaris minor*), Yellow bristle grass.

RESISTANT WEEDS: Gnawed canary grass (*Phalaris paradoxa*), Annual and hybrid ryegrasses, Tall fescue, Chewing's fescue, Cocksfoot, Browntop, Grazing brome (=spiky brome – *Bromus stamineus*), Prairie grass and most broadleaved weeds are not controlled by FOXTROT® at label rates.

DIRECTIONS FOR USE

CROP	WEEDS	RATE	Remarks
Barley, Wheat	Wild oats	750 mL/ha in 250 L of water. Add adjuvant oil at recommended rates	Apply when wild oats are between 3 leaf and 1 st node visible growth stage and when they are growing actively.
	Lesser canary grass (<i>Phalaris minor</i>)		Apply when weeds are between the 3 leaf visible stage and the end of tillering and are growing actively. Avoid application during periods of moisture stress.
Perennial ryegrass (<i>Lolium perenne</i>)	Wild oats	750 mL/ha in 250 L of water. DO NOT use with any other agrochemicals or adjuvants.	Apply in early spring when wild oats are between 3 leaf and end of tillering growth stage and growing actively. Note: Seed yield response is likely to be reduced the longer the delay in the removal of wild oats in spring.
Pasture	Yellow bristle grass	Use 750 ml/ha in 250 L of water. DO NOT use with any other agrichemicals or adjuvants.	Apply to established pasture 7 days after grazing during the period December/January.

WITHHOLDING PERIODS

Barley and Wheat 70 days (**DO NOT** graze treated crop for at least 70 days after treatment).

Perennial ryegrass 42 days (**DO NOT** graze treated crop for at least 42 days after treatment).

Pasture 28 days (**DO NOT** graze treated pasture for at least 28 days after treatment).

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MIXING

Shake well before use. Add the required amount to a partly filled spray tank with agitation system operating. Complete filling the tank with water and apply.

COMPATIBILITY

DO NOT USE FOXTROT® IN ANY MIXTURES WHEN APPLYING TO PERENNIAL RYEGRASS. When used on barley and wheat, FOXTROT® can be tank mixed with Cycocel* and Opus* in 2 and 3 way mixes. Care must be taken to ensure that recommended timings for all products are followed. The addition of a suitable adjuvant oil such as D-C-Trate™ or Hasten™ at label rates is recommended. **DO NOT use FOXTROT® within 5 days of broad-leaved herbicide applications.**

APPLICATION

DO NOT overlap swath width in ryegrass crops as yield suppression may occur. Flush spray system with clean water after use.

RAINFASTNESS

FOXTROT® is rainfast within one hour of application.

SOIL ACTIVITY

FOXTROT® is rapidly broken down by soil microbes causing no restrictions on crop rotations.

UNDERSOWN CROPS

FOXTROT® can be used on crops undersown with certain grasses (See under "Resistant Weeds" above) and clover. **DO NOT use on crops undersown with "Concord" ryegrass.**

RESISTANT WEEDS WARNING:

GROUP **A** HERBICIDE

FOXTROT® Selective Herbicide is a member of the aryloxyphenoxypropionate group of herbicides. FOXTROT® is a herbicide which inhibits production of acetyl CoA carboxylase. For weed resistance management FOXTROT® is a Group A herbicide. Some naturally occurring weed biotypes resistant to FOXTROT®, and other Group A herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FOXTROT® or other Group A herbicides. See www.resistance.nzpps.org for more information.

SHELF LIFE:

When stored appropriately, this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

SAFETY DATA SHEET

Additional information is listed in the SDS available from FMC New Zealand Limited or www.fmccrop.nz

NOTICE TO BUYER

FMC warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use of application, all of which are beyond the control of FMC. To the extent permitted at law FMC shall not be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer.

TO THE EXTENT PERMITTED AT LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

FMC and Foxtrot® are trademarks of FMC Corporation or an affiliate.

© 2020 FMC Corporation. All rights reserved.

*Non FMC Trademark

Registered and Distributed by:

FMC New Zealand Limited
6 Clayton Street
Newmarket
Auckland 1023
New Zealand
Freephone: 0800 65 80 80