

CONTENTS: 5 L

WARNING
Keep out of reach of children
ECOTOXIC

SPORTAK® EW

Fungicide for disease control in
Avocados, Barley, Wheat,
Ornamentals and Radiata pine

Active ingredient: Contains 450 g/L
prochloraz in the form of a emulsion
in water
Also contains 60 g/L aromatic
hydrocarbons



GROUP	3	FUNGICIDE
-------	----------	-----------

**READ LABEL AND LEAFLET BEFORE
MIXING AND APPLICATION**



Registered pursuant to the ACVM Act 1997, No P7750.
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No HSR007837.
See www.epa.govt.nz for approval conditions.



HSNO Classifications: 6.1E, 6.9B, 9.1A, 9.3C

This product must be under control of an approved handler during use.

25-04-2019

WARNING: May be harmful if swallowed. May cause organ damage through prolonged repeated oral exposure at high doses.

ECOTOXIC: Toxic to fish with long-lasting effects. Harmful to terrestrial vertebrates.

PRECAUTIONS:

PERSONAL PROTECTION:

DO NOT eat, drink or smoke while using. Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using. Remove protective clothing and wash hands and face thoroughly before meals and after work. May be harmful if swallowed, inhaled or absorbed through the skin. Avoid skin contact or inhalation. Avoid contact with the eyes.

STORAGE: Store in original container tightly closed and in a locked dry, cool, well ventilated area, away from food stuffs. When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of this product if it is older than this.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

Disposal: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

Container Disposal: Triple rinse container and add residue to mixture. Take to an Agrecovery collection site.

FIRST AID:

If swallowed **DO NOT** induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water, if irritation occurs seek medical advice. If splashed in eyes, wash out immediately with water, if irritation persists get medical advice. If inhaled move the victim to fresh air immediately.

Sportak® EW is a DMI fungicide with eradicator and protectant activity against certain diseases in Avocados, Barley, Wheat, Ornamentals and Radiata pine.

RESISTANCE MANAGEMENT

GROUP	3	FUNGICIDE
-------	----------	-----------

Resistance to this and other DMI fungicides could develop from repeated use. To minimise this risk, use strictly in accordance with label recommendations. Use a maximum of three DMI fungicides per crop, per season, at full label rates.

It is an offence to use this product on animals.

DIRECTIONS FOR USE

Mixing: Add the required amount to the partly filled spray tank with the agitation system operating. Complete filling of spray tank with water and maintain agitation until spraying is complete. Agitate thoroughly before restarting after a stoppage.

Compatibility: Compatible with most commonly used herbicides, fungicides and insecticides.

DO NOT mix with strongly alkaline materials such as Bordeaux mixture and lime sulphur. When used in mixture care should be taken not to compromise timing or method of application for all components of the tank mix. **DO NOT** use in mixture when treating Avocados.



A handwritten signature in blue ink, appearing to be "Mark Gillingham".

25-01-2019

RECOMMENDATIONS:

Crop	Disease	Rate	Remarks
Avocados	Anthraco nose	Post-harvest dip: 110 mL/200 L of water Post-harvest spray: 55 mL/100 L of water	Immerse fruit for 30 seconds. Treat no more than 2 tonnes of fruit (approx 400 packed trays) per 200 L dip solution. Discard used dip solution and replace if further fruit to be treated. Spray fruit for 1 minute in a non-recirculation system. DO NOT re-use solution. NB Thorough coverage of the fruit is essential. Treat fruit at ambient temperature as soon as practicable after picking. After treatment allow fruit to drain and dry before packing. The rate at which the active ingredient is removed is unknown therefore discarding used dip/spray solution is essential. Dirty fruit should be dipped last. DO NOT treat visibly diseased fruit.
Barley & Wheat	Eyespot, Glume blotch, Ear disease complex	1 L/ha	Apply from mid to late tillering (Zadoks GS26) to flag leaf just visible (Zadoks GS37). Apply as soon as the flag leaf ligule is visible to protect the flag leaf from infection. Apply when the flag leaf is fully emerged (Zadoks GS 58).
Barley	Leaf scald, Net blotch, Spot blotch	1 L/ha	Apply at the first signs of disease or as a preventative treatment, when conditions favour infection.
Wheat	Speckled leaf blotch	1 L/ha	Apply as a preventative treatment between first true leaf (Zadoks GS 11) and tillering (Zadoks GS 21) or at the first sign of infection. A second application may be required at ear emergence (Zadoks GS 51) to protect the flag leaf.
Ornamentals	<i>Colletotrichum</i> sp Glomerella leaf spot, <i>Monochaetia</i> sp, <i>Pestalotia</i> sp, Powdery mildew	100 ml/100 L in sufficient water to obtain thorough spray coverage.	Apply at first signs of infection and repeat at 14 day intervals.
		100 ml/100 L as a dip	Immerse whole plants in dip treatment prior to shipping.
Radiata Pine seedlings	Terminal crook disease	1.5 L/ha	Apply as a protectant spray at 7 - 14 day intervals. Use the shorter spray interval under severe infection pressure.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

**WITHHOLDING PERIOD:
BARLEY AND WHEAT: 42 DAYS**



[Handwritten signature]
25-01-2019

NOTICE TO BUYER

FMC warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use of application, all of which are beyond the control of FMC. To the extent permitted at law FMC shall not be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer.

TO THE EXTENT PERMITTED AT LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE

The FMC logo and Sportak® are Trademarks or Registered trademarks of FMC or its affiliates.

Formulated for and distributed by:

FMC New Zealand Limited
6 Clayton Street
Newmarket
Auckland 1023
Freephone: 0800 65 80 80
www.fmccrop.nz

New Zealand Emergency Telephone
Transport Emergency: 0800 387 668
24 hr Emergency Medical Information: 0800 111 174

