






Verimark[®] 20SC

PRODUCT INFORMATION

PRODUCT NAME	Verimark [®] 20SC
ACTIVE INGREDIENT	Cyantraniliprole
CONCENTRATION & FORMULATION	200g/L SC
USE	Insecticide
Mode of Action (MoA)	28
FPA TOX CATEGORY	4
Color Band	Green Label
FPA Registration No.	598-485-4456
CROP REGISTERED	Tomato, Bittergourd, Onion
Systemicity	Systemic
Pest group specificity	Chewing & Sucking Insects
Recommended Timing of Application (relative to insect pest occurrence)	Preventive & Curative

Verimark[®] 20SC is a systemic insecticide containing Cyantraniliprole powered by Cyazypyr[®] belonging to IRAC Group 28 for the control of chewing and sucking insect pests like Cutworm, Whitefly and Leafminer in Tomato, Aphids and Melon worm in Bittergourd and Cutworm in Onion.

DIRECTIONS FOR USE: Verimark® 20SC is a systemic insecticide containing Cyantraniliprole powered by Cyazypyr® active belonging to IRAC Group 28 for the control of chewing and sucking insect pests for the following crops:

Crop	Pests	Use Rate per 16L water	Product Rate Per Hectare	Spray Volume	Application Timing
 Tomato	Cutworm, Whitefly, Leafminer	112ml	700ml	100 L/Ha	2 DBT
 Bittergourd	Aphids, Melon worm	96 - 112ml	1120ml	160 L/Ha	2 DBT
 Onion	Cutworm	30ml	1125ml	600 L/Ha	2 DBT

It is recommended to treat seedlings ONLY ONCE, 2 days before transplanting as protection against early insect pests up to 21 days after treatment application. Apply follow up foliar spray 14-21 days after the pre-plant treatment using another effective compound registered for the control of the above pests damaging the listed crops belonging to different mode of action.