






PRODUCT INFORMATION




PRODUCT NAME	Gezeko® 75WG
ACTIVE INGREDIENT	Tebuconazole + Trifloxystrobin
CONCENTRATION & FORMULATION	500g/kg + 250g/kg WG
USE	Fungicide
Mode of Action (MoA)	3 + 11
FPA TOX CATEGORY	4
Color Band	Green label
FPA Registration No.	598-438-4838
CROP REGISTERED	Potato, Eggplant, Bittergourd, Tomato, Watermelon, Onion, Corn, Mango, Rice
Re-entry Period	12 hours after spraying
Systemicity	Systemic & Mesostemic
Pest group specificity	Fungal diseases
Recommended Timing of Application (relative to disease occurrence)	Preventive & Curative

Gezeko® 75WG

Gezeko® 75WG is a systemic fungicide with the combined power of Tebuconazole and Trifloxystrobin belonging to FRAC Group 3 & 11, respectively under Tox Category 4 (green label) registered for control of various diseases in Potato, Eggplant, Bittergourd, Tomato, Watermelon, Onion, Corn, Mango, Rice

DIRECTIONS FOR USE: Gezeko® 75WG is a systemic fungicide with the combined power of Tebuconazole and Trifloxystrobin belonging to FRAC Group 3 & 11, respectively under Tox Category 4 (green label) registered in the following crops and target diseases.

Crop	Pests	Use Rate per 16L water	Product Rate Per Hectare	Application Timing	Spray Interval	Spray Volume	PHI	Re-Entry Period
 Mango	Anthraco nose	80 - 160g per 200L water	1600g	Apply at first sign of disease infection and repeat after 7 days	7 days	2000L/ha	30 days	As soon as spray deposits have dried
 Rice	Rice blast, Sheath Blight, Brown leaf spot, Dirty Panicle	12 - 16g	160g	Apply at leaf development against early blast attack or if no infection, apply at maximum tillering stage and repeat after 7 days	7 days	160L/ha	35 days	As soon as spray deposits have dried
 Bittergourd	Powdery mildew, Downy Mildew	6 - 8g	200g	Apply at first sign of disease infection and repeat after 7 days	7 days	400L/ha	3 days	As soon as spray deposits have dried
 Eggplant	Cercospora leaf spot	6 - 8g	200g	Apply at first sign of disease infection and repeat after 7 days	7 days	400L/ha	3 days	As soon as spray deposits have dried
 Tomato	Early Blight	6 - 8g	200g	Apply at first sign of disease infection and repeat after 7 days	7 days	400L/ha	3 days	As soon as spray deposits have dried
 Watermelon	Powdery mildew, leafspot	4 - 8g	200g	Apply at first sign of disease infection and repeat after 7 days	7 days	400L/ha	5 days	As soon as spray deposits have dried

Crop	Pests	Use Rate per 16L water	Product Rate Per Hectare	Application Timing	Spray Interval	Spray Volume	PHI	Re-Entry Period
 Corn	Corn rust, Helminthosporium leaf spot	4 - 8g	200g	Apply at first sign of disease infection or about 3 weeks after planting and repeat after 7 days	7 days	400L/ha	5 days	As soon as spray deposits have dried
 Onion	Purple blotch	4 - 8g	200g	Apply at first sign of disease infection or about 3 weeks after planting and repeat after 7 days	7 days	400L/ha	5 days	As soon as spray deposits have dried
 Potato	Early and Late Blight	4 - 8g	200g	Apply at first sign of disease infection or about 15 days after planting and repeat after 7 days	7 days	400L/ha	3 days	As soon as spray deposits have dried

Apply a maximum of 2 sprays of Gezeko® 75WG applied as back-to-back in one cropping season, then alternate with other products belonging to a different fungicide chemistry and registered for the same pest and crop.