






Rovral® Aquaflo 50SC

PRODUCT INFORMATION

PRODUCT NAME	Rovral® Aquaflo 50SC
ACTIVE INGREDIENT	Iprodione
CONCENTRATION & FORMULATION	500g/L SC
USE	Fungicide
Mode of Action (MoA)	Group 2
FPA TOX CATEGORY	3
FPA Registration No.	598-155-4188
CROP REGISTERED	Rice, Banana, Turf
Re-entry Period	As soon as spray deposits have dried
Systemicity	Contact & Locally Sytemic
Pest group specificity	Fungal diseases
Recommended Timing of Application (relative to disease occurrence)	Preventive & Curative

Rovral® Aquaflo 50SC is a broad spectrum contact and locally systemic fungicide containing Iprodione belonging to FRAC Group 2 that has protectant and eradicant properties for the control of various diseases

DIRECTIONS FOR USE: Rovral® Aquaflo 50SC is a broad spectrum contact and locally systemic fungicide containing Iprodione belonging to FRAC Group 2 that has protectant and eradicant properties for the control of various diseases registered in rice, banana and turf

Crop	Pests	Use Rate per 16L water	ProductRate Per Hectare	Application Timing	Spray Interval	Spray Volume	PHI	Re-Entry Period
 Rice	Sheath Blight (Rhizoctonia solani)	60ml per 16L water	600ml	25 DAP & 35 DAP	10 days	160L/ha	14 days	As soon as spray deposits have dried
 Banana	Sigatoka	1.5L / hectare	1,500ml	2 to 3 months after planting	8 - 10 days	30 - 36L/ha	0 days	As soon as spray deposits have dried
 Banana	Crown mold, Crown Rot	32ml per 16L water	N/A	before packing	N/A	N/A	0 days	N/A
Turf Grass	Fairy Ring	100ml per 16L water	6000ml	Curative treatment: 7 to 10 days interval depending on the severity of infection; For Preventiv Treatment - apply as a monthly spray program	7 to 10 days	960L/ha		As soon as spray deposits have dried

For Rice: First application at booting stage or when 5% of panicles have emerged. If favorable disease conditions continue, apply 1 week later but before 75% heading. Do not use more than 3 applications per growing season.

For Banana: Sigatoka Control - Apply 4-5 liters of banana oil and mix thoroughly. Aerial application should be done every 8 to 10 days at a spray volume of 30 to 36 liters per hectare. Crown rot & Mold control - Measure required quantity and mix to appropriate volume of water. Agitate to mix solution thoroughly. Use 250 to 300ml of the solution per 13kg fruit.