










## Steward® 30WDG

### PRODUCT INFORMATION

PRODUCT NAME	Steward® 30WDG
ACTIVE INGREDIENT	Indoxacarb
CONCENTRATION & FORMULATION	300g/Kg WDG
USE	Insecticide
Mode of Action (MoA)	Indoxacarb (MoA 22A)
FPA TOX CATEGORY	3
Color Band	Blue Label
FPA Registration No.	598-353-4412
CROP REGISTERED	Cabbage, Bittergourd, Eggplant, Stringbeans, Corn, Tobacco, Rice
Re-entry Period	As soon as spray mist dries up
Systemicity	Contact
Pest group specificity	Chewing Insects
Recommended Timing of Application (relative to insect pest occurrence)	Preventive

Steward® 30WDG is a contact insecticide containing Indoxacarb belonging to IRAC Group 22A under Tox category 3 (Blue label) for the control of chewing insect pests like Diamondback moth in Cabbage, Leafroller and Cutworm in Bittergourd, Fruit and Shootborer (FSB) in Eggplant, Podborer in Stringbeans, Budworm and Cutworm in Tobacco and Fall armyworm (FAW) in Corn. Steward® 30WDG is recommended to be applied as a preventive control.

**DIRECTIONS FOR USE:** Steward® 30WDG is a contact insecticide containing Indoxacarb belonging to IRAC Group 22A under Tox category 3 (Blue label) for the control of chewing insect pests as indicated below. Best timing of application is during vegetative stage or as alternate product for resistance management during heading / fruiting stage.

Crop	Pests	Use Rate per 16L water	Product Rate Per Hectare	Spray Interval	Spray Volume	PHI	Re-Entry Period
 Cabbage	Diamondback Moth	3.0 - 3.5g	80g	7 days	320 L/Ha	1 day	As soon as spray mist dries up
 Eggplant	Fruit & Shoot Borer (FSB)	4.0 - 5.0g	100g	7 days	320 L/Ha	3 days	As soon as spray mist dries up
 Bittergourd	Leaffolder, Cutworm	4.0 - 5.0g	100g	7 days	320 L/Ha	1 day	As soon as spray mist dries up
 Stringbeans	Podborer	4.0 - 5.0g	100g	7 days	320 L/Ha	3 days	As soon as spray mist dries up
 Tobacco	Budworm, Cutworm	4.0g	125g	7 days	500 L/Ha	1 day	As soon as spray mist dries up
 Rice	Leaffolder	2.0g	40g	7 days	320 L/Ha	3 days	As soon as spray mist dries up
 Corn	Fall Armyworm	24g	360g	7 days	240 L/Ha		As soon as spray mist dries up

Apply a maximum of 2 sprays of Steward® 30WDG is applied back-to-back in any one crop cycle, targeted at a single pest generation and then alternate with other products belonging to a different insecticide chemistry and registered for the same pest and crop.