

OPEN HERE

SIMBA® SX®**Herbicide**contains 200 g/kg
metsulfuron-
methyl**WARNING****Very toxic to aquatic life with
long lasting effects.**

Collect spillage.

Dispose of contents / container to a licensed
hazardous waste-disposal contractor or collection
site except for empty clean containers which may be
disposed of as non-hazardous waste.Do not contaminate water with the product or its
container (Do not clean application equipment near
surface water/Avoid contamination via drains from
farmyards and roads).**To avoid risks to human health and the environment,
comply with the instructions for use.****MAPP 15503**A soluble granule formulation containing 200 g/kg metsulfuron-
methyl, a sulfonylurea, for spring weed control in wheat, oats,
barley, triticale and linseed, and for use on green cover on land
temporarily removed from production e.g. set-aside.

® Registered trademark of E. I. du Pont de Nemours and Company

**Contents: 120 grams e****K-41268/31603 - UK**The (COSHH) Control of Substances Hazardous to Health Regulations may
apply to the use of this product at work.

FOR PROFESSIONAL USE ONLY

Manufactured in the EU

Du Pont (U.K.) Limited, Crop Protection Products
4th Floor, Kings Court, London Road, Stevenage, Hertfordshire - SG1 2NG
Tel: 01438 734450 or enquiry.agproducts@dupont.com
Emergency Tel: 0870 820 0418National Poisons information service tel:
111 (England and Wales),
08454 24 24 24 (Scotland)**The Voluntary Initiative**
This label is compliant
with the CPA
Voluntary Initiative GuidanceTo access the Safety Data Sheet
for this product scan the code, visit
www.cropprotection.dupont.co.uk or
alternatively contact your supplier**Batch N°:****CYAN MAGENTA JAUNE BLACK**

SIMBA® SX®**Herbicide**

contains 200 g/kg
metsulfuron-
methyl

WARNING

**Very toxic to aquatic life with
long lasting effects.**

Collect spillage.

Dispose of contents / container to a licensed
hazardous waste-disposal contractor or collection
site except for empty clean containers which may be
disposed of as non-hazardous waste.

Do not contaminate water with the product or its
container (Do not clean application equipment near
surface water/Avoid contamination via drains from
farmyards and roads).

**To avoid risks to human health and the environment,
comply with the instructions for use.**

**MAPP 15503**

A soluble granule formulation containing 200 g/kg metsulfuron-methyl, a sulfonylurea, for spring weed control in wheat, oats, barley, triticale and linseed, and for use on green cover on land temporarily removed from production e.g. set-aside.

® Registered trademark of E. I. du Pont de Nemours and Company



Contents: **120 grams** e

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

FOR PROFESSIONAL USE ONLY

Manufactured in the EU

Du Pont (U.K.) Limited, Crop Protection Products
4th Floor, Kings Court, London Road, Stevenage, Hertfordshire - SG1 2NG
Tel: 01438 734450 or enquiry.agproducts@dupont.com
Emergency Tel : 0870 820 0418

National Poisons information service tel:
111 (England and Wales),
08454 24 24 24 (Scotland)



The Voluntary Initiative
This label is compliant
with the CPA
Voluntary Initiative Guidance



To access the Safety Data Sheet
for this product scan the code, visit
www.cropprotection.dupont.co.uk or
alternatively contact your supplier



K-41268/31603 - UK



IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Crops: Wheat, oats, barley, triticale, linseed and green cover on land temporarily removed from production e.g. set-aside

Maximum individual dose: 30 g product per hectare

Maximum number of treatments: One per crop for wheat, oats, barley, triticale and linseed
One per year for green cover on land temporarily removed from production

Latest time of application

Wheat, oats, barley and triticale: Before the flag leaf sheath extending stage.

Linseed: Before flower buds visible, or up to 30 cm tall, whichever is the sooner.

Green cover on land temporarily removed from production: Before 1 August in year of application

OTHER SPECIFIC RESTRICTIONS:

This product must only be applied from 1 February in the year of harvest until the specified latest time of application.

SIMBA® SX® must only be applied to green cover on land temporarily removed from production where a full green cover is established. Treated green cover on land temporarily removed from production must not be grazed by livestock or harvested for human consumption or used for animal bedding. SIMBA® SX® must only be applied to green cover on land temporarily removed from production where the cover crop is predominantly (i.e. sufficient to maintain reasonable cover) grassland, wheat, barley, oats or triticale.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

DuPont™
Simba® sx®
CEREA HERBICIDE

MAPP 15503

A soluble granule formulation containing 200 g/kg metsulfuron-methyl, a sulfonylurea, for spring weed control in wheat, oats, barley, triticale and linseed, and for use on green cover on land temporarily removed from production e.g. set-aside.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS

Environmental Protection

Use appropriate containment to avoid environmental contamination
Do not contaminate water with this product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application.
Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.



Storage and Disposal

This material and its container must be disposed of safely

Du Pont (U.K.) Limited, Crop Protection Products
4th Floor, Kings Court, London Road, Stevenage, Hertfordshire - SG1 2NG
Tel: 01438 734450 or enquiry.agproducts@dupont.com
Emergency Tél : 0870 820 0418

National Poisons information service tel:
111 (England and Wales), 08454 24 24 24 (Scotland)

® Registered trademark of E. I. du Pont de Nemours and Company

BLACK


DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Restrictions

- SIMBA® SX®, alone or in tank-mix, must not be applied to any crop suffering from stress as a result of drought, waterlogging, low temperatures, pest or disease attack, nutrient or lime deficiency or other factors reducing crop growth.
- Do not use SIMBA® SX® on cereal crops undersown with grasses, clover or other legumes or any other broad-leaved crop.
- SIMBA® SX® must not be tank-mixed with chlorpyrifos. Allow at least 14 days between application of SIMBA® SX® and chlorpyrifos treatments.
- When using tank-mixtures take care not to overlap spray swaths.
- SIMBA® SX® should not be applied within 7 days of rolling any crop.
- Do not use SIMBA® SX® in tank-mix on oats, triticale and linseed. On Linseed allow at least 7 days between SIMBA® SX® and other treatments (10 days if crop is growing poorly or under stress).
- Contract agents should be consulted before using on crops grown for seed.

- Due to the high level of activity of the herbicide, **special care must be taken to avoid damage by drift onto broad-leaved plants outside the target area, or onto ponds, waterways or ditches. Thorough cleansing of equipment is also very important** – see 'Spray-tank cleanout'

• Refer to 'Compatibility' for details on use of SIMBA® SX® with other 'ALS inhibiting' herbicides

Weed Control

SIMBA® SX® is a highly active herbicide with foliar and root activity against a wide range of broad-leaved weeds. It is most effective when applied to small actively growing weeds. As larger weeds often become less susceptible, it is important to note the size of each weed species at the time of application. Good spray cover of the weeds must be obtained. Weed control may be reduced when soil conditions are very dry. Nevertheless, with adequate soil moisture, susceptible weeds germinating soon after treatment will be controlled.

Weed Resistance

When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered resistant to a herbicide if it survives a correctly-applied treatment

at the recommended dose. Development of resistance within a weed species can be avoided or delayed by sequencing or tank-mixing with suitable products having a different mode of action. A strategy for preventing and managing resistance should be adopted. The Weed Resistance Action Groups have produced guidelines and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

Susceptible Weeds

The susceptibility ratings of weeds in the following table refer to good spray cover and good growing conditions.

Weed Species	Plants up to 2 expanded true leaves	Plants up to 6 expanded true leaves	Plants up to 15 cm across/high
Black-bindweed	MS	MS	—
Charlock	S	S	S
Chickweed, Common	S	S	S
Cranes-Bill, Dove's Foot	S	S	—
Dead-nettle, Red	S	S	MS
Docks	S	S	S
Fat-hen	S	R	R
Field-speedwell, Common	S	S	MS

Weed Species	Plants up to 2 expanded true leaves	Plants up to 6 expanded true leaves	Plants up to 15 cm across/high
Forget-me-not, Field	S	MS	MS
Fool's Parsley	S	S	MS
Hemp-nettle, Common	S	S	S
Knotgrass	S	MS	MS
Mayweeds	S	S	S
Nettle, Small	S	S	—
Pale Persicaria	S	S	S
Pansy, Field	S	MS	MS
Parsley-piert	S	S	S
Poppy, Common	S	S	MS
Redshank	S	S	S
Shepherd's-purse	S	S	S
Sowthistle, Smooth	S	—	—
Volunteer Rape	S	S	—
Volunteer Sugar Beet	S	S	S

S = Susceptible

MR = Moderately Resistant

MS = Moderately Susceptible

R = Resistant

Charlock, Common Chickweed, Red Dead-nettle, Mayweeds, Shepherd's-purse and Volunteer Rape will all be controlled by a dose of 25g/ha at the timings listed above.

Soil

SIMBA® SX® can be used on all soil types.

Volume and application

Apply in 100 - 400 litres of water per hectare, using suitable equipment to give good spray cover of the weeds. The higher volume is recommended when crops are thick or weed growth dense. When applying SIMBA® SX®, care should be taken not to overlap spray swaths.

Mixing

Before using SIMBA® SX®, make sure that the spraying equipment is clean. SIMBA® SX® mixes easily with water, but the following mixing procedure should be followed: Quarter fill the spray tank with water, start the agitation and add the required quantity of SIMBA® SX® directly to the tank without prior creaming. Continue agitation while topping up the tank and while spraying.

Compatibility

To widen the spectrum of activity and improve the control of some weeds, SIMBA® SX® may be applied with other herbicides, provided each product can be applied within the label recommendations for its use. Always add SIMBA® SX® to the spray tank first when using in tank-mix.

The following 'joint application' with other sulfonylurea and 'ALS inhibiting' herbicides may be applied to a crop. Requirements and restrictions on other product labels must be adhered to. 'Joint application' is the use of SIMBA® SX® in tank mixture or sequence with one of the products listed below.

Absolute	Eagle*	Lexus Class	Staka SX
Atlantis WG	Excalibur	Lexus SX	Starane Gold
Barton WG	Exceed SX	Lexus Millennium	Starane Vantage
Boxer	Galaxy	Othello	Starane XL
Broadway Star	GF-184	Octavian	Unite
Broadway Sunrise	Hatra	Pacifica	
Bullion	Hiker	Pursuit*	
Dakota	Horus	Slalom	
Ductis SX	Hunter	Spitfire	

* only cereals as following crop

Additional recommendations may be approved since label printing. Please consult your DuPont distributor for latest details on compatibilities.

CROPS

• Cereals

SIMBA® SX® can be used on all winter and spring varieties of wheat, oats, barley and triticale. SIMBA® SX® must only be applied from 1 February, provided the crop has at least two leaves (wheat, oats and triticale) or three leaves (barley), until before the flag leaf extending stage.

Dose

Apply SIMBA® SX® at 30 g/ha when used alone or in tank-mix with other herbicides.

Following Crops

Only cereals, oilseed rape, field beans or grass may be sown in the same calendar year to succeed a cereal crop treated with SIMBA® SX®.

• Linseed

SIMBA® SX® should be applied to actively growing linseed (all varieties) from the growth stage “first pair of true leaves unfolded”, up to 30 cm high or before flower buds visible, whichever comes first. Most annual weeds are best controlled by the earliest application possible.

Dose

Apply SIMBA® SX® at 30 g/ha.

Following Crops

Only cereals should be planted within 16 months of applying SIMBA® SX® to a linseed crop. In the event of crop failure for any reason, sow only wheat within 3 months of the application of SIMBA® SX®.

• Green cover on land temporarily removed from production *e.g.* set-aside

SIMBA® SX® may be applied to areas temporarily removed from production where the green cover is made up predominantly (i.e. sufficient to maintain reasonable cover) of grassland, wheat, barley, oats or triticale. Where established grass species are the main cover crop, some discoloration and stunting may occur. SIMBA® SX® is selective to seedling cereals from the three-leaf stage (GS 13) but should not be used on seedling grasses.

Dose and timing

Apply SIMBA® SX® at 30 g/ha in 100 - 400 litres of water/ha before 1 August in year of application. The higher volume is recommended when weed growth is dense.

Only one application of a sulfonylurea-containing product must be made per calendar year to green cover on land temporarily removed from production.

Following crops

Sow only cereals within 16 months of applying SIMBA® SX® on green cover on land temporarily removed from production e.g. set-aside.

General

SIMBA® SX® effectively controls broad-leaved docks, with best control achieved from application three weeks before or after cutting. Many other broad-leaved plants occurring in the "Set-aside" area will also be controlled or partially controlled, hence it may not be suitable for use where wild flower borders or other forms of conservation headland are to be developed. Before using on land taken out of production as part of a grant-aided scheme, ensure compliance with the management rules of that scheme.

WARNINGS

EXTREME CARE SHOULD BE TAKEN TO AVOID DAMAGE BY DRIFT ONTO BROAD-LEAVED PLANTS OUTSIDE THE TARGET AREA OR ONTO SURFACE WATERS OR DITCHES OR LAND INTENDED FOR CROPPING. SPRAYING EQUIPMENT SHOULD NOT BE DRAINED OR FLUSHED ONTO LAND PLANTED WITH OR INTENDED FOR PLANTING WITH TREES OR CROPS OTHER THAN CEREALS.

POOR CLEANOUT PRACTICES AND INSUFFICIENT WATER VOLUMES USED

FOR THE RINSE PROCEDURE MAY RESULT IN INADEQUATE REMOVAL OF PRODUCT DEPOSITS. SUBSEQUENT USE OF APPLICATION EQUIPMENT IN THESE CIRCUMSTANCES MAY RESULT IN DAMAGE TO NON-CEREAL CROPS.

SPRAY TANK CLEAN-OUT

TO AVOID SUBSEQUENT DAMAGE TO CROPS OTHER THAN CEREALS, IMMEDIATELY AFTER SPRAYING SIMBA® SX® THOROUGHLY CLEAN ALL SPRAY EQUIPMENT, INCLUDING INSIDE AND OUTSIDE OF LID, USING THE FOLLOWING PROCEDURE:

Always start with a clean tank and spray system. Clean spray equipment thoroughly immediately after use.

1. Thoroughly and completely rinse all interior tank surfaces (including lid) with water (use at least 10% of the tank capacity), taking care to remove any visible deposits. Flush pump, filters and boom after removing in-line strainers, nozzle tips and screens (clean these parts separately). Drain the remainder of the rinsate from the tank.
2. Repeat the rinse, flush and drain.

GENERAL NOTES:

Consult label tank cleanup procedures for all tank mix partners and be sure to use the most rigorous procedure recommended.

NOTICE TO BUYER

All goods supplied by us are of a high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

All manufacturers' trademarks are duly acknowledged.

BLACK



BLACK

