



# Stoker™

## Fertiliser

A high load formulation delivering key nutrients  
for oilseed rape, pulses and other crops.

Liquid fertiliser for foliar application		
Nitrogen (N):	2.1% w/w	(32 g/l)
Magnesium (MgO):	3.6% w/w	(53 g/l)
Sulphur (SO <sub>2</sub> ):	9.9% w/w	(148 g/l)
Boron (B):	4.0% w/w	(60 g/l)
Manganese (Mn):	1.5% w/w	(22 g/l)
Molybdenum (Mo):	0.1% w/w	(2 g/l)
Zinc (Zn):	0.1% w/w	(2 g/l)

Net content: 7.3 kg

To be used only where there is a recognised need. Do not exceed the appropriate dose rates.

### FORMULATION

Stoker™ fertiliser is an advanced suspension formulation of manganese, magnesium, boron, zinc and molybdenum with nitrogen and sulphur for foliar application to oilseed rape, brassicas, peas, beans and other arable crops. The formulation provides superior foliar cover and adhesion, optimising availability, uptake and absorption of the application.

### RATES OF USE

Apply 2.5 l/ha in at least 200 litres of water. Ensure thorough cover of the foliage.

### TIMINGS

**Oilseed rape:** Start at the 4 leaf stage and repeat if necessary at 10 - 14 day intervals. Repeat when stems begin to elongate rapidly.

**Leaf brassicas:** Apply early in the season when the plants have sufficient leaf area to intercept the spray. Repeat if necessary at 10 - 14 day intervals until 4 weeks before harvest.

**Pulses (beans, peas, soya beans):** Apply early in the season when the plants have sufficient leaf area to intercept the spray. Repeat if necessary at 10 - 14 day intervals until 4 weeks before harvest.

**Other agricultural and horticultural crops:** Stoker™ fertiliser is suitable for application to other agricultural and horticultural crops where nutrient deficiency has been identified or is suspected.

Applications may begin as soon as the target crop has sufficient foliage to intercept the spray and repeated as required.

It is recommended that soil and tissue analysis results are used whenever possible to guide micro and macronutrient applications to optimise crop nutrient status.

Do not apply this product in very bright sunlight or in temperatures above 25°C. If possible, spray in the evening or early morning for best results.

### APPLICATION

**Shake the container well before opening and use.**

Spray equipment should be clean.

Half fill the spray tank with water, begin agitation and add the required quantity of Stoker™ fertiliser. Rinse the container thoroughly, add the rinsings to the spray-tank, complete the filling and apply without delay. Maintain agitation throughout.

Clean sprayer thoroughly after use.

### COMPATIBILITY

Stoker™ fertiliser is physically and chemically compatible with a wide range of agrochemical products. Consult your supplier for the latest information. Up-to-date information is available at [www.fmc-agro.co.uk](http://www.fmc-agro.co.uk). Always read the label of the partner product carefully before mixing with Stoker™ fertiliser.

### STORAGE

**Protect from frost. Store above 5°C.**

Keep in cool, well ventilated suitable store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs.

### SAFETY PRECAUTIONS

The product is not classified as hazardous under GB CLP and CLP (REG EC 1272/2008). Contains 1,2-Benzisothiazolin-3-one. May produce an allergic reaction.

Batch No: See neck of the bottle

# Contents: 5 litres

Packed in boxes of 2 x 5 litres



FMC Agro Ltd, Rectors Lane, Pentre, Flintshire, CH5 2DH, UK.  
Tel: + 44 (0) 1244 537370 - E-mail: [fmc.agro.uk@fmc.com](mailto:fmc.agro.uk@fmc.com)  
[www.fmc-agro.co.uk](http://www.fmc-agro.co.uk)

FMC, the FMC Logo and Stoker are trademarks of FMC Corporation or an affiliate. ©2018-2022 FMC Corporation. All rights reserved.