

## A high load formulation delivering key micronutrients for cereal crops.

**Trice**<sup>™</sup>

### COMPATIBILITY

Trice is physically and chemically compatible with a wide range of agrochemical products. Consult your supplier for the latest information. Up-to-date information is available at www.fmc-agro.co.uk. Always read the label of the partner product carefully before mixing with Trice.

### **STORAGE**

### Protect from frost. Store above 5°C.

Keep in cool, well ventilated suitable store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs.

### SAFETY PRECAUTIONS

H400: Very toxic to aquatic life.

H411: Toxic to aquatic life with long lasting effects.

### Warning

P273: Avoid release to the environment.

P391: Collect spillage.

P501: Dispose of contents/container as hazardous

waste in accordance with local regulations. Contains 1,2-Benzisothiazolin-3-one. May produce an allergic reaction.

Made in UK Batch No: See neck of the bottle

# Contents: 5 litres

Packed in boxes of 2 x 5 litres

-FMC

FMC Agro Ltd., Rectors Lane, Pentre, Flintshire, CH5 2DH, UK Tel: + 44 (0) 1244 537370 - E-mail: fmc.agro.uk@fmc.com

www.fmc-agro.co.uk

22010850 19012021

### CONTAINS

 Copper (Cu):
 110 g/l

 Manganese (Mn):
 330 g/l

 Zinc (Zn):
 84 g/l

**RATES OF USE** 

Apply 1.0 l/ha in at least 200 litres of water. Repeat if deficiency is severe.

### TIMING

Trice can be applied to crops whenever deficiency is observed or expected, or as a routine maintenance treatment. Applications may commence after the 3 leaf stage in cereals, and in other crops when sufficient foliage has developed to absorb the spray.

Autumn sown crops will benefit from an application before the onset of Winter.

It is recommended that soil and tissue analysis results are used whenever possible to guide micro and macronutrient applications to optimise crop nutrient status.

Do not apply this product in very bright sunlight or in temperatures above 25°C. If possible, spray in the evening or early morning for best results.

### **APPLICATION**

### Shake the container well before opening and use.

Spray equipment should be clean.

Half fill the spray tank with water, begin agitation and add the required quantity of Trice. Rinse the container thoroughly, add the rinsings to the spray-tank, complete the filling and apply without delay. Maintain agitation throughout.

Clean sprayer thoroughly after use.

¥2