

# **Zinc 69**<sup>™</sup>

### Zinc nutritional supplement for agricultural and horticultural

### crops by foliar application

### COMPATIBILITY

Zinc 69 is physically and chemically compatible with a wide range of agrochemical products. Consult your supplier for the latest information. Up-to-date information is available at www.fmc-agro.co.uk. Always read the label of the partner product carefully before mixing with Zinc 69.

### STORAGE

#### Protect from frost. Store above 5°C.

Keep in cool, well ventilated suitable store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs.

### SAFETY PRECAUTIONS

H410: Very toxic to aquatic life with long lasting effects.

Warning

P273: Avoid release to the environment.

P391: Collect spillage.

P501: Dispose of contents/container as hazardous waste in accordance with local regulations.

Contains 1,2-Benzisothiazolin-3-one. May produce an allergic reaction.

Verdi-Crop

Made in UK Batch No: See neck of the bottle

## Contents: 5 litres

Packed in boxes of 2 x 5 litres

### -FMC

FMC Agro Ltd., Rectors Lane, Pentre, Flintshire, CH5 2DH, UK. Tel: + 44 (0) 1244 537370 - E-mail: fmc.agro.uk@fmc.com

www.fmc-agro.co.uk

15096959 23032020

### CONTAINS

Zinc (Zn): 700 g/l

### FORMULATION

Zinc 69 is fully formulated for ease of application and maximum availability to the growing crop.

### RATES OF USE

Apply 1 l/ha in at least 200 litres of water.

Repeat applications may be required where moderate/severe deficiency is known or identified by analysis. Sub-clinical (maintenance) dressing – apply 0.5 l/ha twice during the Spring/Summer period.

### TIMING

**Cereals:** 3 leaf stage to second node detectable (GS 13 - 32). For Autumn-sown crops, an application prior to Winter dormancy is strongly recommended.

Maize: 3 - 8 leaf stage.

**Oilseed rape:** October/November before Winter dormancy. Repeat in the Spring at the beginning of stem extension.

Sugar beet and root vegetables: Apply from the 4 - 6 leaf stage Potatoes: From 1 week after 100 % emergence.

Peas and beans: From the 4 - 6 true leaf stage.

Top fruit: Apply post petal-fall.

Other crops susceptible to zinc deficiency: Apply as early in the growing season as possible providing there is adequate leaf cover.

It is recommended that soil and tissue analysis results are used whenever possible to guide micro and macronutrient applications to optimise crop nutrient status.

Do not apply this product in very bright sunlight or in temperatures above 25°C. If possible, spray in the evening or early morning for best results.

### APPLICATION

### Shake the container well before opening and use.

Spray equipment should be clean.

Half fill the spray tank with water, begin agitation and add the required quantity of Zinc 69. Rinse the container thoroughly, add the rinsings to the spray tank, complete the filling and apply without delay. Maintain agitation throughout.

Clean sprayer thoroughly after use.