



TIZCA® Potato Fungicide

GROUP 29 FUNGICIDE

MAPP No. 18813

A suspension concentrate containing 500 g/litre fluazinam (ISO) and 1,2-benzisothiazol-3(2H)-one

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Authorisation holder: FMC Agro Limited, Rectors Lane, Pentre, Flintshire, CH5 2DH. Tel: 01244 537370. Email: fmc.agro.uk@fmc.com. Website: www.fmc-agro.co.uk. Technical Helpline: 01423 205011

Emergency No.s - Medical Incidents Involving Crop Protection Products: England & Wales: 111, Scotland: 08454 24 24 24, Northern Ireland: Local GP or Pharmacist, Republic of Ireland: 01 837 9964.

Healthcare professionals seeking poisons information: www.toxbase.org.

Emergency No.s - Transportation, Warehousing & After Sale Incidents: CHEMTRAC help centre Dialling from the UK and NI: 0870 820 0418, Dialling from Ireland: 01 901 4670 Batch Number: see elsewhere on the container

Contents: **5 litres**

RISK AND SAFETY INFORMATION

TIZCA® FUNGICIDE

Contains fluazinam (ISO) and 1,2-benzisothiazol-3(2H)-one

WARNING

May cause an allergic skin reaction. Suspected of damaging the unborn child.

Very toxic to aquatic life with long lasting effects.

Obtain special instructions before use.

Avoid breathing mist or vapours. Wear protective gloves, protective clothing and eye/face protection.

IF exposed or concerned: Get medical attention/advice.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



The
Voluntary
Initiative

This product label is compliant with the CropLife UK Voluntary Initiative guidance.

IMPORTANT INFORMATION FOR USE ONLY AS A PROFESSIONAL FUNGICIDE.

Crops	Maximum individual dose (mL product/ha)	Maximum total dose (L product/ha/crop)	Latest time of application	Aquatic buffer zone distance
Potato	400	3	A minimum of seven days must be observed between applications at the 300 mL product/ha rate and a minimum of ten days at the 400 mL product/ha rate	7 metres (see Environmental Protection Phrase 1)

READ THE LABEL BEFORE USE. USING THE PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.
FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

11006156_22067571_GREAT BRITAIN AND NORTHERN IRELAND

To access the Safety Data Sheet for this product scan the QR code or use the web link:
https://ag.fmc.com/uk/sites/default/files/2025/11/Tizca_SDs_GB.pdf



Alternatively, contact your supplier

P 032C P 2149C CYAN MAGENTA YELLOW BLACK



	FILL IN	DIMENSIONS (WXH)
Country	GREAT BRITAIN AND NORTHERN IRELAND	Single label
Product SKU	11006156	Base for Booklet
Product SKU Description	TIZCA_5L_GB	Booklet
Label SKU	22067571	Number of pages (Booklet) 3 pages
Label SKU Description	LBL_ML-B_5L_TIZCA_GB_2	Carton label
Label SKU (earlier version(s))		Preprinted Material

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section should be read carefully in order to obtain safe and successful use of this product.

Disease control

TIZCA® is a fungicide for the control of foliar blight (*Phytophthora infestans*) and protection against tuber blight in potatoes. TIZCA® is a protectant fungicide and therefore applications must begin prior to the development of blight.

Crops

TIZCA® can be used in early and maincrop potatoes.

Rate and Time of Application

Apply TIZCA® at 300-400 ml/ha

Initial application: Irrespective of crop growth stage, apply TIZCA® at the first blight warning or when local weather conditions are favourable for the disease, whichever is the sooner. In the absence of weather conducive to disease development, the first application should be made just before the crop meets in the rows.

Repeat applications:

Risk of disease development:	Interval between applications:	Dose rate:
Low risk (before conditions conducive to blight development have occurred)	14 days	300-400 ml/ha
High risk (following two warm, humid days or in irrigated crops)	7-10 days	300-400 ml/ha
Severe risk (during continued warm, humid weather and when blight is in nearby crops, potato dumps or volunteer plants)	7 days or 10 days	300 ml/ha or 400 ml/ha Apply according to blight warning and disease prediction systems

Tuber blight: To reduce the risk of tuber blight at harvest, TIZCA® should be used regularly at the recommended intervals from full canopy development to complete haulm desiccation. A complete haulm desiccation programme must be carried out to reduce the risk of development of subsequent tuber blight. The crop should not be lifted for at least 10 days after complete kill of the haulm, and stored crops should not be lifted whilst any green tissue remains.

Potato blight fungicides may not give complete protection when the disease risk is severe. However, disease development will be delayed significantly with a well timed spray programme.

Mixing and Spraying

Apply TIZCA® in 200-500 litres of water per hectare. Use adequate water to ensure complete coverage of the foliage and stems, increasing the volume as haulm growth progresses, particularly in dense crops and under high or severe blight risk conditions. Apply as a MEDIUM spray, as defined by BCPC. Half fill the spray tank with clean water and start the agitation. Shake the container well before opening and pour in the required amount of product. Rinse empty containers thoroughly and add rinsings to the spray tank. Add the remainder of the water and continue agitation until spraying is completed. When tank mixes are to be used, each product should be added separately to the spray tank, taking due note of any instructions given as to the order of mixing. Do not leave the spray liquid in the sprayer for long periods.

Advice for operators

- TIZCA® may cause transient skin reaction in a small number of sensitive individuals.
- People who have been sensitised by TIZCA® should not use or have further contact with the product.
- Avoid contact with spray drift. Ensure spraying is carried out under good conditions and that the sprayer is adjusted to minimise drift.
- Do not allow unprotected skin to come into contact with contaminated clothing.
- Do not allow unprotected skin to come into contact with contaminated sprayer parts.
- Ensure sprayer is washed down daily.
- Wash all protective clothing regularly, preferably daily, when working with TIZCA®.
- When entering crops do not allow unprotected skin to come into contact with treated foliage.

	FILL IN		DIMENSIONS (WXH)
Country	GREAT BRITAIN AND NORTHERN IRELAND	Single label	
Product SKU	11006156	Base for Booklet	
Product SKU Description	TIZCA_5L_GB	Booklet	129/124 x 149 mm
Label SKU	22067571	Number of pages (Booklet)	3 pages
Label SKU Description	LBL_ML-B_5L_TIZCA_GB_2	Carton label	
Label SKU (earlier version(s))		Preprinted Material	

Compatibility

For information on tank mixing please contact your FMC distributor or refer to the FMC web-site (www.fmc-agro.co.uk).

Resistance

TIZCA® contains fluazinam, a protectant multisite fungicide (FRAC Group 29).

Disease control may be reduced if strains of pathogen less sensitive to TIZCA® develop.

Make a timely start to spray programmes, when there is warning or risk, or, in the absence of weather conducive to disease development, just before the crop foliage begins to meet in the rows. Use forecast schemes and / or local knowledge to time application more accurately (see details under 'Rate and Time of Application').

Alternate with other fungicides having a different mode of action to FRAC Group 29 fungicides.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

SAFETY PRECAUTIONS**Operator Protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

Environmental Protection

Crops with >5m buffer zone:

Since there is a risk to aquatic life from use, direct spray from horizontal boom sprayers must not be allowed to fall within the distance specified for the crop of the top of the bank of any static or flowing waterbody or within 1m of the top of a ditch which is dry at the time of application. Spray must be aimed away from water.

Buffer zones must be measured in accordance with the guidance set out in the booklet 'Local Environment Risk Assessment for Pesticides - Horizontal Boom Sprayers' available from HSE Chemicals Regulation Directorate's website and any amendments made to it.

The results of the LERAP must be recorded in written form and must be available for a period of three years for inspection to any person entitled to exercise enforcement powers under or in connection with the Plant Protection Products Regulations 2011 or the Plant Protection Products (Sustainable Use) Regulations 2012.

(An electronic record will satisfy the requirement for a written record, providing it is similarly available for inspection and can be copied).

Where the buffer zone is greater than 5m then it must not be reduced under the LERAP scheme.

Other specific restrictions

This product must not be applied via hand-held equipment.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with the 'Local Environment Risk Assessment for Pesticides Horizontal Boom Sprayers' booklet available from the HSE Chemicals Regulation Directorate's website or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years. A minimum of 7 days must be observed between applications of 300 ml product/ha and a minimum of 10 days between applications at 400 ml product/ha.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY, by using an integrate pressure rinsing device or manually rinsing three times. Add washings to sprayer at the time of filling and dispose of safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

NOTICE TO BUYER

FMC as the manufacturer warrants that the composition of the product in sealed original FMC packaging complies with the information given on this label. FMC is not liable for any direct or indirect consequences resulting from improper storage, handling or use of the product contrary to or otherwise not in compliance with this label, law and regulations. Various factors, especially local conditions, such as soil condition, plant varieties and weather conditions, may either cause the product not to have the full desired effect or result in damage to the crops treated. FMC does not assume any liability for such results.

All manufacturers trademarks are duly acknowledged. FMC, the FMC logo, and TIZCA® are trademarks of FMC Corporation and/or an affiliate.

©2014-2025 FMC Corporation. All rights reserved.

	FILL IN		DIMENSIONS (WXH)
Country	GREAT BRITAIN AND NORTHERN IRELAND	Single label	
Product SKU	11006156	Base for Booklet	
Product SKU Description	TIZCA_5L_GB	Booklet	129/124 x 149 mm
Label SKU	22067571	Number of pages (Booklet)	3 pages
Label SKU Description	LBL_ML-B_5L_TIZCA_GB_2	Carton label	
Label SKU (earlier version(s))		Preprinted Material	