

OPEN HERE

UFI: HUYW-N2XQ-0N4N-3E4J

Risk and Safety Information**BIPLAY® SX®**

Herbicide

Contains tribenuron-methyl.
May produce an allergic
reaction

**May cause damage through
prolonged or repeated exposure.
Very toxic to aquatic life with long
lasting effects.**

Do not breathe dust or spray.
Avoid release to the environment.
Get medical attention if you feel
unwell.

Collect spillage.
Dispose of contents/
container to a licensed
hazardous waste disposal
contractor or collection site except
for triple rinsed empty containers
which can be
disposed of as non-hazardous
waste.

Do not contaminate water with the product or its
container (Do not clean application equipment near
surface water/Avoid contamination via drains from
farmyards and roads).

**To avoid risks to human health and the environment,
comply with the instructions for use PCS No: 03434**

**WARNING****FMC**

BIPLAY® SX®

Cereal Herbicide

GROUP 2 HERBICIDE

PCS No: 03434

A water soluble granule formulation
containing 111 g/kg metsulfuron methyl and
222 g/kg tribenuron methyl for spring weed
control in wheat, barley, oats and triticale.

FMC Agro Limited, Rectors Lane, Pentre, Flintshire,
CH5 2DH. Tel: 01244 537370. Email: fmc.agro.uk@fmc.com
Website: www.fmc-agro.co.uk
Technical Helpline: 01423 205011

* is a trademark of FMC Corporation or an affiliate

Emergency No.s - Medical Incidents Involving

Crop Protection Products: England & Wales: 111,
Scotland: 08454 24 24 24, Northern Ireland: Local GP
or Pharmacist, Republic of Ireland: 01 837 9964.

Healthcare professionals seeking poisons information :
www.toxbase.org.

Emergency No.s - Transportation, Warehousing &

After Sale Incidents: CHEMTREC help centre Dialling
from the UK and NI: 0870 820 0418, Dialling from
Ireland: 01 901 4670

FOR PROFESSIONAL USE ONLY**SAFETY PRECAUTIONS**

Operator protection: When using do not eat, drink
or smoke.

Environmental Protection: DO NOT CONTAMINATE
WATER WITH THE PRODUCT OR ITS CONTAINER
(DO NOT CLEAN APPLICATION EQUIPMENT NEAR
SURFACE WATER/AVOID CONTAMINATION VIA
DRAINS FROM FARMYARDS AND ROADS).

Do not contaminate ponds, waterways or ditches with
chemical or used container.

To protect aquatic organisms respect an unsprayed
buffer zone of 5 m to surface water bodies. Direct spray
away from water.

Storage and Disposal: Do not re-use container for any
purpose and dispose of safely.

Batch N°:

Contents: **135 g**

11005808_22072106_IRELAND

| | FILL IN | | DIMENSION (WxH) |
|-----------------------------|------------------------------|---------------------------|------------------------------|
| Country | IRELAND - IE | Single label | |
| Product SKU | 11005808 | Base for Booklet | |
| Product SKU Description | BIPLAY SX_135G_IE | Booklet | BKL 230x95 mm (190x95 mm) |
| Label SKU | 22072106 | Number of pages (Booklet) | 6 pages |
| Label SKU Description | LBL_ML-F_135G_BIPLAY SX_IE_3 | Carton label | |
| Label SKU (earlier version) | 22045342, 22046664 | Preprinted Material | No |

BIPLAY® SX®

Cereal Herbicide

GROUP **2** HERBICIDE

PCS No: 03434

A water soluble granule formulation containing 111 g/kg metsulfuron methyl and 222 g/kg tribenuron methyl for spring weed control in wheat, barley, oats and triticale.

FMC Agro Limited, Rectors Lane, Pentre, Flintshire, CH5 2DH. Tel: 01244 537370.
Email: fmc.agro.uk@fmc.com. Website: www.fmc-agro.co.uk.
Technical Helpline: 01423 205011

Emergency No.s - Medical Incidents Involving Crop Protection Products:

England & Wales: 111, Scotland: 08454 24 24 24, Northern Ireland: Local GP or Pharmacist, Republic of Ireland: 01 837 9964.

Healthcare professionals seeking poisons information : www.toxbase.org.

Emergency No.s - Transportation, Warehousing & After Sale Incidents:

CHEMTREC help centre Dialling from the UK and NI: 0870 820 0418, Dialling from Ireland: 01 901 4670

® is a trademark of FMC Corporation or an affiliate

| IMPORTANT INFORMATION FOR USE ONLY AS A PROFESSIONAL HERBICIDE | | | | |
|---|--|---|---|---|
| Crops | Maximum individual and total dose (g product / ha) | Maximum number of treatments (per crop) | Latest time of application | Method of application |
| Wheat, barley, oats and triticale | 45 | 1 | Up to and including flag-leaf ligule / collar just visible (GS39) | Tractor mounted/ trailed horizontal sprayer |
| READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS | | | | |

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

- BIPLAY® SX® should not be applied to any crop suffering from stress as a result of drought, water-logging, low temperatures, pest or disease attack, nutrient or lime deficiency or other factors reducing crop growth.
- Do not use BIPLAY® SX® on cereal crops undersown with grasses, clover or other legumes or any other broad-leaved crop.
- Due to the high level of activity of the herbicide, special care must be taken to avoid damage by drift onto broad-leaved plants outside the target area, or onto ponds, waterways or ditches. Thorough cleansing of equipment is also very important - see below.
- Contract agents should be consulted before using on crops grown for seed.
- BIPLAY® SX® should not be applied within 7 days of rolling the crop.
- Do not apply BIPLAY® SX® more than once to any cereal crop.
- All requirements or restrictions on other product labels must be adhered to when applied to the same crop as BIPLAY® SX®. Contact your FMC distributor for further information.
- Do not use BIPLAY® SX® in tank-mix on oats or triticale. Allow at least 7 days between BIPLAY® SX® and other treatments (10 days if crop is growing poorly or under stress).
- Refer to 'Compatibility' for details on use of BIPLAY® SX® with other 'ALS inhibiting' herbicides.

WEED CONTROL

BIPLAY® SX® contains metsulfuron methyl and tribenuron methyl, sulfonylurea (ALS inhibitor) herbicides which work mainly by contact action and is most effective if applied when the weeds are small and actively growing. Good spray cover of the weeds must be obtained for best results. Residual effects may be reduced by heavy rain. Susceptible plants cease growth almost immediately after application and symptoms can be seen about two weeks after application. Weed control may be reduced when soil conditions are very dry. The susceptibility rating of weeds in the following table refer to good spray cover and good growing conditions.

| Weed Species | Plants up to 2 fully expanded true leaves | Plants up to 6 fully expanded true leaves |
|-------------------------|---|---|
| Chickweed, Common | S | S |
| Dead-nettle, Red | S | S |
| Field Pansy, Common | S | MS |
| Field-speedwell, Common | S | S |
| Forget-me-not, Field | S | MS |
| Fool's Parsley | MS | MS |
| Hemp-nettle, Common | MS | MS |
| Knotgrass | S | MS |
| Mayweeds | S | S |
| Poppy, Common | S | S |

S = Susceptible; MS = Moderately susceptible

RESISTANCE

This product contains tribenuron-methyl and metsulfuron-methyl, which are ALS inhibitors, also classified by the Herbicide Resistance Action Committee as 'Group 2'. When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered resistant to a herbicide if it survives a correctly applied treatment at the recommended dose. Development of resistance with a weed species can be avoided or delayed by alternating (or tank mixing) with suitable products having a different mode of action which have activity against the target weeds.

The use of cultural methods of control and herbicides with non-ALS modes of action within the cropping season and/or throughout the cropping rotation will help reduce the risk of developing resistance.

Do not apply to weeds where target site resistance to ALS inhibiting herbicides is confirmed.

To reduce the risk of developing resistance applications should be made to young, actively growing weeds.

For the control of Chickweed, this product must always be applied in a mixture with an alternate mode of action product at recommended rates.

A strategy for preventing and managing resistance should be adopted. Further details and advice on how to implement such a strategy may be obtained from your crop adviser or supplier. The Herbicide Resistance Action Committee (HRAC) also produces guidelines that may be consulted for additional information.

CROPS

BIPLAY® SX® can be used on all varieties of winter wheat, spring wheat, winter barley, spring barley, winter oats, spring oats and triticale between the growth stages given below.

DOSE

Apply BIPLAY® SX® at 45 g product/ha.

TIMING

BIPLAY® SX® should be applied in the spring after 1st February from the three-leaf stage (GS13) up to and including the flag-leaf ligule / collar just visible (GS39) stage of crop growth.

FOLLOWING CROPS

Only cereals, oilseed rape, field beans or grass may be sown in the same calendar year as harvest of a cereal crop treated with BIPLAY® SX®.

VOLUME AND APPLICATION

BEFORE USING BIPLAY® SX®, SPRAYING EQUIPMENT MUST BE CLEAN AND FREE FROM CONTAMINATION WITH OTHER PESTICIDES.

Application should be made in 100-200 litres of water per hectare using suitable equipment to give good spray cover of the weeds. When crops are thick or weed growth dense, use 400 litres of water per hectare. Care should be taken not to overlap spray swaths.

MIXING

Quarter fill the spray tank with water, start the agitation and add the required quantity of BIPLAY® SX® directly to the tank without prior creaming. Continue agitation while topping up the tank and while spraying.

COMPATIBILITY

In any tank-mix add BIPLAY® SX® to the tank first and ensure it is fully dispersed before adding the partner product. Do not allow BIPLAY® SX® to come into contact with undiluted pesticide concentrate. Products should only be tank-mixed if each product can be applied within the label recommendations for its use.

For further information contact your FMC distributor.

SPRAY TANK CLEAN-OUT

POOR CLEANOUT PRACTICES AND INSUFFICIENT WATER VOLUMES USED FOR THE RINSE PROCEDURE MAY RESULT IN INADEQUATE REMOVAL OF PRODUCT DEPOSITS. SUBSEQUENT USE OF APPLICATION EQUIPMENT IN THESE CIRCUMSTANCES MAY RESULT IN DAMAGE TO NON-CEREAL CROPS.

TO AVOID SUBSEQUENT DAMAGE TO CROPS OTHER THAN CEREALS, IMMEDIATELY AFTER SPRAYING BIPLAY® SX® THOROUGHLY CLEAN ALL SPRAY EQUIPMENT, INCLUDING INSIDE AND OUTSIDE OF LID, USING THE FOLLOWING PROCEDURE:

Always start with a clean tank and spray system. Clean spray equipment thoroughly immediately after use.

1. Thoroughly and completely rinse all interior tank surfaces (including lid) with water (use at least 10% of the tank capacity), taking care to remove any visible deposits. Flush pump, filters and boom after removing in-line strainers, nozzle tips and screens (clean these parts separately). Drain the remainder of the rinsate from the tank.

2. Repeat the rinse, flush and drain.
3. Dispose of washings safely. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

Note:

Consult label tank cleanup procedures for all tank mix partners and be sure to use the most rigorous procedure recommended.

WARNING

EXTREME CARE SHOULD BE TAKEN TO AVOID DAMAGE BY DRIFT OF SPRAY ONTO BROAD-LEAVED PLANTS OUTSIDE THE TARGET AREA OR ONTO PONDS, WATERWAYS OR DITCHES. SPRAYING EQUIPMENT SHOULD NOT BE DRAINED OR FLUSHED ON LAND PLANTED WITH OR INTENDED FOR PLANTING WITH TREES OR CROPS OTHER THAN CEREALS.

NOTICE TO BUYER

FMC as the manufacturer warrants that the composition of the product in sealed original FMC packaging complies with the information given on this label. FMC is not liable for any direct or indirect consequences resulting from improper storage, handling or use of the product contrary to or otherwise not in compliance with this label, law and regulations. Various factors, especially local conditions, such as soil condition, plant varieties and weather conditions, may either cause the product not to have the full desired effect or result in damage to the crops treated. FMC does not assume any liability for such results.

FMC, the FMC logo, BIPLAY® SX® are trademarks of FMC Corporation and/or an affiliate. ©2016-2026 FMC Corporation. All rights reserved.

