


INKA® SX®
Herbicide

Contains Tribenuron-methyl.
May produce an allergic
reaction

**May cause damage to
organs through prolonged
or repeated exposure.
Very toxic to aquatic life with long
lasting effects.**

Do not breathe dust or spray.
Get medical advice/attention
if you feel unwell.

Collect spillage.

Dispose of contents /
container to a licensed
hazardous-waste disposal
contractor or collection
site except for empty clean
containers which can be disposed of as non-
hazardous waste.

Do not contaminate water with the product or its
container (Do not clean application
equipment near surface water/Avoid
contamination via drains from farmyards and
roads).

**To avoid risks to human health and the
environment, comply with the instructions for
use**


WARNING
FMC
INKA® SX®
Cereal Herbicide

GROUP

2

HERBICIDE

MAPP 21882


A soluble granule formulation containing 250 g/kg
tribenuron methyl and 250 g/kg thifensulfuron
methyl, both sulfonylureas, for weed control in
cereals.

The (COSHH) Control of Substances Hazardous
to Health Regulations may apply to the use of this
product at work.

* is a trademark of FMC Corporation or an affiliate


 Contents: **240 g**

FMC Agro Limited, Rectors Lane, Pentre, Flintshire,
CH5 2DH. Tel: 01244 537370. Email: fmc.agro.uk@
fmc.com. Website: www.fmc-agro.co.uk. Technical
Helpline: 01423 205011

**Emergency No.s - Medical Incidents Involving
Crop Protection Products:** England & Wales:
111, Scotland: 08454 24 24 24, Northern Ireland:
Local GP or Pharmacist, Republic of Ireland:
01 837 9964. Healthcare professionals seeking
poisons information : www.toxbase.org.

**Emergency No.s - Transportation, Warehousing
& After Sale Incidents:** CHEMTREC help centre
Dialling from the UK and NI: 0870 820 0418, Dialling
from Ireland: 01 901 4670

**READ DIRECTIONS FOR USE ON ATTACHED
LEAFLET**

FOR PROFESSIONAL USE ONLY


The Voluntary Initiative
This label is compliant
with the CropLife UK
Voluntary Initiative
Guidance



To access the Safety
Data Sheet for this
product scan the code

Batch N°:

11005586_22066733_LBL_ML-F_240G_INKA SX_GB_2_BOOKLET,8pages.indd 1

16/09/2026 13:07

P. 2408 P. 032 CYAN MAGENTA JAUNE NOIR

	FILL IN		DIMENSION (WxH)
Country	GREAT BRITAIN AND NORTHERN IRELAND	Single label	
Product SKU	11005586	Base for Booklet	
Product SKU Description	INKA SX_240G_GB	Booklet	BKL 230x95 mm (190x95 mm)
Label SKU	22066733	Number of pages (Booklet)	8 pages
Label SKU Description	LBL_ML-F_240G_INKA SX_GB_2	Carton label	
Label SKU (earlier version)	22051561	Preprinted Material	No

11005586_22066733_GREAT BRITAIN AND NORTHERN IRELAND

INKA® SX®

Cereal Herbicide

GROUP

2

HERBICIDE

MAPP 21882

A soluble granule formulation containing 250 g/kg tribenuron methyl and 250 g/kg thifensulfuron methyl, both sulfonylureas, for weed control in cereals.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

FMC Agro Limited, Rectors Lane, Pentre, Flintshire, CH5 2DH. Tel: 01244 537370. Email: fmc.agro.uk@fmc.com. Website: www.fmc-agro.co.uk. Technical Helpline: 01423 205011

Emergency No.s - Medical Incidents Involving Crop Protection Products:

England & Wales: 111, Scotland: 08454 24 24 24, Northern Ireland: Local GP or Pharmacist, Republic of Ireland: 01 837 9964. Healthcare professionals seeking poisons information : www.toxbase.org.

Emergency No.s - Transportation, Warehousing & After Sale Incidents:

CHEMTREC help centre Dialling from the UK and NI: 0870 820 0418, Dialling from Ireland: 01 901 4670

® is a trademark of FMC Corporation or an affiliate

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Restrictions

- Do not apply to any crop suffering from stress as a result of drought, water-logging, low temperatures, pest or disease attack, nutrient or lime deficiency or other factors reducing crop growth.
- Do not apply to crops undersown with grass, clover or other legumes or any other broad-leaved crop.
- Do not apply more than once to any cereal crop.
- Do not apply within seven days of rolling the crop.
- Contract agents should be consulted before using on crops grown for seed.
- Due to the high level of activity of the herbicide, special care must be taken to avoid damage by drift onto broad-leaved plants outside the target area, or onto ponds, waterways or ditches. Thorough cleansing of equipment is also very important - see below.
- Refer to 'Compatibility' for details on use of INKA® SX® with other 'ALS inhibiting' herbicides.

Weed Control

INKA® SX® contains thifensulfuron-methyl and tribenuron-methyl, sulfonylurea (ALS inhibitor) herbicides that work predominantly by foliar action with some root activity. It controls a wide range of broad-leaved weeds and is most effective if applied when the weeds are small and actively growing. Good spray cover of weeds must

be obtained and since larger weeds often become less susceptible, it is important to note the size of each weed at the time of application. Susceptible plants cease growth almost immediately after post-emergence treatment and symptoms can be seen about two weeks after spring application. It is also important to identify the weeds occurring in the crop and refer to the weed table to ensure that the weeds present are those susceptible to INKA® SX®. The susceptibility ratings of weeds in the following table refer to good spray cover and good growing conditions.

Weed Resistance

This product contains tribenuron-methyl and thifensulfuron-methyl which is an 'ALS inhibitor', also classified by the Herbicide Resistance Action Committee as 'Group 2 (previously Group B')'.

When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered resistant to a herbicide if it survives a correctly-applied treatment at the recommended dose. Development of resistance with a weed species can be avoided or delayed by using cultural methods of control; using herbicides with non-ALS modes of action within the cropping season and/or throughout the cropping rotation and sequencing or tank-mixing with suitable products having a different mode of action where two or more components are active against the target weeds.

A strategy for preventing and managing resistance should be adopted. Always monitor effectiveness of weed control and investigate any patches of poor control. If unexplained, contact your agronomist, who may consider a resistance test appropriate. The use of cultural methods of control and herbicides with non-ALS modes of action within the cropping season and/or throughout the cropping rotation will help reduce the risk of developing resistance. To reduce the risk of developing resistance applications should be made to young, actively growing weeds. Avoid using herbicides with a single mode of action, such as ALS herbicides, in the same fields over a number of years. The Weed Resistance Action Group has produced guidelines and copies that are available from the HCP, CropLife UK, your distributor, crop advisor or product manufacturer.

Susceptible Weeds

The susceptibility ratings of weeds in the following table refer to good spray cover and good growing conditions.

Weed Species	Plants up to 2 expanded true leaves	Plants up to 6 expanded true leaves	Plants up to 15 cm across/high
Black-bindweed	S	S	MS
Charlock	S	S	S
Chickweed, Common	S	S	S
Dead -nettle, Red	S	S	MR
Dock, Broad-leaved	S	S	S
Fat-hen	S	S	MS
Forget-me-not	S	S	MS
Fumitory	S	S	R
Hemp-nettle, Common	S	S	-
Mayweed#	S	S	S
Poppy, Common	S	S	R
Redshank	S	S	MS
Shepherd's-purse	S	S	MS

S = Susceptible MS = Moderately Susceptible MR = Moderately Resistant R = Resistant

Mayweeds can be controlled up to 20 cm

In addition to the above weeds, volunteer beans and oilseed rape will be controlled by a dose of 20 g/ha.

Soil and weather

INKA® SX® can be used on all soil types. Weed control may be reduced when conditions are very dry.

Volume and application

BEFORE USING INKA® SX®, SPRAYING EQUIPMENT MUST BE CLEAN AND FREE FROM CONTAMINATION WITH OTHER PESTICIDES.

Application should be made in 200 litres of water per hectare using suitable ground equipment to give good spray cover of the weeds. Use a conventional field crop sprayer at a pressure of 2 - 3 bars and apply as a MEDIUM spray (as defined by BCPC). Care should be taken not to overlap spray swaths.

Mixing

Quarter fill the spray tank with clean water, start the agitation and add the required quantity of INKA® SX® directly to the tank without prior creaming. Continue agitation while topping up the tank and while spraying.

Compatibility

To widen the spectrum of activity and improve the control of some weeds, INKA® SX® may be applied with other herbicides, provided each product can be applied within the label recommendations for its use. Always add INKA® SX® to the spray tank first when using in tank-mix.

The following 'joint application' with other sulfonylurea and 'ALS inhibiting' herbicides may be applied to a crop. Requirements and restrictions on other product labels must be adhered to. 'Joint application' is the use of INKA® SX® in tank mixture or sequence with one of the products listed below.

Absolute	Eagle*	Hunter	Spitfire
Atlantis WG	Excalibur	Lexus Class	Spitfire Solo
Barton WG	Exceed SX	Lexus SX	Staka SX
Broadway Star	Galaxy	Oklar SX	Starane Gold
Broadway Sunrise	GF-184	Oriel 50 SX	Starane Vantage
Boxer	GF-2070\$	Othello	Starane XL
Bullion	Hatra	Pacifica	Unite\$
Dakota	Hiker	Pursuit*	
Ductis SX	Horus	Slalom	

* only cereals as following crop

\$ use only in sequence with INKA® SX®

Additional recommendations may be approved since label printing. Please consult your FMC distributor for latest details on compatibilities.

Crops

INKA® SX® can be used on all varieties of winter and spring wheat and barley, oats, rye and triticale.

Timing

INKA® SX® should be applied from the three-leaf stage up to and including the flag leaf ligule/collar just visible. Safety to crops of oats, rye and triticale treated before four tillers (GS24) has not been assessed.

Dose

Apply INKA® SX® at 60 g/ha.

Following Crops

Only cereals, field beans or oilseed rape may be sown in the same calendar year as harvest of a cereal crop treated with INKA® SX®. In the event of a crop failure for any reason, sow only a cereal crop within three months of application of INKA® SX®. Before sowing, soil should be ploughed and cultivated to a depth of at least 15cm. After three months field beans and oilseed rape may also be sown.

WARNINGS

EXTREME CARE SHOULD BE TAKEN TO AVOID DAMAGE BY DRIFT ONTO BROAD-LEAVED PLANTS OUTSIDE THE TARGET AREA OR ONTO SURFACE WATERS OR DITCHES OR LAND INTENDED FOR CROPPING. SPRAYING EQUIPMENT SHOULD NOT BE DRAINED OR FLUSHED ONTO LAND PLANTED WITH OR INTENDED FOR PLANTING WITH TREES OR CROPS OTHER THAN CEREALS.

POOR CLEANOUT PRACTICES AND INSUFFICIENT WATER VOLUMES USED FOR THE RINSE PROCEDURE MAY RESULT IN INADEQUATE REMOVAL OF PRODUCT DEPOSITS. SUBSEQUENT USE OF APPLICATION EQUIPMENT IN THESE CIRCUMSTANCES MAY RESULT IN DAMAGE TO NON-CEREAL CROPS.

SPRAY TANK CLEAN-OUT

TO AVOID SUBSEQUENT DAMAGE TO CROPS OTHER THAN CEREALS, IMMEDIATELY AFTER SPRAYING INKA® SX® THOROUGHLY CLEAN ALL SPRAY EQUIPMENT, INCLUDING INSIDE AND OUTSIDE OF LID, USING THE FOLLOWING PROCEDURE:

Always start with a clean tank and spray system. Clean spray equipment thoroughly immediately after use.

1. Thoroughly and completely rinse all interior tank surfaces (including lid) with water (use at least 10% of the tank capacity), taking care to remove any visible deposits. Flush pump, filters and boom after removing in-line strainers, nozzle tips and screens (clean these parts separately). Drain the remainder of the rinsate from the tank.
2. Repeat the rinse, flush and drain.

GENERAL NOTES:

- Consult label tank cleanup procedures for all tank mix partners and be sure to use the most rigorous procedure recommended.

COMPANY ADVISORY INFORMATION

This section is not part of the Product label under Regulation 1107/2009 5 and provides additional advice on product use at the discretion of FMC Agro Limited.

NOTICE TO BUYER

FMC as the manufacturer warrants that the composition of the product in sealed original FMC packaging complies with the information given on this label. FMC is not liable for any direct or indirect consequences resulting from improper storage, handling or use of the product contrary to or otherwise not in compliance with this label, law and regulations.

Various factors, especially local conditions, such as soil condition, plant varieties and weather conditions, may either cause the product not to have the full desired effect or result in damage to the crops treated. FMC does not assume any liability for such results.

FMC, the FMC logo, INKA® and SX® are trademarks of FMC Corporation and/or an affiliate.

SAFETY PRECAUTIONS

Operator Protection

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

Environmental Protection

Crops/situations with 5m buffer zone:

Since there is a risk to aquatic life from use, users not applying the statutory buffer zone must either themselves carry out or ensure that someone else has carried out a Local Environment Risk Assessment for Pesticides (LERAP) on their behalf before each spraying operation from a horizontal boom sprayer. Users must not allow direct spray from horizontal boom sprayers to fall within 5m of the top of the bank of any static or flowing waterbody or within 1m of a ditch which is dry at the time of application (these distances to be measured as set out in the guidance documents available from HSE Chemical Regulation Division's website and any amendments that are made to it) unless:

- (a) The LERAP indicates that a narrower buffer zone will be sufficient; and
- (b) Any measures indicated by the LERAP as justifying the narrower buffer zone are complied with in full and in accordance with any conditions applicable to them.

Spray must be aimed away from water.

The results of the LERAP must be recorded in written form and must be available for a period of three years for inspection to any person entitled to exercise enforcement powers under or in connection with the Plant Protection Products Regulations 2011 or the Plant Protection Products (Sustainable Use) Regulations 2012. (An electronic record will satisfy the requirement for a written record, providing it is similarly available for inspection and can be copied).

Detailed guidance on LERAPs and how to conduct a LERAP are contained in the guidance documents available from HSE Chemicals Regulation Division's website. All LERAPs must be carried out in accordance with this Guidance and any amendments that are made to it.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with HSE's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Storage and Disposal

Do not re-use container for any purpose.

KEEP AWAY FROM FOOD, DRINK, AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

INKA™ SX®

Herbicide

Contains Tribenuron-methyl. May produce an allergic reaction

May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects.

Do not breathe dust or spray.

Get medical advice/attention if you feel unwell.

Collect spillage.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To avoid risks to human health and the environment, comply with the instructions for use

**WARNING**
**IMPORTANT INFORMATION
FOR USE ONLY AS A PROFESSIONAL HERBICIDE**

Crops/ situations:	Maximum individual dose: (g product / ha)	Maximum number of treatments: (per crop)	Latest time of application:	Aquatic buffer zone distance (metres):
Barley, oats, rye, triticale, wheat	60	1	Up to and including flag leaf ligule/ collar just visible (GS39)	5 (see Environmental Protection Phrase 1)

Other Specific Restriction:

- (a) This product is not to be applied before end of February in the year of harvest.
- (b) This product must not be applied via hand-held equipment.
- (c) Non-returnable containers must not be re-used for any purpose.
- (d) This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with HSE's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.