

Lucento® Fungicide



WHEAT

Best-in-class protection on key wheat diseases while preserving yield potential.

Powered by a novel premix of active ingredients from FMC, new Lucento® fungicide provides an exclusive combination of benefits for smart disease protection and high yielding performance against the toughest diseases in wheat. Under heavy pressure, field studies demonstrated that Lucento fungicide provides excellent control of all labeled diseases. Lucento fungicide provides a plant health response and yield advantage over competitors. It can also be used to reduce the burden on other important fungicides as part of a resistance management plan.

Benefits of Lucento Fungicide:

- The only premix of a proven SDHI, bixafen, and FMC-patented flutriafol active ingredients. Highly systemic fungicide with preventative and curative control of leaf rust and powdery mildew.
- Both acropetal and translaminar movement improve coverage and disease control.
- Dual modes of action reduce risk of resistance.

Application Guidelines

Rate per acre: 3.0 - 5.5 fl. oz.

May be applied via ground, air or chemigation

Preharvest Interval (PHI): 21 days

Maximum applications per crop year: 2

Restricted Entry Interval (REI): 12 hours



Diseases Controlled:

- Leaf rust
(*Puccinia triticina*)
- Stripe rust
(*Puccinia striiformis*)
- Stagonospora leaf/ glume blotch
(*Parastagonospora nodorum*)
- Tan spot
(*Pyrenophora tritici repentis*)
- Septoria leaf blotch
(*Zymoseptoria tritici*)
- Stem rust
(*Puccinia graminis*)
- Powdery mildew
(*Blumeria graminis*)

Application Instructions:


Apply Lucento fungicide in a protective spray schedule or when conditions are favorable for disease development.

For optimum control of foliar diseases, a surfactant must be tank mixed with Lucento fungicide.

For management of fungal pathogen resistance development, do not make more than two sequential applications per year before alternating to another fungicide from a FRAC Group different from FRAC Groups 7 and 3.

Yield bu / Acre

Xianming Chen – Strip Rust Trial - 2018



Rotational Crop Guidelines

Lucento fungicide labeled crops	0 Days
Leafy crops	30 Days
Unlabeled crops	365 Days

Timing	Treatment	SY Ovation (Moderately Tolerant) Yield (Bu/A)		Brundage (Susceptible) Yield (Bu/A)	
Flag Leaf - Feekes 10	13.7 fl. oz. Trivapro® fungicide	134	a	95	a
Flag Leaf - Feekes 10	5.55 fl. oz. Lucento fungicide	134	a	93	ab
Flag Leaf - Feekes 10	5.5 fl. oz. Nexicor™ fungicide + 3 fl. oz. Tilt fungicide	129	bc	94	a
Flag Leaf - Feekes 10	4 oz. Priaxor Xemium fungicide + 4 fl. oz. Tilt fungicide	130	bc	92	ab
Flag Leaf - Feekes 10	14 fl. oz. Quilt® XL fungicide	128	bc	91	ab
Flag Leaf - Feekes 10	7 fl. oz. Alto® fungicide	126	c	90	b
Flag Leaf - Feekes 10	4 oz. Tilt fungicide	122	d	90	b

Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. Lucento® fungicide may not be registered for sale or use in all states. Contact your local FMC retailer or representative for details and availability in your state. FMC, the FMC logo and Lucento are trademarks of FMC Corporation or an affiliate. Priaxor, Xemium and Nexicor are trademarks of BASF. Tilt, Trivapro, Quilt and Alto are trademarks of Syngenta Group Company. Zolera is a trademark of Arysta Life Science. ©2019 FMC Corporation. All rights reserved. 19-FMC-0074 01/19



An Agricultural Sciences Company