
Brigade® 2EC Insecticide/Miticide



Product Information:

Scouting for and identifying mid- to late-season cotton pests is an important step to manage boll damage and harvest quality. Because eggs and larvae are often found underneath the canopy, the level of insect pressure growers may be facing is deceiving. Brigade® 2EC insecticide/miticide provides excellent results controlling chewing, sucking and piercing pests that impact cotton.

Controlling these pests can help avoid substantial losses. Stink bugs, for example, can lead to economic losses from reduced yields and diminished fiber quality. The broad-spectrum formula in Brigade 2EC insecticide/miticide reduces stink bugs, plant bugs, bollworms as well as mites. Plus, its all-weather chemistry reduces wash-off and increases residual activity for prolonged control. This insecticide/miticide can be applied via ground or air and is a flexible tank-mix partner.

Quick Facts:

- Excellent control of chewing, sucking and piercing pests including stink bugs, plant bugs, bollworms and mites.
- Reduced wash-off due to all-weather chemistry that increases residual control.
- Flexible tank-mix partner.
- · Can be applied by ground or air.
- Low water solubility.



Label Information:

Active Ingredient: Bifenthrin REI: 12 hours
Group 3A Insecticide PHI: 14 days

Signal Word: WARNING

Cotton Application Recommendations

PEST	RATE OF APPLICATION	
	LB. AI/A	FL. OZ./A
European Corn Borer Soybean (Banded) Thrips Tobacco Thrips	0.02 – 0.10	1.3 – 6.4
Boll Weevil Bollworm Cabbage Looper Cotton Aphid Cotton Fleahopper Cotton Leafperforator Cutworms Fall Armyworm Plant Bug Saltmarsh Caterpillar Southern Garden Leafhopper Stink Bugs Tobacco Budworm Whitefly Yellow Striped Armyworm	0.04 – 0.10	2.6 – 6.4
Beet Armyworm Carmine Spider Mite Lygus Spp. Pink Bollworm Twospotted Spider Mite	0.06 – 0.10	3.8 – 6.4

For more information about Brigade® 2EC insecticide/miticide, contact your FMC retailer or visit FMCCrop.com.



