## **Affinity** BroadSpec herbicide with TotalSol® soluble granules **Affinity** TankMix herbicide with TotalSol® soluble granules



#### **Product Information:**

Affinity® BroadSpec herbicide with TotalSol® soluble granules and Affinity® TankMix herbicide with TotalSol® soluble granules offer a flexible, reliable solution for postemergence broadleaf weed control in cereal crops. Affinity Broadspec herbicide may be safely tank mixed with all Group 2 grass herbicides without compromising weed control. Affinity TankMix herbicide is available to be safely mixed with Group 1 grass herbicides in small grains.

Affinity brand herbicides complement other broadleaf herbicides to offer a more effective and complete broad-spectrum weed control solution. Affinity brand herbicides are designed to provide maximum flexibility and ease of use. Both herbicides are formulated with TotalSol soluble granules, a patented formulation that dissolves completely for more consistent weed control and simplified sprayer clean-out. Affinity brand herbicides also allow for crop rotation to most crops after harvest.

For maximum flexibility and cleaner fields, choose an Affinity brand herbicide as a tank-mix partner for your cereal fields.

### **Quick Facts**

- · Labeled for use on wheat, durum and barley.
- Controls key broadleaf weeds, including wild buckwheat and volunteer canola.
- For best results, use Affinity brand herbicides on newly emerged, actively growing weeds.





# Weed list combines Affinity® TankMix and Affinity® BroadSpec herbicides, each with TotalSol® soluble granules. See labels for a full list of weeds controlled specific to each product.

Common cocklebur\*\* Flixweed
Common lambsquarters Marshelder
Common sunflower\*\* Henbit

Kochia\*\*\* Annual sowthistle
Redroot pigweed\*\*\* Field pennycress
False chamomile/Mayweed Wild mustard

Volunteer peas Nightflowering catchfly

Volunteer flax\*\*

Volunteer lentils

Volunteer canola

Marestail\*\*

Curly dock

Prickly lettuce

Lady's thumb

Lanceleaf sage\*

Common mallow\*\*
Canada thistle\*\*
Wild buckwheat

#### **Application Guidelines:**

- Re-Entry Interval: 12 hours
- Pre-Harvest Interval:
   Wheat, barley, triticale: 45 days
- Product Use Rates:
   Aim® EC herbicide: 0.5 1.0 fl oz./A
   Affinity BroadSpec herbicide: 0.4 1.0 oz./A

Affinity TankMix herbicide: 0.6 - 1.0 oz./A

- Use only clean water for preparing product solutions.
- Any tank-mix partners and adjuvants should be added to the spray tank only after the Affinity brand product granules have fully disolved.

#### **Application Guidelines:**

Wheat, barley and triticale: Make application after crop is in the two-leaf stage, but before flag leaf is visible.

#### **Spray Adjuvants:**

Under arid/semi-arid conditions, non-ionic surfactant or crop oil may be used. See label for specific recommendations.

#### Label:

Active Ingredient:

Thifensulfuron methyl and Tribenuron methyl.

Mode of Action:

Group 2 ALS/AHAS inhibitor. Blocks normal function of the enzyme acetolactate acid (ALS) and acetohydroxy acid synthase (AHAS).

For more information, contact your local FMC retailer or representative to learn more about Affinity BroadSpec and Affinity TankMix herbicides from FMC and visit FMCCrop.com.

Always read and follow all label directions, restrictions and precautions for use. Some products may not be registered for sale or use in all states. As of November 1, 2017, the USEPA registrations for DuPont™ Affinity® BroadSpec herbicide with TotalSol® soluble granules and Affinity® TankMix herbicide with TotalSol® soluble granules and TotalSol® soluble granules were sold to FMC by DuPont. FMC, Aim, Affinity and TotalSol are trademarks of FMC Corporation or an affiliate. ©2018 FMC Corporation. All rights reserved. 18-FMC-2102 10/18







<sup>\*</sup> See specific weed instructions within the label for more information.

<sup>\*\*</sup>Partial control/suppression.

<sup>\*\*\*</sup>Naturally occurring resistant biotypes are known to occur.