

# ALLY® EXTRA SG HERBICIDE WITH TOTALSOL® SOLUBLE GRANULES

## RELIABLE CONTROL OF KEY BROADLEAF WEEDS FOR A STRONG START

### ENHANCED CONTROL OF BROADLEAF WEEDS IN PROGRAMS WITH GROUP 1 GRASS HERBICIDES.

PRODUCT ATTRIBUTES	BENEFITS
Flexibility	<ul style="list-style-type: none"> <li>Can be applied with topdress fertilizer to control weeds and contribute to a cleaner harvest.</li> <li>Ally® Extra SG herbicide with TotalSol® soluble granules can be used as an early postemergence or topdress herbicide application in the spring, summer or fall when a majority of weeds are actively growing.</li> </ul>
Crop safety	Offers excellent crop tolerance to help maximize yields.
TotalSol soluble granule formulation	TotalSol soluble granules dissolve completely in water, providing more consistent weed control with faster, easier spray tank clean-out.
Three active ingredients	Delivers control against many tough broadleaf weeds. See back side for complete weed list.

### RATE RECOMMENDATION

#### For most varieties of wheat, barley, triticale and fallow

- Apply Ally Extra SG herbicide at 0.3-0.5 oz./A.
- Apply to wheat and barley after the crop is in the two-leaf stage, but before the flag leaf is visible except for durum wheat and wampum varieties.

#### For durum wheat and wampum spring wheat

- Apply Ally Extra SG herbicide at 0.3-0.5 oz./A.
- Apply after the crop is tillering but before boot. Always apply in combination with 2,4-D.

### APPLICATION INFORMATION

- Tank mix with MCP ester or dicamba.
- Dissolve in water prior to adding MCPA ester and add the surfactant last.

### TECHNICAL INFORMATION

Active Ingredient(s)	Thifensulfuron methyl, Tribenuron methyl, Metsulfuron methyl
IRAC Class / MOA	Group 2 ALS/AHAS inhibitor. Blocks normal function of the enzyme acetolactate acid (ALS) and acetohydroxy acid synthase (AHAS).
Formulation	TotalSol soluble granules
Signal Word	CAUTION
Packaging	12 x 20 oz. or 480 oz. refillable container

### USAGE GUIDELINES

PHI	45 days
REI	12 hours

# ALLY® EXTRA SG HERBICIDE WITH TOTALSOL® SOLUBLE GRANULES

TECHNICAL BULLETIN | **WHEAT**

Group 2 Herbicide

## RELIABLE CONTROL OF KEY BROADLEAF WEEDS FOR A STRONG START

WEEDS CONTROLLED		
<ul style="list-style-type: none"> <li>• Annual knawel</li> <li>• Annual sowthistle</li> <li>• Black mustard</li> <li>• Blue/purple mustard*</li> <li>• Broadleaf dock</li> <li>• Bur buttercup (testiculate)</li> <li>• Bushy wallflower/Treacle mustard</li> <li>• Canada thistle*</li> <li>• Carolina geranium</li> <li>• Claspig pepperweed</li> <li>• Coast fiddleneck (tarweed)</li> <li>• Common buckwheat</li> <li>• Common chickweed</li> <li>• Common cocklebur</li> <li>• Common mallow</li> <li>• Common purslane</li> <li>• Common radish</li> <li>• Common ragweed</li> <li>• Common sunflower*</li> <li>• Conical catchfly</li> <li>• Corn chamomile</li> <li>• Corn gromwell</li> <li>• Corn spurry</li> <li>• Cowcockle</li> <li>• Cress (mouse-ear)</li> </ul>	<ul style="list-style-type: none"> <li>• Curly dock</li> <li>• Cutleaf evening primrose</li> <li>• False chamomile</li> <li>• Field chickweed</li> <li>• Field pennycress (fanweed)</li> <li>• Filaree (redstem, Texas)</li> <li>• Flixweed*</li> <li>• Groundsel (common)</li> <li>• Henbit</li> <li>• Kochia‡</li> <li>• Knotweed (prostrate)*</li> <li>• Lambsquarter (common, slimleaf)</li> <li>• London rocket</li> <li>• Marshelder</li> <li>• Mayweed chamomile</li> <li>• Miner's lettuce</li> <li>• Narrowleaf lambsquarter</li> <li>• Nightflowering catchfly</li> <li>• Pennsylvania smartweed</li> <li>• Pigweed (prostrate, redroot, smooth, tumble)</li> <li>• Pineappleweed</li> <li>• Plains coreopsis</li> <li>• Prickly lettuce‡</li> <li>• Redmaids</li> </ul>	<ul style="list-style-type: none"> <li>• Russian thistle‡</li> <li>• Scentless chamomile/mayweed</li> <li>• Shepherd's purse</li> <li>• Smallflower buttercup</li> <li>• Smallseed falseflax</li> <li>• Smartweed (green, lady's thumb, pale)</li> <li>• Snow speedwell</li> <li>• Sticky chickweed</li> <li>• Stinking mayweed/dog fennel</li> <li>• Swinecress</li> <li>• Tansy mustard*</li> <li>• Tarweed fiddleneck</li> <li>• Tumble/Jim Hill mustard</li> <li>• Volunteer lentils</li> <li>• Volunteer peas</li> <li>• Volunteer sunflower</li> <li>• Waterpod</li> <li>• Wild buckwheat*</li> <li>• Wild chamomile</li> <li>• Wild garlic*</li> <li>• Wild mustard</li> <li>• Wild radish*</li> </ul>
WEEDS PARTIALLY CONTROLLED**		
<ul style="list-style-type: none"> <li>• Catchweed bedstraw</li> <li>• Mallow (little)</li> </ul>	<ul style="list-style-type: none"> <li>• Nightshade (cutleaf, hairy)</li> <li>• Sowthistle (annual)*</li> </ul>	<ul style="list-style-type: none"> <li>• Tall waterhemp</li> <li>• Vetch (common, hairy)*</li> </ul>

\* See the Specific Weed Problems section of the Ally® Extra SG herbicide with TotalSol® soluble granules label for more information.

\*\* Partial control: For better results, use the highest labeled rate of Ally Extra SG herbicide and include a tank-mix partner such as 2,4-D, MCPA, bromoxynil (such as Buctril® herbicide, Bison® herbicide, Bronate® or Bronate® Advanced herbicides) or dicamba (such as Banvel® herbicide/ Clarity® herbicide). Refer to the Tank Mixtures section of the Ally Extra SG herbicide label.

‡ Naturally occurring resistant biotypes of kochia, prickly lettuce and Russian thistle are known to occur. See the Tank Mixtures and Specific Weed Problems sections of the Ally Extra SG herbicide label for additional details.