

Anthem® Flex Herbicide

**Flexible
Powerful
Accurate
Against Resistance**

WHEAT

PRODUCT INFORMATION:

Anthem® Flex herbicide from FMC provides residual control of grasses and broadleaf weeds in winter and spring wheat. This low-use-rate herbicide has a wide application window and comes in an easy-to-mix formulation.

Anthem Flex herbicide delivers long residual control and suppression of broadleaves, including downy brome*, Italian ryegrass, rattail fescue, yellow foxtail, Palmer amaranth and other tough weeds.

With more flexibility and no known resistance issues, Anthem Flex herbicide is an excellent weed management tool. Anthem Flex herbicide is even effective at controlling weeds with known resistance to glyphosate and ALS herbicides.

- Labeled for spring and winter wheat
- Applied in all tillage systems
- Preplant, preemergence and early postemergence applications
- Low use rate:
 - Spring/winter wheat:** 2.0 – 4.5 fl. oz./A
- Control and suppression of key annual grasses and broadleaf weeds
- No known resistance



Target Weeds:

Annual bluegrass
Chickweed
Crabgrass
Downy brome*
Italian ryegrass
Palmer amaranth
Rattail fescue
Pigweed
Yellow foxtail
Waterhemp

Application Timing:

****Spring and winter wheat, except in Washington, Oregon, Idaho and Montana:** Delayed preemergence and early postemergence

****Spring and winter wheat in Washington, Oregon, Idaho and Montana:** Preplant surface, preemergence, early postemergence

Product Use Rate:

Winter/spring wheat: 2.0 – 4.5 fl. oz./A

Restrictions:

Spring/winter wheat: Do not apply more than 0.133 lb. ai/A of pyroxasulfone and 0.0095 lb. ai/A of carfentrazone-ethyl in a 12-month cropping season.

Please see the label for a complete list of restrictions.

Label:

Active Ingredients: Pyroxasulfone (Group 15);
carfentrazone-ethyl (Group 14)

Mode of Action: Pyroxasulfone inhibits very-long-chain fatty acid synthesis as a Group 15 herbicide. It is a root and shoot growth inhibitor that controls susceptible germinating seedlings before or soon after they emerge from the soil.


Spring and Winter Wheat

RATE SELECTION/SOIL TEXTURE

WHEAT (SPRING AND WINTER) GROWN IN WASHINGTON, OREGON, IDAHO AND MONTANA CROP SECTION			
APPLICATION RATES FOR ANTHEM® FLEX HERBICIDE WHEN APPLIED ALONE, IN TANK MIX, OR SEQUENTIALLY			
Application Timing	Use rate (oz./A) by soil texture ¹ Anthem Flex herbicide fl. oz./A (lb. ai./A)		
	Coarse	Medium	Fine
Preplant Surface	2.5 – 2.7 (0.078 – 0.085)	2.75 – 3.5 (0.086 – 0.102)	3.5 – 4.5 (0.109 – 0.141)
Delayed preemergence	2.0 – 2.7 (0.063 – 0.085)	2.75 – 3.5 (0.086 – 0.102)	3.5 – 4.5 (0.109 – 0.141)
Early postemergence	2.0 – 2.7 (0.063 – 0.085)	2.75 – 3.6 (0.086 – 0.114)	2.7 – 4.5 (0.085 – 0.141)

Anthem Flex Herbicide Delayed Preemergence in Wheat

Charlie Cahoon, Virginia Tech, Painter, VA 2016



To learn more about Anthem Flex herbicide, contact your FMC Star Retailer or visit FMCcrop.com.

*Please see the Anthem Flex herbicide FIFRA SECTION 2(ee) recommendation for control of downy brome in Montana, Idaho, Oregon and Washington.

**Please see the Anthem Flex herbicide 24c SLN labels for a list of additional states that can apply preplant or preemergence in wheat.

Always read and follow all label directions, restrictions and precautions for use. Some products may not be registered for sale or use in all states. FMC and Anthem are trademarks of FMC Corporation or an affiliate. Axiom and Osprey are trademarks of Bayer CropScience. ©2018 FMC Corporation. All rights reserved. 18-FMC-0089 03/18

