Group 14, 15 Herbicide

FOR SUPERIOR CONTROL OF GRASSES AND SMALL-SEEDED BROADLEAF WEEDS IN SOYBEANS.

- Excellent foundational herbicide. Unmatched residual control of small-seeded broadleaf weeds and grasses.
- For use in LibertyLink®, Roundup Ready® and Roundup Ready 2 Xtend®* trait systems, Enlist®* soybean systems, as well as conventional soybeans.
- Controls waterhemp, Palmer amaranth, nightshade, lambsquarters, kochia, morningglories and smartweed.
- Provides excellent residual control of grasses, up to two weeks longer than competitors, to allow for reduced postemerge trips in season.
- For resistant weed populations, use higher recommended rate for longer residual control.
- For no-till: Add Aim® EC herbicide 0.75-1 fl. oz. + 2,4-D 1 pt./A + COC 1% or appropriate burndown program.

'Please see individual tank mix product labels for XtendiMax, Engenia, FeXapan and Enlist brands for approved tank mix use.

USE DIRECTIONS			
Rates - Soybeans	5.9-15.7 fl. oz./A		
Application Timing	Apply preplant, preemergence up to three days after planting.		
Tank-Mix Recommendations- Refer to labels for rate selection by soil type	 Metribuzin – 5-7 oz./A FirstRate® herbicide 0.4-0.6 oz./A Classic® herbicide 1-1.5 oz./A 		
Restrictions	Do not make more than two applications per year. Do not apply within seven days if a second application is necessary. On coarse soils - DO NOT apply more than a maximum cumulative amount of 9.42 fl. oz/A Authority® Edge herbicide (0.313 lb. ai/A containing 0.111 lb. ai of pyroxasulfone and 0.202 lb. ai of sulfentrazone) per year. On medium and fine soils – DO NOT apply more than a maximum cumulative amount of 15.67 fl. oz/A of Authority Edge herbicide (0.521 lb. ai/A containing 0.186 lb. ai of pyroxasulfone and 0.335 lb. ai of sulfentrazone) per year.		

ORGANIC MATTER %	COARSE	MEDIUM	FINE	
PRE USE RATES - OUNCES / ACRE				
< 1.5	5.9-7.9	7.0-10.0	9.0-12.0	
1.5 - 3.0	6.9-8.8	8.0-11.0	10.0-13.0	
> 3.0	7.0-9.4	9.0-12.0	12.0-15.7	

Use higher rates for soils of pH less than 7.2 and lower rates for pH greater than 7.2 within the rate range.

TECHNICAL INFORMATION			
ACTIVE INGREDIENT	Sulfentrazone + Pyroxasulfone		
LOADING	4.25 lbs. ai/gal.		
WSSA GROUP	14,15		
FORMULATION	SC - Suspension Concentrate		
SIGNAL WORD	Caution		

ROTATIONAL CROP GUIDELINES**		
CORN (FIELD AND SEED)	4 months	
WHEAT	4 months	
PEANUTS, POTATOES	4 months	
SUNFLOWERS	Anytime	
CORN (SWEET AND POP)	10-12 months	
COTTON	12-18 months	
SUGAR BEETS	24 months	

** Rotational crop guidelines based on up to 13.4 fl. oz./A. See label for specific rotational intervals.





