# **Command<sup>®</sup> 3ME** Microencapsulated Herbicide

Keep competitive grasses and broadleaf weeds out of your tobacco fields with advanced herbicide technology.

#### TOBACCO

#### **Product Information:**

Command<sup>®</sup> 3ME microencapsulated herbicide provides excellent weed control in a convenient package. Apply to soil at preemergence before or after transplant for control of problem grasses like foxtails and key broadleaf weeds like lambsquarters and smartweed. An advanced microencapsulation formula protects the integrity of the herbicide in tough environmental conditions for maximum protection. Likewise, its favorable environmental profile protects the integrity of your land.

Command 3ME microencapsulated herbicide can be tank mixed with other herbicides for broader weed control and is compatible with conventional tillage, reduced tillage and no-till practices.

#### **Quick Facts:**

- Advanced microencapsulation technology protects the integrity of the herbicide so you can protect the integrity of your tobacco harvest.
- Formulation reduces off-target effects and volatile movement.
- Strong control of key broadleaf and annual grass species.
- Convenient for tank mixing and compatible with any tilling practices.
- Can be applied preemerge before or after transplant.



# Weeds Controlled / Partially Controlled\*:

Barnyardgrass	Jimsonweed*
Bermudagrass	Panicum, Fall and Texas
Broadleaf Signalgrass	Velvetleaf
Cocklebur*	Lambsquarters
Common Ragweed*	Purslane
Crabgrass	Prickly Sida
Field Sandbur	Smartweed, Ladysthumb
Foxtail, Giant, Green	and Pennsylvania
and Yellow	Spurred Anoda
Goosegrass	

\*The weeds listed are controlled or partially controlled at varying rates, see label for complete listing of rates of control depending on weed spectrum.

# **Application Timing:**

**Preemergent (Prior to Transplant):** Make a single application in a minimum of 20 gallons of water per acre at a rate of 2.0-2.667 pints (0.750-1.0 lb. active) per acre. For heavy pressure or heavy soils, use the higher specified rate, otherwise less than desirable weed control may result.

Apply immediately before or up to 30 days prior to transplanting. Apply as a surface treatment prior to emergence of weeds. If weeds emerge before preemergence application can be made, cultivate before or immediately preceding treatment.

If additional seedbed preparation is needed, use equipment to move the soil no deeper than 1-1/2-2 inches.

**Preemergent (Post Transplant):** Make a single broadcast application in a minimum of 20 gallons of water per acre at the rate of 2.0-2.667 pints (0.75 -1.0 lb. active) per acre. For heavy weed pressure or heavy soils, use the higher specified rate, otherwise less than desirable weed control may result.

Apply no more than once per year using a maximum rate of 2.667 pints (1.0 lb. ai./A) per acre. Apply over the top of tobacco plants immediately or up to seven days after transplanting but prior to emergence of weeds. If weeds emerge before application, cultivate before treatment

### Product Use Rate:

Typical Use Rate: 2.0 pt. / A REI: 12 hours

#### Label:

#### Active Ingredient: Clomazone (Group 13)

**Mode of Action:** Clomazone is an isoxazolidinone herbicide, which acts to inhibit pigment synthesis in the plant chloroplast as a Group 13/Group F3 herbicide. It is quickly taken up by the roots and shoots to block plant chlorophyll to control susceptible germinating seedlings before or soon after they emerge from the soil.

# **Restrictions:**

Do not graze or harvest for food or feed cover crops planted less than nine months after Command<sup>®</sup> 3ME microencapsulated herbicide application. Do not use on tobacco seedling beds. Do not apply more than 1.0 lb. ai. per acre per year.

Do not retreat field with a second application of Command 3ME microencapsulated herbicide.

# Please see the label for a complete list of restrictions and bufferzone requirements.

### FMC Tobacco Portfolio:

The complete tobacco portfolio line of crop protection products from FMC includes the following.

#### Herbicides:

Aim® EC herbicide Command 3ME microencapsulated herbicide Spartan® 4F herbicide Spartan® Charge herbicide **Insecticides:** Capture® LFR® insecticide Coragen® insect control Exirel® insect control

Verimark® insect control

# For more information about Command 3ME microencapsulated herbicide and the complete tobacco portfolio, contact your FMC retailer or visit FMCcrop.com.

Capture LFR insecticide is a Restricted Use Pesticide. Always read and follow all label directions and precautions for use. Some products may not be registered for sale or use in all states. As of November 1, 2017, the USEPA registrations for DuPont<sup>™</sup> Coragen<sup>®</sup>, Exirel<sup>®</sup> and Verimark<sup>®</sup> insect controls were sold by to FMC by DuPont. FMC, Aim, Capture, LFR, Command, Coragen, Exirel, Spartan and Verimark are trademarks of FMC Corporation or an affiliate. ©2018 FMC Corporation. All rights reserved. 18-FMC-1045 07/18



