

PRODUCT ATTRIBUTES	BENEFITS
<b>One-Pass Weed Control</b>	Finesse® cereal and fallow herbicide is compatible with liquid fertilizers, making it easy and economical to achieve weed control after only one application.
<b>Contact and Residual Activity</b>	Ability to burn down weeds that are emerged at the time of application. One application provides residual control of winter and summer weeds through harvest.
<b>Superior Broadleaf Weed Control</b>	University studies prove Finesse cereal and fallow herbicide delivers excellent control of winter annual weeds such as henbit and mustards. It also provides residual activity to keep spring-germinating weeds like wild buckwheat under control.
<b>Application Flexibility</b>	<ul style="list-style-type: none"> <li>Finesse cereal and fallow herbicide can be applied in the fall, winter or spring, giving growers more control over the timing of their fertilizer application.</li> <li>One fall treatment of Finesse cereal and fallow herbicide at a 0.4 oz./A rate will control winter weeds like mustards and henbit while still providing control of wild buckwheat and other weeds, saving growers the extra step of a spring treatment.</li> </ul>
<b>Excellent Crop Safety</b>	Finesse cereal and fallow herbicide has excellent crop tolerance and no grazing restrictions.

WEEDS CONTROLLED OR SUPPRESSED			
0.2-0.3 oz./A		0.3-0.4 oz./A	0.5 oz./A (preemergence only)
Beakchervil, bur	Groundsel	Bedstraw <sup>2,3</sup>	Bromus species: cheat <sup>1,2,3</sup> , downy brome <sup>1,2,3</sup> , Japanese brome <sup>1,2,3</sup>
Bushy wallflower (treacle mustard)	Hempnettle	Bluegrass, annual <sup>2,3</sup>	Corn, volunteer <sup>3</sup>
Buttercup, bur (testiculate)	Henbit	Bromus species (cheat, downy brome, Japanese brome) <sup>1,2,3</sup>	Ryegrass, annual <sup>1,2,3</sup>
Carrot, wild	Lambsquarters <sup>1</sup>	Buckwheat, wild <sup>3,4</sup>	
Catchfly, conical	Lettuce, prickly <sup>1,3</sup>	Corn, volunteer <sup>3</sup>	
Chamomile: false, mayweed	Miner's lettuce	Darnel, Persian <sup>2,3</sup>	
Chickweed: common <sup>1</sup> , jagged, mouse ear	Mustard: blue, tumble (Jim Hill), wild <sup>1</sup>	Fiddleneck, coast (tarweed)	
Cockle, white	Pennycress, field	Foxtail: green (pigeon grass) <sup>1,2</sup> , yellow <sup>1,2,3</sup>	
Coreopsis, plains	Pepperweed, Virginia	Geranium, dovefoot	
Cow cockle	Pigweed: prostrate, redroot <sup>1</sup> , smooth <sup>1</sup>	Gromwell, corn <sup>2,3</sup>	
Dock: broadleaf, curly	Pineappleweed	Knotweed, prostrate <sup>2,3</sup>	
Evening primrose, cutleaf	Purslane	Kochia <sup>1,2,3</sup>	
False flax, small seed <sup>1</sup>	Shepherd's purse	Poppy, prickly (pinnate)	
Filaree, redstem	Ladysthumb	Radish, wild <sup>3</sup>	
Flixweed <sup>1,2,3</sup>	Spurry, corn	Ryegrass, annual <sup>1,2,3</sup>	
Geranium, Carolina	Tansy mustard <sup>2,3</sup>	Smartweed, Pennsylvania <sup>2</sup>	
		Sowthistle, annual	
		Speedwell: common <sup>2</sup> , ivyleaf <sup>2</sup>	
		Sunflower <sup>1,3</sup>	
		Thistle: Canada <sup>2,3</sup> , Russian <sup>1,2,3</sup>	
		Vetch <sup>3</sup>	

<sup>1</sup>Naturally occurring resistant biotypes of these weeds are known to occur. See the Tank Mixtures, Specific Weed Instructions and Weed Resistance sections of the Finesse cereal and fallow herbicide label for additional details.

<sup>2</sup>When used as directed, weeds are suppressed and/or controlled. Weed suppression is visible reduction in weed competition, reduced population and vigor as compared to an untreated area. Degree of suppression will vary with rate used, size of weeds and environmental conditions following treatment.

<sup>3</sup>See the Specific Weed Problems section on the Finesse cereal and fallow herbicide label for more information regarding control and suppression of these weeds.

<sup>4</sup>For best results, apply Finesse cereal and fallow herbicide preemergence at 0.4 fl. oz./A to wild buckwheat in the fall or early spring.

### APPLICATION RECOMMENDATIONS

- For best results, apply Finesse® cereal and fallow herbicide at 0.2-0.4 oz./A.
- Apply preemergence or anytime from the two-leaf stage to boot.
- Unless otherwise directed, always add a nonionic surfactant to the spray solution. Do not use low rates of liquid nitrogen fertilizer as a substitute for a surfactant.

### APPLICATION GUIDELINES

When applying Finesse cereal and fallow herbicide with liquid fertilizer, it is advisable to slurry it in water, then thoroughly mix the slurry into the liquid fertilizer. For best results on emerged weeds, Finesse cereal and fallow herbicide should always be tank mixed with MCPA, 2,4-D, dicamba or fluroxypyr.

### TECHNICAL INFORMATION

Active Ingredient(s)	Group	Formulation	Signal Word	PPE	PHI	REI	Packaging
Chlorsulfuron, Metsulfuron Methyl	Group 2 ALS/AHAS inhibitor. Blocks normal function of the enzyme acetolactate acid (ALS) and acetohydroxy acid synthase (AHAS).	White and brown granules	CAUTION	Mixers, loaders, applicators and other handlers: long-sleeved shirt and long pants, shoes plus socks and chemical resistant gloves.	45 days (grain)	4 hours	12 x 20 fl. oz. or 480 fl. oz. refillable container.